

University of Arkansas, Fayetteville

ScholarWorks@UARK

---

Theses and Dissertations

---

5-2021

## Impact of Different Plant-Based Foods Consumption on the Composition and Diversity of Gut Microbiota

Ayoub Al Othaim

*University of Arkansas, Fayetteville*

Follow this and additional works at: <https://scholarworks.uark.edu/etd>



Part of the [Food Microbiology Commons](#), and the [Human and Clinical Nutrition Commons](#)

---

### Citation

Al Othaim, A. (2021). Impact of Different Plant-Based Foods Consumption on the Composition and Diversity of Gut Microbiota. *Theses and Dissertations* Retrieved from <https://scholarworks.uark.edu/etd/3992>

This Dissertation is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of ScholarWorks@UARK. For more information, please contact [ccmiddle@uark.edu](mailto:ccmiddle@uark.edu).

Impact of Different Plant-Based Foods Consumption on the Composition and Diversity of Gut  
Microbiota

A dissertation submitted in partial fulfillment  
of the requirements for the degree of  
Doctor of Philosophy in Cell and Molecular Biology

by

Ayoub Al Othaim  
Qassim University  
Bachelor of Medical Microbiology, 2009  
Quinnipiac University  
Master of Health Science, Medical Laboratory Sciences, 2014

May 2021  
University of Arkansas

This dissertation is approved for recommendation to the Graduate Council.

---

Franck Carbonero, Ph.D.  
Dissertation Director

---

Luke R. Howard, Ph.D.  
Committee Member

---

Jiangchao Zhao, Ph.D.  
Committee Member

---

Charles Rosenkrans, Ph.D.  
Committee Member

## ABSTRACT

Plant-based diets are primarily derived from plant sources, including fruits and vegetables. Such patterns are associated with health benefits by promoting general health and preventing many diseases and metabolic disorders. Fruits and vegetables are key sources of high-quality nutrients such as dietary fibers and polyphenols. Since they are rich in dietary fibers and polyphenols, whole fruits and vegetables represent unique and somewhat understudied modulators of the gut microbiota and its associated functions. This study aims to (i) investigate the impact of varying dilutions of cherry juices concentrate on the composition of the murine gut microbiota, (ii) investigate the impact of cranberry juice consumption on the bacterial composition of gut and vaginal microbiota in postmenopausal women, and (iii) investigate the impact of the amounts and botanical diversity of fruits and vegetables on gut microbiota profiles in healthy adults. Fecal samples were collected from mice consuming increasing tart cherry juice concentrations in their drinking water. A randomized, crossover controlled dietary intervention was conducted with cranberry juice and a matched control, and postmenopausal female volunteers provided stool and vaginal swabs for microbiota analyses. Two pilot studies looking at the impact of low and adequate intake of fruits and vegetables (SMART-Diet), as well as low versus high botanical diversity (SMART-Valid), were conducted at Colorado State University; stool samples were collected for gut microbiota analyses and short-chain fatty acids measurements. Microbial analyses were performed using Illumina following a dual-indexing 16S rRNA gene amplicon. Each dietary intervention resulted in distinct microbiota modulations and effects. Cherry juices consumption significantly increased the relative abundance of *Akkermansia* and *Barnesiella* and decreased the relative abundance of *Bacteroides*. Consumption of cranberry juice modulated the gut microbiota composition by elevating the relative abundance

of *Bifidobacterium*, *Clostridium* XIVa members, *Eggerthella*, and *Prevotella*. Cranberry juice consumption also resulted in a beneficial impact on vaginal microbiota by decreasing the relative abundance of *Streptococcus* compared to a placebo. Finally, the SMART-Diet study showed that fruit and vegetable consumption resulted in a reduction of the Firmicutes/Bacteroidetes ratio whereas the SMART-Valid study showed an increase in the Firmicutes/Bacteroidetes ratio after fruit and vegetable consumption, and a significant increase in production of beneficial short-chain fatty acids. Our findings suggest that consumption of cherry juices, cranberry juice, and fruit and vegetable inclusive diets all have a beneficial impact on the composition of gut microbiota. This body of work is consistent with the emerging concept that whole foods should be considered rather than extracts when it comes to gut microbiota and microbiome studies. Human dietary studies with larger sample size are warranted by these suggested beneficial modulations.

## **ACKNOWLEDGMENTS**

First and foremost, praise and thanks be to God, the Almighty, the greatest of all, for the blessings He has bestowed upon me and for giving me the strength and wisdom to achieve this dream. Without His grace, this research would not have been possible.

This doctoral thesis exists in its current form due to the assistance and encouragement of several people. With immeasurable love and appreciation, I would like to express my deepest gratitude to the people who have had the greatest impact on my success during my Ph.D. journey. First, I would like to extend my sincere gratitude to my supervisor, Dr. Franck Carbonero, who expertly guided me through this study. I gratefully acknowledge his motivation, encouragement, invaluable insights, constructive suggestions and feedback, and continuous support throughout this work. I would also like to extend my gratitude to my thesis committee, Dr. Luke Howard, Dr. Charles Rosenkrans, and Dr. Jiangchao Zhao, for their constructive suggestions, insightful comments, and wonderful contributions and support; to Dr. Tiffany Weir for the opportunity to use her lab at Colorado State University and for providing the necessary equipment; and to my laboratory colleagues, who have generously supported me with their skill. Lastly, I extend my special thanks and appreciation to my sponsor, Majmaah University, for supporting me throughout my graduate studies.

## **DEDICATION**

This study is dedicated to my beloved parents, Abdulaziz Al Othaim and Helah Al Othaim, who have been my source of inspiration and have given me endless love and encouragement to complete my studies. My dream of completing this research would not have come true without their support, sacrifice, and prayers. I also dedicate this work to my wife, soulmate, and best friend, Dr. Sarah Alajaji, with my heartfelt thanks for her mental strength, support, encouragement, and patience; and to my dear little boy, Feras, the most precious gift in my life, to whom I express my deep love. Though I did not deliberately neglect him, he has missed out on some of my love, care, and affection during this busy period. I also extend my gratitude to my siblings, Huda, Wafa, Sumayyah, Mohammed, Ibraheem, Younes, Yousef, Motaz, Adeeb, and Jory, whose continued support has motivated me to pursue my education and complete this accomplishment. Last but not least, I express my sincere thanks to my parents-in-law, Mansour and Zainab, for their blessings, affection, and support during my studies.

## TABLE OF CONTENT

GENERAL INTRODUCTION.....	1
CHAPTER 1: .....	3
Impact of Fruit and Vegetable Inclusive Diets on Gut Microbiota Composition and Diversity ....	3
Abstract .....	3
Keywords: .....	4
1.1 Introduction .....	4
1.2 Plant-based dietary patterns .....	5
1.2.1 Health benefits of plant-based diets .....	6
1.2.1.1 Lower risk of obesity .....	6
1.2.1.2 Lower risk of diabetes.....	8
1.2.1.3 Lower risk of cardiovascular diseases .....	9
1.2.1.4 Lower risk of cancers.....	10
1.3 Gut Microbiota Modulation by Plant-Based Foods .....	11
1.3.1 Gut Microbiota: General Aspects .....	11
1.3.2 Effect of Plant-Based Diets on Gut Microbiota Composition.....	15
1.3.3 Impact of Fruits and Vegetables on Gut Microbiota Composition .....	18
1.3.3.1 Berries .....	18
1.3.3.2 Cherries .....	26
1.3.3.3 Mangoes .....	28

1.3.3.4 Apples .....	29
1.3.3.5 Grapes .....	30
1.3.3.6 Kiwifruits .....	33
1.3.3.7 Oranges .....	33
1.3.3.8 Pomegranates .....	35
1.3.3.9 Avocados.....	36
1.3.3.10 Bitter melons .....	37
1.3.3.11 Broccoli.....	38
1.3.3.12 Mushrooms .....	39
1.3.3.13 Jerusalem artichokes .....	41
1.4 Conclusion.....	41
References .....	55
CHAPTER 2: .....	80
Impact of Increasing Concentration of Tart and Sweet Cherries Juices Concentrates on Healthy Mice Gut Microbiota.....	80
Abstract .....	80
Keywords: .....	81
2.1 Introduction .....	81
2.2 Materials and Methods .....	85
2.2.1 Animals.....	85



2.2.2 Treatments .....	85
2.2.3 Experiment intervention .....	85
2.2.4 DNA extraction .....	86
2.2.5 Universal polymerase chain reaction.....	87
2.2.6 Libraries Preparation and Sequencing.....	87
2.2.7 Statistical analysis .....	88
2.3 Results and Discussion.....	89
2.3.1 Nutritional analyses .....	89
2.3.2 Gut microbiota modulation.....	89
2.4 Conclusion.....	92
Acknowledgments .....	93
References .....	101
CHAPTER 3: .....	108
Impact of Cranberry Juice Consumption on Gut and Vaginal Microbiota in Postmenopausal Women.....	108
Abstract .....	108
Keywords: .....	109
3.1 Introduction .....	109
3.2 Materials and Methods .....	112
3.2.1 Subjects.....	112

3.2.2 Study Design and sample collection.....	112
3.2.3 DNA extraction .....	113
3.2.4 Universal polymerase chain reaction.....	113
3.2.5 Libraries Preparation and Sequencing.....	114
3.2.6 Statistical analyses.....	115
3.3 Results and Discussion.....	115
3.3.1 Impact of the cranberry juice on the gut microbiota .....	116
3.3.2 Impact of the cranberry juice on the vaginal microbiota.....	117
3.3.2.1 Lactobacillus dominant vaginal microbiota.....	117
3.3.2.2 Diverse baseline vaginal microbiota.....	117
3.3.2.3 Streptococcus dysbiosis .....	119
3.4 Conclusion.....	120
Acknowledgement:.....	120
Author Contributions: .....	120
References .....	127
CHAPTER 4: .....	134
Botanical Diversity and Amount of Dietary Fruits and Vegetables Distinctly Affect the Human Gut Microbiome.....	134
Abstract .....	134
Keywords: .....	135

4.1 Introduction .....	135
4.2 Materials and Methods .....	139
4.2.1 Subjects.....	139
4.2.1.1 SMART-Diet study.....	139
4.2.1.2 SMART-Valid study.....	139
4.2.2 Study Design and sample collection.....	140
4.2.2.1 SMART-Diet study.....	140
4.2.2.2 SMART-Valid study.....	140
4.2.3 DNA extraction .....	142
4.2.4 Universal polymerase chain reaction.....	142
4.2.5 Amplicon Library Preparation and Sequencing .....	143
4.2.6 Short-chain fatty acids analyses .....	144
4.2.7 Statistical analysis .....	144
4.3 Results and Discussion.....	145
4.3.1 SMART-Diet study: Longitudinal assessment of the impact of low and adequate FV intake .....	145
4.3.2 SMART-Valid: Impact of botanical diversity of a high FV diet.....	147
4.4 Conclusion.....	150
References .....	170
GENERAL CONCLUSION .....	178

APPENDIXES .....	180
------------------	-----

## LIST OF TABLES

### Chapter 1

<b>Table 1. 1:</b> Modulation of the gut microbiota by berries. ....	43
<b>Table 1. 2:</b> Modulation of the gut microbiota by cherries. ....	46
<b>Table 1. 3:</b> Modulation of the gut microbiota by mangos.....	47
<b>Table 1. 4:</b> Modulation of the gut microbiota by apples.....	47
<b>Table 1. 5:</b> Modulation of the gut microbiota by grapes.....	48
<b>Table 1. 6:</b> Modulation of the gut microbiota by kiwifruits.....	49
<b>Table 1. 7:</b> Modulation of the gut microbiota by oranges.....	50
<b>Table 1. 8:</b> Modulation of the gut microbiota by pomegranates.....	50
<b>Table 1. 9:</b> Modulation of the gut microbiota by avocados. ....	51
<b>Table 1. 10:</b> Modulation of the gut microbiota by bitter melons. ....	51
<b>Table 1. 11:</b> Modulation of the gut microbiota by vegetables. ....	52

### Chapter 3

<b>Table 3. 1:</b> The analytical data for the cranberry and placebo beverages. ND indicates non-detectable .....	122
<b>Table 3. 2:</b> Volunteers demographics. ....	122

### Chapter 4

<b>Table 4. 1:</b> Inclusion/Exclusion criteria for the SMART-Diet and SMART-Valid Studies.....	151
---	-----

## LIST OF FIGURES

### Chapter 2

**Figure 2. 1:** Study design and intervention. .... 94

**Figure 2. 2:** (A) Palatability of cherries juices concentrates as represented by daily water consumption.. (B) The impact of cherries juices consumption on weight trajectories for 23 days.. .... 95

**Figure 2. 3:** Cherries type (A) have less impact on gut microbiota than concentration and amounts consumed (B, C, and D). .... 96

**Figure 2. 4:** The impact of concentration of cherries juices on gut microbiota diversity.. .... 98

**Figure 2. 5:** Impact of varying concentrations of cherries juices on the phylum level..... 99

**Figure 2. 6:** Impact of varying concentrations of cherries juices on the genus level..... 100

### Chapter 3

**Figure 3. 1:** Timeline of intervention..... 122

**Figure 3. 2:** Impact of cranberry juice and placebo on the relative abundance of genera from Actinobacteria, Bacteroidetes, Verrucomicrobia, and Firmicutes. .... 123

**Figure 3. 3:** Non-metric Multidimensional Scaling (Bray-Curtis index) sowing the three different clusters of baseline vaginal microbiota. .... 124

**Figure 3. 4:** Diversity indices and changes in relative abundances in volunteers with *Lactobacillus*-dominated vaginal microbiota. .... 124

**Figure 3. 5:** Diversity indices changes in subjects with diverse vaginal microbiota at baseline. .... 125

**Figure 3. 6:** Impact of cranberry juice and the placebo on the relative abundance of the major genera in subjects with diverse vaginal microbiota at baseline.. .... 125

**Figure 3. 7:** Diversity indices changes in subjects with *Streptococcus* dysbiosis. .... 126

**Figure 3. 8:** Impact of cranberry juice and the placebo on the relative abundance of the major genera in subjects with *Streptococcus* dysbiosis at baseline. .... 126

### Chapter 4

**Figure 4. 1:** SMART-Diet study design and intervention..... 151

**Figure 4. 2:** SMART-Valid study design and intervention..... 152

<b>Figure 4. 3:</b> Impact on general microbiota profiles of the low and adequate FV treatment interventions at day 3, 6, and 9 compared to subjects' baseline. ....	153
<b>Figure 4. 4:</b> Impact on individuals' gut microbiota profiles (NMDS) of the low and high FV botanical biodiversity diet interventions.....	154
<b>Figure 4. 5:</b> Impact on diversity indices (Number of taxa, Shannon index) of the low and adequate FV treatment interventions at day 3, 6, and 9 compared to subjects' baseline.....	155
<b>Figure 4. 6:</b> Impact of low FV and adequate FV diets at the phylum level for each time point.....	156
<b>Figure 4. 7:</b> Impact of low FV and adequate FV diets on the composition of genera that belong to the Bacteroidetes phyla.....	157
<b>Figure 4. 8:</b> Impact of low FV and adequate FV diets on the composition of genera that belong to the Firmicutes phyla.....	158
<b>Figure 4. 9:</b> Impact of low FV and adequate FV diets on the composition of genera that belong to the Firmicutes phyla.....	159
<b>Figure 4. 10:</b> Impact of low FV and adequate FV diets on the composition of <i>Bifidobacterium</i> , <i>Akkermansia</i> , and Enterobacteriaceae. ....	160
<b>Figure 4. 11:</b> Impact on general microbiota profiles (NMDS) of the low and high FV botanical biodiversity diet interventions.....	161
<b>Figure 4. 12:</b> Impact on diversity indices (Number of taxa, Shannon index) of the low and high FV botanical diversity diet interventions. ....	162
<b>Figure 4. 13:</b> Impact of the low and high FV botanical diversity consumption at the phylum level.....	163
<b>Figure 4. 14:</b> Impact of the low and high FV botanical diversity consumption of the genera belong to the Bacteroidetes phyla.....	164
<b>Figure 4. 15:</b> Impact of the low and high FV botanical diversity consumption of the genera belong to the Firmicutes phyla.....	165
<b>Figure 4. 16:</b> Impact of the low and high FV botanical diversity consumption of <i>Bifidobacterium</i> , <i>Akkermansia</i> , and Enterobacteriaceae Unclassified. ....	168
<b>Figure 4. 17:</b> Impact of low and high FV botanical diversity consumption on SCFAs levels. .	169

## **LIST OF PUBLISHED AND SUBMITTED PAPERS**

Al Othaim, A., & Carbonero, F. (2021). *Impact of Fruit and Vegetable Inclusive Diets on Gut Microbiota Composition and Diversity*. [Submitted for publication; Chapter one]. Cell and Molecular Biology, University of Arkansas.

Al Othaim, A., Marasini, D., & Carbonero, F. (2020). Impact of increasing concentration of tart and sweet cherries juices concentrates on healthy mice gut microbiota. *Food Frontiers*, 1(3), 224-233. [Published; Chapter two].

Al Othaim, A., Marasini, D., & Carbonero, F. (2021). Impact of cranberry juice consumption on gut and vaginal microbiota in postmenopausal women. *Food Frontiers*, 1– 12. [Published; Chapter three].



## **GENERAL INTRODUCTION**

Diet plays a crucial role not only on the host's general health but also on modulating the gut microbiota ecosystem. This connecting role of gut microbiota has been illustrated by their impact on many aspects of human homeostasis including immune system modulation and metabolic and physiologic effects. The imbalance of the gut microbiota, often termed dysbiosis, has been associated with several diseases and metabolic disorders. With either beneficial or deleterious consequences, specific dietary patterns can impact the gut microbiota composition and its impact on health. A growing body of studies has shown the beneficial impact of plant-based diet on general health by preventing diseases and disorders as well as improving the gut microbiota ecosystem.

Plant-based diets typically consist of foods that are derived from plant sources, mainly fruits and vegetables. Fruits and vegetables are rich in bioactive compounds such as dietary fibers and polyphenols, as well as other essential nutrients, including vitamins and minerals. The impact of dietary fibers and polyphenols on the gut microbiota composition and diversity has been receiving attention in recent years. Gut microbes use dietary fibers as the primary source of energy and produce bioactive metabolites, such as short-chain fatty acids which are the main fermentation end-products (acetate, propionate, and butyrate). Some metabolites have been shown to exert beneficial impact on various physiological processes. Additionally, gut microbiota plays a vital role in polyphenols bioavailability. Polyphenols with high molecular weights cannot be absorbed in the small intestine; however, gut microbiota in the large intestine converts unabsorbed polyphenols to smaller bioactive metabolites which make them available to the host. These bioactive metabolites might be responsible for potential health benefits rather than the original native polyphenols.

The vast majority of published papers in the human nutrition and gut microbiome field have focused on studying the influence of extracted food compounds on gut microbiota ecosystem instead of whole foods. Therefore, this present dissertation brings attention to the impact of whole foods — including cherry juices, cranberries juice, and the amount and botanical diversity of fruit and vegetable diets — on the gut microbiota composition.

## CHAPTER 1:

### Impact of Fruit and Vegetable Inclusive Diets on Gut Microbiota Composition and Diversity

Ayoub Al Othaim<sup>1,2</sup> and Franck Carbonero<sup>1,3,4\*</sup>

<sup>1</sup> Cell and Molecular Biology Program; University of Arkansas, Fayetteville, AR, 72701 USA; [franck.carbonero@wsu.edu](mailto:franck.carbonero@wsu.edu)

<sup>2</sup>Department of Medical Laboratories, College of Applied Medical Sciences, Majmaah University, Al-Majmaah, 11952, Saudi Arabia

<sup>3</sup>Department of Food Science; University of Arkansas, Fayetteville, AR, 72704 USA

<sup>4</sup>Department of Nutrition and Exercise Physiology, Elson Floyd School of Medicine, Washington State University-Spokane, 412 East Spokane Falls Boulevard, Spokane, WA, 99202 USA

\* Correspondence: [franck.carbonero@wsu.edu](mailto:franck.carbonero@wsu.edu)

† The work is a part of Ayoub Al Othaim's doctoral thesis

#### Abstract

Fruits and vegetables are a fundamental source of dietary fibers, polyphenols, and essential nutrients, including vitamins and minerals. Nutritional recommendations emphasize fruit and vegetable consumption due to their health benefits on the host. Fruits and vegetables rich in dietary fibers and polyphenols can impact the gut microbiota composition and associated functions, whereas gut microbiota can metabolize components in fruits and vegetables into bioactive metabolites. Gut microbiota and a diet-rich in fruits and vegetables play a significant role in promoting the host's general health and preventing multiple diseases and metabolic disorders. This review recapitulates the latest findings on the impact of whole fruit and vegetable consumption on gut microbiota composition as well as the role of gut microbiota and a diet-rich in fruits and vegetables in disease and disorder prevention. This body of literature, while still limited especially in terms of randomized human dietary intervention, strongly suggests overall

beneficial modulation of the gut microbiome by common and less common fruits, and possibly vegetables.

**Keywords:** Fruits, Vegetables, Gut microbiota, Polyphenols, Dietary fibers.

## **1.1 Introduction**

Since the emergence of human nutrition as a scientific discipline and the development of nutritional guidelines, what constitutes a healthful diet is still intensely debated. Recently, the debate has become even more complicated with emerging evidence of the complex modulation of nutrients because of overlooked metabolic processes and their influence on health and disease. The mutualistic relationship between diet and gut microbiota is an example of exponential scientific interest. Diet is the most effective factor for shaping gut microbiota composition and consequently metabolites production and absorption with potential impacts on the host's general health.

Gut microbiota refers to the ecological communities that live in the gastrointestinal tract. There are trillions of microbes with thousands of life bacterial species in the human gut. The gut microbiota profile is not constant among humans, and is strongly individualized. This variation occurs due to several factors including host genetics, diet, age, mode of birth, and antibiotics. Disruption of the gut microbiota composition can alter gut homeostasis and thereby cause a variety of diseases. Therefore, the maintenance of the gut microbiota equilibrium is important to maintain a healthy status.

Diet is one of the most important factors that impact the shape, composition, and diversity of gut microbiota. This is to be expected, since microorganisms use undigested foods as fuel for their essential biological processes. Thus, changes in dietary patterns alter the bacterial

metabolism, which in turn influence the abundance of different types of bacteria in the gut. Specific dietary patterns strongly alter gut microbiota and can affect its impact on health, with either beneficial or deleterious consequences.

Western diet is typically characterized by high total energy due to saturated fat (butter, red meat) and sugar with an associated reduced consumption of fruits and vegetables. The high intake of energy-dense and processed foods is a risk factor for many chronic diseases including obesity, diabetes, inflammatory bowel diseases (IBD), and cardiovascular diseases. Inflammation is one of the underlying mechanisms linked with Western diet. There is a growing recognition for the influence of gut microbiota to the complexity of these disorders.

For decades, many studies have sought to substantiate the purported beneficial effects of fruits and vegetables-based diet. Epidemiological and dietary intervention studies have consistently indicated that plant-based diets provide a beneficial impact on human health and biomarkers. The present review discusses the importance of plant-based diets in shaping the gut microbiota and the potential health effects of these diet-microbiota interactions.

## **1.2 Plant-based dietary patterns**

A plant-based diet is a diet that is mostly or completely derived from plant sources. It can be obtained by increasing the intake of plant based foods such as fruits, vegetables, beans, seeds, pulses, herbs, whole grains, and nuts and reducing or excluding processed foods and animal products, including red meat, poultry, fish, eggs, and dairy products (Ostfeld, 2017; Tuso, Ismail, Ha, & Bartolotto, 2013). Plant-based dietary patterns have exploded in popularity over the past several decades. In the United States, the percentage of adults who chose a plant-based diet was approximately 3.3% in 2016 ("How Many Adults in the U.S are Vegetarian and Vegan?," 2016).

More recently, the department of agricultural economics at Oklahoma State University developed a project to track consumer's food preferences. This project was a series of monthly surveys called "Food Demand Survey (FooDS)" and was conducted between 2013 and 2018. The project data showed an increase in the number of self-identified vegetarians in the United States, around 4.1% between 2013 and 2018 ("Food demand Survey," 2018). Furthermore, the interest in the potential benefits of plant-based diets has increased in the last few decades. This not only underscores the increased interest in plant-based diets among consumers and food industries but also indicates the improved awareness and understanding of how plant-based diets impact human health.

### **1.2.1 Health benefits of plant-based diets**

Plant-based dietary patterns are typically rich in dietary fiber and a variety of phytonutrients, antioxidants, vitamins, and minerals and lower in fat, particularly saturated fat (Hever & Cronise, 2017). Plant-based diets have become a topic of formal scientific inquiry. A substantial number of studies have been conducted to confirm the beneficial effect of plant-based nutrition on the host. Plant-based foods have been associated with a reduced risk of many diseases and metabolic disorders, including obesity, diabetes, cardiovascular diseases, and cancer.

#### ***1.2.1.1 Lower risk of obesity***

Obesity is a serious pandemic worldwide. The World Health Organization (WHO) estimated that more than 1.9 billion people are obese or overweight in 2016 ("Obesity and overweight," 2020). According to the Centers for Disease Control and Prevention (CDC), the prevalence of obesity in the United States among adults was 42.4% in 2017-2018 (Hales, Carroll,

Fryar, & Ogden, 2020). Obesity is associated with severe health conditions and triggers obesity-related diseases, including type 2 diabetes, cardiovascular diseases, and cancers ("Obesity and overweight," 2020; Wright, Wilson, Smith, Duncan, & McHugh, 2017).

Poor diet plays a key role in the genesis of obesity. In contrast, plant-based diets show evidence to achieve and maintain healthy body weight (Melina, Craig, & Levin, 2016). A growing body of studies has examined the use of plant-based dietary patterns for weight loss and obesity prevention. Berkow and Barnard reviewed numerous published articles containing information on vegetarians and plant-based diets. While the systematic review indicated the impact of plant-derived foods in weight loss compared with other diets, there was also a clear pattern for subjects who follow plant-based diets to be at a lower risk of obesity. The observational studies showed a lower average body weight in vegetarian individuals compared to non-vegetarians, approximately 3% to 20%. In addition, vegetarians have a consistently lower obesity prevalence rate, 0% to 6%, compared to non-vegetarians, 5% to 45% (Berkow & Barnard, 2006). The Oxford component of the European Prospective Investigation into Cancer and Nutrition (EPIC) compared changes in weight and body mass index (BMI) over a five-year period in approximately 22,000 meat-eating, fish-eating, vegetarian, and vegan participants in the United Kingdom. The study results showed that the lowest weight gain was associated with individuals who followed a vegan diet and those who had changed to a diet containing less animal products. The result also showed a significant difference in BMI between the four diet groups, with the meat-eaters having the highest BMI and the vegans had the lowest BMI (Rosell, Appleby, Spencer, & Key, 2006). In 2016, a meta-analysis of randomized controlled trials was performed to compare the influence of vegetarian diets and non-vegetarian diets on weight loss among individuals. This review included twelve randomized controlled studies with

approximately 1,150 subjects. The results showed a significant weight reduction among individuals assigned to vegetarian diets compared to individuals assigned to non-vegetarian diets (Huang, Huang, Hu, & Chavarro, 2016). A randomized controlled trial study was performed to investigate the impact of a whole food plant-based diet on either obese or overweight individuals with a diagnosis of either type 2 diabetes, ischaemic heart disease, or the cardiovascular risk factors of hypertension or hypercholesterolaemia. At 6 and 12 months, the whole food plant-based diet reduced the BMI to 4.4 and 4.2 kg m<sup>-2</sup>, respectively, compared with the subjects' typical diet. Furthermore, the total cholesterol tended to decrease after the implementation of whole food plant-based diet in both intervention periods (Wright et al., 2017). Data from the Adventist Health Study-2 showed that obesity and abdominal adiposity were significantly lower among non-Hispanic white adults who followed a vegetarian diet (Matsumoto et al., 2019).

#### ***1.2.1.2 Lower risk of diabetes***

Diabetes is one of the most significant global health crises of this century ("Diabetes," 2020). Globally, the number of people diagnosed with diabetes in 2017 was approximately 425 million (Cho et al., 2018). In the United States, around 13% of all US adults over 18 years of age had been diagnosed diabetes in 2018 (Centers for Disease Control and, 2020). Lifestyle changes, mainly dietary changes, can play a crucial role in preventing, treating, or promoting type 2 diabetes (McMacken & Shah, 2017). Unlike calorie-dense foods, such as diets high in saturated fat and sugar, plant-based diets may effectively prevent or treat type 2 diabetes (McMacken & Shah, 2017; Papier et al., 2019).

The association between dietary patterns and the incidence of diabetes was investigated using data from the Adventist Health Study-2. The results showed that vegans, lacto ovo-



vegetarians, and semi-vegetarians were less likely to develop diabetes (Tonstad et al., 2013). A meta-analysis study showed that fruit and vegetable consumption, especially green leafy vegetables, is associated with a significant reduction of type 2 diabetes incidence (M. Li, Fan, Zhang, Hou, & Tang, 2014). Data from the Nurses' Health Study, Nurses' Health Study 2, and the Health Professionals Follow-up Study suggested that plant-based diets, mainly diets with high-quality plant foods, are associated with a lower risk of developing type 2 diabetes (Satija et al., 2016). A study was conducted to investigate the correlation between a vegetarian diet and the risk of diabetes in a Taiwanese population which stated that a vegetarian diet might protect against diabetes (Chiu, Pan, Lin, & Lin, 2018). The EPIC-Oxford cohort study was used to examine the association between vegetarian diets and the risk of type 2 diabetes in British adults. The study results suggested that vegetarian and low-meat diets may reduce the risk of developing diabetes development (Papier et al., 2019).

#### ***1.2.1.3 Lower risk of cardiovascular diseases***

Cardiovascular diseases are the most prevalent cause of death worldwide, with more than 17.6 million deaths per year. Approximately one-third of all global deaths are caused by cardiovascular diseases ("Cardiovascular diseases (CVDs)," 2017; "Heart Disease and Stroke Statistics-2019 At-a-Glance," 2019). Plant-based diets are associated with a reduction in the risk of cardiovascular diseases and cardiovascular disease mortality (Dinu, Abbate, Gensini, Casini, & Sofi, 2017; H. Kim et al., 2019). Plant-derived foods also showed evidence of reducing heart disease risk factors such as serum total cholesterol, blood glucose, abdominal obesity, and blood pressure (Rizzo, Sabaté, Jaceldo-Siegl, & Fraser, 2011; Wang et al., 2015; Yokoyama, Barnard, Levin, & Watanabe, 2014; Yokoyama, Nishimura, et al., 2014).

The EPIC-Oxford cohort study results were used to investigate the impact of different dietary patterns on serum lipid concentration. The study showed that vegans had the lowest total cholesterol in serum compared to individuals assigned to other diet patterns (Bradbury et al., 2014). A meta-analysis of 11 randomized controlled trials found that the total low-density lipoprotein and high-density lipoprotein cholesterol levels were low among vegetarians. These findings, perhaps, explain why plant-based diets reduce the risk of cardiovascular disease and cardiovascular disease mortality (Wang et al., 2015). A comprehensive meta-analysis reported that vegetarian diets significantly decrease the incidence and/or mortality from ischaemic heart disease by about 25% (Dinu et al., 2017). Data from the Adventist Health Study-2 (AHS-2) were analyzed to assess the association between dietary patterns and cardiovascular disease risk factors among non-Hispanic white adults. The investigation showed that vegetarian diets were associated with significantly lower levels of cardiovascular disease risk factors than non-vegetarian diets (Matsumoto et al., 2019).

#### ***1.2.1.4 Lower risk of cancers***

Cancer is a major public health issue worldwide and is the second leading cause of death globally. In 2018, the total number of deaths caused by cancer was around 9.6 million deaths, meaning that one of every six deaths was cancer-related ("Cancer," 2018). According to The American Cancer Society (ACS), the estimated number of new cancer cases and cancer deaths in the United States was 1.8 million and 606,520, respectively, in 2020 (Siegel, Miller, & Jemal, 2020). Plant-based dietary patterns have been associated with reducing the risk of cancer occurrence.

Data from the Adventist Health Study-2 was used to investigate the association between plant-based dietary patterns and the overall cancer incidence among 69,120 participants. The results showed that vegetarian diets seem to be linked with lower overall cancer risks, especially a lower risk of gastrointestinal cancer. The vegan diet appeared to confer greater protection against overall cancer incidence, especially female-specific cancer, than any other dietary patterns (Tantamango-Bartley, Jaceldo-Siegl, Fan, & Fraser, 2013). A study using the Adventist Health Study-2 data stated that vegetarian diets potentially associated with lowering the incidence of colorectal cancers (Orlich et al., 2015). Other papers reported that a vegan diet might correlate with a lower risk of prostate cancer (Tantamango-Bartley et al., 2016). A comprehensive meta-analysis study revealed that vegan and vegetarian diets significantly reduced the incidence and mortality of all cancers by around 15% and 8% respectively (Dinu et al., 2017).

Based on prior studies, it is clear that dietary patterns that are higher in plant foods and lower in animal foods are promoting general health and lowering the incidence of several diseases and metabolic disorders.

### **1.3 Gut Microbiota Modulation by Plant-Based Foods**

#### **1.3.1 Gut Microbiota: General Aspects**

The human gastrointestinal tract harbors diverse and complex microbial community composed mainly of bacteria and termed gut microbiota. The actual numbers of microorganisms inhabiting the gastrointestinal tract have been the subject of controversies and exaggerated claims. The initial estimate of the number of bacteria inhabiting the gastrointestinal tract was  $10^{14}$ , which is approximately ten times more than the cells of human body (Savage, 1977);

however, a more recent calculation estimated there to be  $10^{13}$  bacteria inhabiting the gastrointestinal tract making the ratio of bacteria to human cells closer to 1:1 (Sender, Fuchs, & Milo, 2016). The bacterial density and diversity in the intestinal tract are quite different depending on the location. The colon has the highest bacterial biodiversity, whereas the upper part has the lowest number of microbes (Hillman, Lu, Yao, & Nakatsu, 2017). Gut microbiota is referred to as the “hidden metabolic organ” due to its substantial impact on general human health, including metabolism, nutrition, and physiological and immune function (Thursby & Juge, 2017).

In earlier times, members of a microbial community were studied and identified using culture-dependent approaches. These methods are broadly used in the microbiology field; however, they have some limitations, especially in the study of microbial ecology. The culture-dependent methods are limited in terms of bacterial identification as a wide range of bacterial strains cannot be grown and cultured in the laboratory. Additionally, these methods are limited to distinguish bacteria at lower taxonomic levels (Morgan & Huttenhower, 2012).

Recently, culture-independent approaches, including high-throughput and low-cost sequencing methods, have been widely used to investigate numerous characteristics of microbial communities such as taxonomic diversity and functional metagenomics. 16S ribosomal RNA (rRNA) gene has been by far the most common phylogenetic marker used to study gut microbial diversity and taxonomy (Poretsky, Rodriguez-R, Luo, Tsementzi, & Konstantinidis, 2014). Many aspects make the 16S rRNA gene the ideal targeted gene for gut microbial phylogeny and taxonomy studies. These aspects include (1) the presence in all bacteria and archaea; (2) the evolution at relatively constant rates; (3) and the existence of conservative regions, used for designing amplification primers, and highly variable regions (V1-V9), used for distinguishing

and identifying bacterial species (Janda & Abbott, 2007; Mizrahi-Man, Davenport, & Gilad, 2013).

Based on 16S rRNA-based studies, human intestinal bacteria involve around 1000 bacterial species belonging to six different bacterial phyla. Bacteroidetes and Firmicutes are the most dominant phyla, whereas Actinobacteria, Proteobacteria, Verrucomicrobia, and Fusobacteria mostly present in lower proportions (Eckburg et al., 2005; Huttenhower et al., 2012; Rajilić-Stojanović & de Vos, 2014; Rowland et al., 2018). In lower taxonomic levels, bacteria like *Bifidobacterium*, *Lactobacillus*, Clostridiaceae, Lachnospiraceae, Ruminococcaceae, *Prevotella*, and *Akkermansia* are considered beneficial bacteria, while *Escherichia coli*, *Salmonella*, *Staphylococcus*, *Bacillus*, and *Enterococcus* are potential pathogens (Cui et al., 2019; Lavefve, Howard, & Carbonero, 2020).

Gut microbiota collaborates with the host immune system to form a complex and reciprocally beneficial relationship (Bäckhed, Ley, Sonnenburg, Peterson, & Gordon, 2005; Lazar et al., 2018; Round & Mazmanian, 2009, 2010; H.-J. Wu & Wu, 2012). Gut microbiota reinforces the function of the immune system and prevents the colonization of pathogens (Okumura & Takeda, 2017; Ubeda, Djukovic, & Isaac, 2017). On the other hand, the immune system has co-evolved to live in a cooperative relationship with the healthy microbiota and fight against pathogens (Jandhyala et al., 2015; Round & Mazmanian, 2009, 2010). In addition to the immunological protective functions, gut microbiota helps to maintain the integrity of the mucosal barrier and regulate host digestion and metabolism (Okumura & Takeda, 2018).

Changing in the gut microbial composition can lead to beneficial or harmful effects on human health (Thursby & Juge, 2017). The imbalance of the gut microbiota symbiotic

ecosystem, known as dysbiosis, links to several pathologies including IBD, celiac disease, and metabolic disorders (Biragyn & Ferrucci, 2018; Schippa & Conte, 2014). Life style changes might disrupt the gut microbiota ecosystem, which may explain the increased incidence of microbiome-linked pathologies over the past few decades (Logan, Jacka, & Prescott, 2016). Many factors can abnormally shift in the composition of gut microbiota and, in turn, increase the incidence of diseases including diet, environmental changes, physical and psychological stress, and antibiotic exposure (Karl et al., 2018). Many scientific studies have linked the domination of some gut microbiota groups with positive health outcomes, and these microbes are commonly identified as "potentially beneficial bacteria." The most common bacterial strains that fall in this category belong to the genera *Bifidobacterium*, *Lactobacillus*, *Akkermansia*, *Faecalibacterium*, *Eubacterium*, *Roseburia*, and *Blautia* (Hiippala et al., 2018; LeBlanc et al., 2017; Quigley, 2013; Q. Yang et al., 2020; Y.-J. Zhang et al., 2015). Other bacterial groups have been associated with the development or progression of several diseases and disorders, and these groups include some species from the genera *Clostridium*, *Enterobacter*, *Enterococcus*, and *Bacteroides* (Alhinai, Walton, & Commane, 2019; Gagnière et al., 2016; Kowalska-Duplaga et al., 2019; Q. Yang et al., 2020). The potential correlation between the Firmicutes and Bacteroidetes population and BMI has been intensively studied. Several studies demonstrated that the increase of the Firmicutes/Bacteroidetes ratio might contribute to the pathophysiology of obesity (Barlow, Yu, & Mathur, 2015; Fernandes, Su, Rahat-Rozenbloom, Wolever, & Comelli, 2014; Kasai et al., 2015; Koliada et al., 2017; Mathur & Barlow, 2015; Turnbaugh et al., 2006). Other studies, however, have contradictory results and showed no statistically significant difference in the Firmicutes/Bacteroidetes ratio between obese and lean individuals (Aguirre & Venema, 2015; H.-J. Hu et al., 2015; Jumpertz et al., 2011; Tims et al., 2013; Walters, Xu, & Knight, 2014). The

contradicting results from previous studies suggest that the Firmicutes/Bacteroidetes ratio may not be valid as an eventual biomarker for obesity.

### **1.3.2 Effect of Plant-Based Diets on Gut Microbiota Composition**

The contribution of dietary patterns to modulate and modify the community structure and function of gut microbiota has been a hot topic in recent decades. Diet can directly influence the microbial composition in the gastrointestinal tract and play a key role in promoting or inhibiting their growth. Different dietary patterns, such as plant-based diets and the Western-style diet, have a significant impact on the gut microbiota ecology (Tomova et al., 2019). The Western-style diet, which is low in dietary fibers and high in fat content, has been linked to microbial disruptions in the digestive system and a reduction of microbial diversity (Sonnenburg et al., 2016). On the other hand, plant-based diets promote healthy and stable gut microbiota due to their high content of two critical nutrients, dietary fibers and phytonutrients (Hever & Cronise, 2017; Tomova et al., 2019).

Dietary fibers are edible carbohydrate polymers with three or more monomeric units that resist hydrolysis by digestive enzymes in the small intestine but can be fermented by microbial residents in the large intestine (Gentile & Weir, 2018; Williams, Grant, Gidley, & Mikkelsen, 2017). Gut microbiota uses dietary fibers as the primary source of energy through a metabolic process called fermentation. This fermentative activity results in the generation of bioactive metabolites such as short-chain fatty acids (SCFAs), primarily acetate, propionate, and butyrate. These organic products have a beneficial impact on various physiological processes, including the regulation of host metabolism, immune system, and cell proliferation (Koh, De Vadder, Kovatcheva-Datchary, & Bäckhed, 2016). Dietary fiber intake is not only promoting a healthy

environment in the gut but also enhances the richness and diversity of the gut microbiota (Cui et al., 2019).

Phytonutrients are a wide variety of chemical compounds produced by plants. One of the major types of these plant products is dietary polyphenols. Dietary polyphenols are naturally found in many foods and beverages, namely in fruits, vegetables, cereals, tea, coffee, and wine (Kumar Singh et al., 2019). During recent decades, there has been a growing interest in the bioactivity of polyphenol-rich foods. Many clinical and preclinical studies have been published showing the antioxidant and anti-inflammatory properties of phenolic compounds. This may suggest a potential link between polyphenol-rich food consumption and the reduction of several chronic disorders including obesity, diabetes, cancers, IBD, cardiovascular disease, and neurodegenerative diseases (Costa et al., 2017; Crascì, Lauro, Puglisi, & Panico, 2018; King, Kumar, & Kumar, 2018; Rahman et al., 2018).

Chemically, polyphenols are a large heterogeneous group of compounds characterized by hydroxylated phenyl moieties. Generally, polyphenols are classified into three major classes, flavonoids, stilbenoids, and phenolic acids, and flavonoids are the most prevalent class among them (Papuc, Goran, Predescu, Nicorescu, & Stefan, 2017). There are more than 9,000 different chemical structures of flavonoids that have been identified in nature. Based on their chemical structure differences, flavonoids are classified into subclasses including, anthocyanidins, flavanones, flavones, flavanonols, flavonols, flavan-3-ols or flavanols, isoflavones, neoflavonoids, and chalcones (Cardona, Andrés-Lacueva, Tulipani, Tinahones, & Queipo-Ortuño, 2013; Kawabata, Yoshioka, & Terao, 2019; Papuc et al., 2017).



The beneficial properties of dietary polyphenols are primarily reliant on their bioavailability. Gut microbial residents play a vital role in polyphenols bioavailability, especially in the colon. In general, approximately 5–10% of the total digested polyphenols are absorbed in the small intestine, while the remaining 90–95% pass to the large intestine (Peron et al., 2019). Typically, polyphenols with a low molecular weight are partially absorbed in the small intestine. However, the vast majority of polyphenols have a high molecular weight in their native state, making them unabsorbed in the small intestine (Cassidy & Minihane, 2017; Kawabata et al., 2019). The high molecular weight polyphenols and other unabsorbed low molecular weight are transported into the large intestine where the gut microbial community in the colon breaks them down into absorbable low molecular weight phenolic metabolites (Cardona et al., 2013; Cassidy & Minihane, 2017). These metabolic products might be responsible for the potential health benefits derived from polyphenol-rich foods, rather than the original native polyphenols (Cardona et al., 2013).

Consuming a diet rich in fruits and vegetables exerts an important influence on gut microbiota diversity and ecological structure. Fruits and vegetables are important foods in daily meals (Sharoba, Farrag, & Abd El-Salam, 2013). In accordance with the US Department of Health and Human Services (DHHS) and the United States Department of Agriculture (USDA), fruits and vegetables should make up half of the meal ("2015–2020 Dietary Guidelines," 2015). Due to the high content of dietary fibers and phenolic compounds, fruit and vegetable intake improves the richness and diversity of the gut microbial community. The increased interest and awareness among consumers in maintaining a healthy diet urges scientists and researchers to intensify their efforts in studying health-promoting food.

### 1.3.3 Impact of Fruits and Vegetables on Gut Microbiota Composition

Fruits and vegetables are a fundamental source of dietary fibers, polyphenols, and micronutrients such as vitamins and minerals. Many studies have investigated the potential impact of the consumption of fruits and vegetables and their extracts on the gut microbiota ecosystem. This review summarizes the latest findings on the effects of whole fruits and vegetables as well as their products, such as juices, on the gut microbiota composition and diversity.

#### 1.3.3.1 Berries

The impact of berry consumption on the composition and diversity of the gut microbial community has been studied intensively in recent years. In general, berries are an excellent source of bioactive compounds, mainly phenolic compounds, including anthocyanins, flavonols, flavanols, tannins, or phenolic acids (Nile & Park, 2014). Recently an exhaustive was published review which covers the specific area of berries and gut microbiome and health (Lavefve et al., 2020). This review will focus only on gut microbiota composition reports (**Table 1. 1**).

##### 1.3.3.1.1 Strawberries

Strawberry consumption altered the gut microbial composition in diabetic *db/db* mice. The phylum Actinobacteria slightly increased after the strawberry supplementation with a significant increase of its commensal genera *Bifidobacterium*. The phylum *Verrucomicrobia* and the genera *Bacteroides* and *Akkermansia* were significantly reduced in the mice that ate strawberries with no impact observed to *Dehalobacterium*, *Dorea*, *Lactobacillus*, and *Turicibacter* (Petersen et al., 2019). Moreover, strawberry administration alleviated the gut microbiota dysbiosis caused by the dextran-sulfate-sodium (DSS)-induced colitis in mice.

Strawberry supplementation induced the abundance of *Bifidobacterium* and *Lactobacillus* and reduced the abundance of *Dorea*, *Bilophila*, *Akkermansia*, and *Bacteroides* in colitis mice compared to the control group (Y. Han et al., 2019). The increased abundance of *Bilophila*, an opportunistic pathogen, was observed in colitis mice (I. Yang et al., 2013). *Bifidobacterium* and *Lactobacillus* spp. are generally considered probiotic microorganisms (Aoki et al., 2017). Although *Akkermansia* has been claimed to associate mostly with healthier clinical profiles, recent studies showed that the relative abundance of *Akkermansia* was elevated in several diseases, such as Parkinson's disease and Alzheimer's disease (Cirstea, Radisavljevic, & Finlay, 2018; Heintz-Buschart et al., 2018; Hill-Burns et al., 2017; Vogt et al., 2017).

A randomized controlled trial showed that the mixture of strawberries, raspberries, and cloudberries potentially altered the gut microbiota composition in human subjects with symptoms of metabolic syndrome. Unfortunately, the individual variations in the gut microbiota composition were not provided (Puupponen-Pimiä et al., 2013).

#### 1.3.3.1.2 Cranberries

Whole cranberries altered the gut microbiota community of a human *in vitro* by increasing the relative abundance of the families Porphyromonadaceae and Bacteroidaceae and stabilizing the commensal families Ruminococcaceae and Lachnospiraceae. Additionally, a reduction in the relative abundance of Enterobacteriaceae was observed after the cranberry treatment (O'Connor et al., 2019). A high level of the Enterobacteriaceae has often been reported in patients with IBD (Zeng, Inohara, & Nuñez, 2017).

*In vitro*, dietary cranberry (*Vaccinium macrocarpon*) ameliorated the gut microbiota dysbiosis caused by DSS-induced colitis in mice by reversing the reduction of the  $\alpha$ -diversity. At

the genus level, cranberry consumption altered the gut microbiota composition by increasing the relative abundance of potentially beneficial bacteria, e.g., *Bifidobacterium*, *Lactobacillus*, and *Akkermansia*, and decreasing the relative abundance of potentially harmful bacteria, e.g., *Sutterella* and *Bilophila* (Cai et al., 2019).

Cranberry consumption altered gut microbiota relative abundance in a short-period (5 days) in healthy human volunteers. The Firmicutes/Bacteroidetes ratio was influenced by the cranberry intake with an increase in Bacteroidetes level and a decrease in Firmicutes level. Cranberry intake also impacts bacterial abundance in the genus level by increasing the abundance of *Lachnospira* and *Anaerostipes*, SCFAs-producing bacteria, and decreasing the abundance of *Oribacterium* (Rodríguez-Morató, Matthan, Liu, de la Torre, & Chen, 2018).

#### 1.3.3.1.3 Black and red raspberries

The impact of red and black raspberries on reshaping the gut microbial composition has also been studied. Black raspberries significantly increased the genera *Akkermansia* and *Desulfovibrio* abundance in healthy rats. A number of butyrate-producing genera, such as *Anaerostipes*, were also increased after the black raspberry consumption (Pan et al., 2017). A similar impact was observed in a study using specific-pathogen-free mice where the black raspberry supplementation remarkably increased the abundance of the phylum Verrucomicrobia and its genera *Akkermansia muciniphila*. Black raspberries also altered the Firmicutes/Bacteroidetes ratio by increasing the relative abundance of Bacteroidetes and reducing Firmicutes abundance (Tu et al., 2018). *A. muciniphila* is believed to have several health benefits to humans and inversely correlated with various diseases (Cani & de Vos, 2017; Jayachandran, Chung, & Xu, 2019). Another mice study also supported the effect of black

raspberries on lowering the Firmicutes/Bacteroidetes ratio. At the genus level, *Barnesiella* and *Turicibacter* were significantly higher in mice fed a black raspberry-rich diet compared to the control group, while *Clostridium* and *Lactobacillus* were significantly lower (Gu et al., 2019). *Barnesiella*, a recently discovered genus, belongs to the Porphyromonadaceae family which was found to metabolize glucose to produce butyrate and isobutyrate (Sakamoto, Lan, & Benno, 2007; Sakamoto et al., 2009; Wylie et al., 2012). Additionally, a positive correlation between *Barnesiella* and the level of SCFAs has been reported (Ying Zhao et al., 2013). *Turicibacter* was found to have a positive correlation with butyric acid (Zhong, Nyman, & Fåk, 2015).

Supplementation with red raspberries was associated with significant changes in the fecal microbiota of obese diabetic mice. The relative abundance of Lachnospiraceae was increased in the red raspberry supplemented group. Additionally, a decrease in the relative abundance of *Lactobacillus*, *Odoribacter*, and the fiber degrader S24-7 family as well as unknown groups of Bacteroidales and Enterobacteriaceae was observed after red raspberry supplementation (Garcia-Mazcorro, Pedreschi, et al., 2018).

#### 1.3.3.1.4 Blueberries

Blueberries markedly impacted the gut microbiota composition. In the IBD mouse model, a blueberry-rich diet significantly reduced *Clostridium perfringens*, *Enterococcus*, *E. coli*, and *Lactobacillus* with no impact to *Bacteroides-Prevotella-Porphyromonas* group and *Bifidobacterium* (Paturi et al., 2012). *C. perfringens* and *Enterococcus faecalis* are potentially associated with IBD (Balish & Warner, 2002; Falk et al., 2007; Paturi et al., 2012; Tannock, 2010). Furthermore, blueberry consumption led to lowering the Firmicutes/Bacteroidetes ratio in

mice. The abundance of the phylum Tenericutes increased, and the family Deferribacteres decreased in blueberry-fed mice (Wankhade et al., 2019). Another study showed that blueberry administration increased the relative abundance of *Turicibacter* sp. H121 in mice fed a high-fat and high-sucrose diet (Morissette et al., 2020).

In rats, a Lowbush wild blueberry-enriched diet influenced the gut microbiota profile by significantly increasing the relative abundance of the phylum Actinobacteria, the order Actinomycetales, and a number of genera under the family Bifidobacteriaceae and Coriobacteriaceae, and significantly decreasing the relative abundance of *Lactobacillus* and *Enterococcus*. The Lowbush wild blueberry consumption increased the *Bifidobacterium*/Coriobacteriaceae ratio compared with the control diet (Lacombe et al., 2013). The increased *Bifidobacterium*/Coriobacteriaceae ratio positively impacted plasma cholesterol levels in hamsters (Martínez et al., 2009). Blueberry juice supplementation contributed to ameliorating the alteration in the gut microbiota composition caused by carbon tetrachloride (CCl<sub>4</sub>) in a CCl<sub>4</sub>-induced rat liver fibrosis model (Yan et al., 2019).

In healthy humans, wild blueberry (*Vaccinium angustifolium*) consumption modulated the intestinal microbiota. The abundance of commensal *Bifidobacterium* was significantly increased following the wild blueberry treatment with no significant impact to *Bacteroides*, *Prevotella*, *Enterococcus*, and *Clostridium coccoides* (Vendrame et al., 2011). Another human study showed that the consumption of a wild blueberry drink modulated the most abundant bifidobacterial taxonomic groups with significant stimulation of *Bifidobacterium longum subsp. infantis* strain (Guglielmetti et al., 2013).

#### 1.3.3.1.5 Other berries

A variety of eight Nordic berries; strawberry (*Fragaria ananassa* Senga Sengana), bilberry (*Vaccinium myrtillus*), red raspberry (*Rubus idaeus*, var. Ottawa), cloudberry (*Rubus chamaemorus*), blackcurrant (*Ribes nigrum*, var. Öjeby), and sea buckthorn berry (*Hippophae rhamnoides*) showed antimicrobial activity against selected Gram-positive and Gram-negative bacteria *in vitro*. *Staphylococcus* and *Salmonella* were the most sensitive bacteria to the berry intervention (Puupponen-Pimiä et al., 2005). *Salmonella* is a well-known foodborne pathogen and considered the leading bacterial cause of acute gastroenteritis (Majowicz et al., 2010; Rivera-Chávez et al., 2016). Sea buckthorn berry juice influenced gut microbiota *in vitro* by increasing the growth of *Bifidobacteria*, *Bacteroides/Prevotella*, and Lactic acid bacteria (Attri, Sharma, Raigond, & Goel, 2018). A similar impact on the beneficial microbial population was observed in a following *in vitro* study when the sea buckthorn berry juice supplementation stimulated the growth of *Bifidobacteria*, *Lactobacilli*, and *Bacteroides/Prevotella* (Attri & Goel, 2018). *Bifidobacterium* and *Lactobacillus* spp. are well-known probiotic microorganisms (Aoki et al., 2017).

Juçara (*Euterpe edulis*) significantly increased the relative abundance of *Bifidobacterium* after 24 hours of fermentation *in vitro* (Guergoletto, Costabile, Flores, Garcia, & Gibson, 2016). Similarly, juçara pulp reinstated *Bifidobacterium* content in rats fed a high-fat diet (Jamar et al., 2018). *Bifidobacterium* has been found to play a role in preventing cognitive impairment in Alzheimer's disease (Kobayashi et al., 2017).

*Lonicera caerulea* L. Berry (LCBP) modified the gut microbiota composition in a high-fat diet-induced mice model (S. Wu et al., 2018). The results revealed that supplementation with

LCBP impacted the Firmicutes/Bacteroidetes ratio by lowering the abundance of Firmicutes and elevating Bacteroidetes abundance as well as increasing the abundance of Parabacteroides and Verrucomicrobia. In the genus level, the LCBP diet induced the relative abundance of *Akkermansia*, which was suggested to play a role against obesity (Everard et al., 2013; Plovier et al., 2017). LCBP supplementation increased the relative abundance of *Bacteroides* and *Parabacteroides*, which were found to be reduced with non-alcoholic fatty liver disease and the high fat-induced diet (Del Chierico et al., 2017; Liu et al., 2016). Additionally, the relative abundance of *Staphylococcus*, *Lactobacillus*, *Oscillospira*, and *Ruminococcus* was reduced after LCBP supplementation. The increased abundance of *Lactobacillus*, *Ruminococcus*, and *Oscillospira* was associated with intestinal permeability, systemic inflammation, and macrophage dysfunction (Santisteban et al., 2017; Thevaranjan et al., 2017; S. Wu et al., 2018). These findings express the potential health benefits of LCBP.

Goji berry (*Lycium barbarum*) modulated the gut microbial population in IL-10-deficient mice, which will develop a chronic IBD spontaneously. Dietary Goji markedly enhanced the growth of Actinobacteria phylum and its genera *Bifidobacterium*. Butyrate-producing bacteria, including Lachnospiraceae family and its genera *Roseburia* spp. and Ruminococcaceae family and its genera *Faecalibacterium prausnitzii*, were also promoted by Goji berry (Kang, Yang, Zhang, Ross, & Zhu, 2018). These genera have been described as potentially protective against human IBD (Kumari, Ahuja, & Paul, 2013; Martín et al., 2014; Rossi et al., 2016; Tamanai-Shacoori et al., 2017). *F. prausnitzii* dysbiosis is associated with several intestinal and metabolic disorders such as IBD, irritable bowel syndrome (IBS), colorectal cancer (CRC), obesity, and celiac disease (Balamurugan, Rajendiran, George, Samuel, & Ramakrishna, 2008; De Palma et al., 2010; Furet et al., 2010; Neish, 2009; Rajilić-Stojanović et al., 2011; Sokol et al., 2008).



Furthermore, many studies have reported the anti-inflammatory effects of *F. prausnitzii* (Martín et al., 2017; Miquel et al., 2015; Quévrain et al., 2016; Rossi et al., 2015; Sarraiyayrouse et al., 2014).

Lingonberries lowered the Firmicutes/Bacteroidetes ratio in *Apoe*<sup>-/-</sup> mice, atherosclerosis-prone mice, fed a high-fat diet. The genera *Bacteroides*, *Clostridium*, and *Parabacteroides*, and the species *A. muciniphila*, *Blautia producta*, *Clostridium difficile*, and *Eubacterium dolichum* were increased after Lingonberries supplementation. A decrease in the abundance of *Mucispirillum* and *Oscillospira* was observed in mice fed the high-fat diet (Matziouridou, Marungruang, Nguyen, Nyman, & Fåk, 2016). *E. dolichum* is a butyrate-producer bacterium, and butyrate has anti-inflammatory properties and can play a role against the development of IBD (M. Zhang et al., 2016).

A berry mixture is consisting of blueberries, blackberries, raspberries, Portuguese crowberries, and strawberries mitigated the gut microbiota alteration caused by a salt-induced hypertension diet in rats (Gomes et al., 2019). The Bacteroidetes population was significantly increased after the berry mixture intake while the Firmicutes and Proteobacteria were decreased. The berry mixture supplementation increased the abundance of Bacteroidaceae, S24-7, and Ruminococcaceae families. Moreover, the berry mixture intake significantly decreased the relative abundance of Erysipelotrichaceae, which has been correlated with inflammation (Dinh et al., 2015; Gomes et al., 2019).

Blackcurrants and raspberries modulated the gut microbiota composition in rats. The result showed a clear difference in the diversity index between the rats fed blackcurrants and raspberries as the diversity index was higher for the rats fed raspberries than blackcurrants.

However, the modulation in the composition of specific bacteria has not been reported (Jakobsdottir et al., 2013).

A randomized, double-blind placebo-controlled study revealed that *Schisandra shinensis* fruit modulates the gut microbiota composition in obese women. *Schisandra chinensis* fruit intake lowered the Firmicutes/Bacteroidetes ratio by increasing Bacteroidetes abundance and decreasing Firmicutes abundance. At the genus level, *Bifidobacterium*, *Akkermansia*, *Roseburia*, *Prevotella*, and *Bacteroides* showed a superior increase in the treatment group compared to the placebo group (Song, Wang, Eom, & Kim, 2015). These bacteria were found to be associated with reducing and controlling obesity (Dao et al., 2016; Everard et al., 2013; Hjorth et al., 2018; H. I. Kim et al., 2019; Neyrinck et al., 2012; Plovier et al., 2017; Tamanai-Shacoori et al., 2017; J. Y. Yang et al., 2017).

A variety of eight Nordic berries; bilberry (*Vaccinium myrtillus*), red raspberry (*Rubus idaeus*, var. Ottawa), cloudberry (*Rubus chamaemorus*), strawberry (*Fragaria ananassa* Sengana Sengana), blackcurrant (*Ribes nigrum*, var. Öjeby) and sea buckthorn berry (*Hippophae rhamnoides*) showed antimicrobial activity against selected Gram-positive and Gram-negative bacteria, including *Staphylococcus* and *Salmonella*, which were the most sensitive to the berries (Puupponen-Pimiä et al., 2005). *Salmonella* is a well-known foodborne pathogen and considered the leading bacterial cause of acute gastroenteritis (Majowicz et al., 2010; Rivera-Chávez et al., 2016).

#### **1.3.3.2 Cherries**

Cherries contain a high level of dietary fibers and bioactive compounds, such as polyphenols (Budak, 2017; Mikulic-Petkovsek, Stampar, Veberic, & Sircelj, 2016; Redondo,

Arias, Oria, & Venturini, 2017). The impact of cherries on gut microbiota composition and diversity has been receiving attention in recent years (**Table 1. 2**). A recent study investigated the impact of tart cherries and tart cherry juice on the human gut microbiota composition *in vitro* and *in vivo*. In the *in vitro* study, fermentation of tart cherry powder and tart cherry juice significantly increased the abundance of *Bacteroides*, *Collinsella*, *Veillonella*, *Bilophila*, *Escherichia*, *Clostridium* XIVa, and Enterobacteriaceae (Mayta-Apaza et al., 2018). In the *in vivo* study, tart cherry juice intake influenced the gut microbiota profile among healthy human volunteers. Individuals with an initial high abundance of *Bacteroides* showed a decrease in the abundance of *Bifidobacterium*, *Bacteroides*, *Parabacteroides*, and *Alistipes* after the tart cherry intervention, while *Lactobacillus*, Lachnospiraceae, *Ruminococcus*, *Streptococcus*, and *Collinsella* were increased. Interestingly, the treatment showed an opposite effect on the gut microbiota profile in individuals with an initial low abundance of *Bacteroides*. Individuals with low-*Bacteroides* abundance showed induced levels of *Bifidobacterium*, *Bacteroides*, and *Lactobacillus* (numerical) along with reduced levels of Lachnospiraceae, *Collinsella*, *Ruminococcus*, *Streptococcus*, *Dialister*, *Blautia*, and *Roseburia* (Mayta-Apaza et al., 2018). In another study, dark sweet cherries (*Prunus avium*) modulated the gut bacterial composition in obese diabetic *db/db* mice by increasing the abundance of *Akkermansia* spp. and fiber-degraders (the S24-7 family) and decreasing the abundance of *Lactobacillus* and Enterobacteriaceae (Garcia-Mazcorro, Lage, et al., 2018). In a more recent study, the increasing concentration of sweet and tart cherry juices resulted in significant growth of *Barnesiella* and *Akkermansia* in healthy mice (Al Othaim, Marasini, & Carbonero, 2020).

### 1.3.3.3 Mangoes

Mangoes exhibit a wide range of health benefits due to the high content of phytochemicals such as polyphenols, carotenoids, vitamin E, and vitamin C with excellent antioxidant properties. Additionally, mangoes have a high content of dietary fiber; the pulp alone contains approximately 1.3–3.8g/100g total fiber (Ajila, Naidu, Bhat, & Rao, 2007; Gondi, Basha, Bhaskar, Salimath, & Prasada Rao, 2015; Masibo & He, 2009). Several studies have investigated the impact of mangoes on preventing disease and metabolic disorders through, for example, inhibiting adipogenesis, ameliorating hyperglycemia and hyperlipidemia, reducing blood glucose, and preventing the growth of colonic cancer cells (Evans et al., 2014; Gondi et al., 2015; Noratto et al., 2010; Taing et al., 2012). The high content of dietary fiber and polyphenols make mangoes promising candidate for modulating the gut microbiota profile (Table 1. 3).

A study published in 2016 showed the ability of mango supplementation to restore the relative abundance of the beneficial genera *Bifidobacteria*, *Akkermansia*, and *Aldercreutzia* in the mice previously exposed to a high-fat diet. Additionally, mango supplementation increased the relative abundance of *Ruminococcus* and reduced the abundance of *Bacteroides* and *Parabacteroides* despite the high-fat feeding (Ojo et al., 2016). *Aldercreutzia* was characterized to produce equol, an antioxidant from dietary isoflavonoids (Maruo, Sakamoto, Ito, Toda, & Benno, 2008). A more recent study showed that mango intake caused a trend toward increasing the relative abundance of *Lactococcus lactis*, a tannase-producing bacterium. It reduced the relative abundance of *Clostridium leptum*, an endotoxin-producing bacterium, and *Bacteroides thetaiotaomicron*, an associated bacterium with obesity (Barnes et al., 2019).

#### 1.3.3.4 Apples

Apples are a rich source of bioactive compounds such as dietary polyphenols and dietary fiber, mainly pectin, which is the primary soluble fiber in apples. Pectin resists degradation by gastric acid and intestinal enzymes and passes to the large intestine where it is subjected to fermentation activity by the bacterial residents. Therefore, pectin is known to have a positive impact on gut microbiota composition and diversity and promoting health (Chung et al., 2017; Denis et al., 2016; Koutsos, Tuohy, & Lovegrove, 2015; Masumoto et al., 2016). Pectin has a prebiotic activity and has been shown to enhance beneficial bacteria such as *F. prausnitzii* (Chung et al., 2017).

To the best of our knowledge, a few studies have investigated the impact of whole apple consumption on gut microbiota (**Table 1. 4**). Three commercial apple varieties, Renetta Canada, Golden Delicious, and Pink Lady, modulated the gut microbiota composition *in vitro*. At the phylum level, all the apple varieties significantly increased Actinobacteria proportion with no effect on Firmicutes proportion. Only Renetta Canada and Golden Delicious varieties significantly increased the abundance of Proteobacteria and significantly decreased the abundance of Bacteroidetes. At the genus level, the population of *Bifidobacterium* spp. was significantly increased after the administration of Renetta Canada and Golden Delicious. Renetta Canada administration significantly increased the abundance of *F. prausnitzii*, unlike Golden Delicious and Pink Lady, which showed no impact on *F. prausnitzii* population (Koutsos et al., 2017).

*In vivo*, apple consumption modulated the gut microbiota composition in rats fed whole apples for four and 14 weeks. In the short-term intervention, *Bacteroidetes* level was decreased

after the apple supplementation (Licht et al., 2010). In a more recent *in vivo* study, apple supplementation impacted the Firmicutes/Bacteroidetes ratio by increasing Firmicutes population and decreasing Bacteroidetes in rats fed a high-fat diet compared to the control group. Additionally, apple supplementation increased the abundance of *Blautia* in rats fed a low-fat diet compared to the control group. At lower taxonomic levels, apple supplementation significantly increased the abundance of an unknown group of Clostridiales, with a decrease in the abundance of Bacteroidaceae (Garcia-Mazcorro et al., 2019). In mice fed a high fat diet, whole apples increased the relative abundance of *Akkermansia*, whereas the apple peel supplementation increased the level of *Bacteroides* (Elkahoui, Levin, Bartley, Yokoyama, & Friedman, 2019).

In humans, apple intake significantly increased the abundance of *Bifidobacteria* with a suggestive increase in the abundance of *Lactobacillus*, *Streptococcus*, and *Enterococcus*. A significant reduction was observed in the abundance of Enterobacteriaceae and Lecithinase-positive clostridia, including *C. perfringens* and a suggestive decrease in the *Pseudomonas* abundance (Shinohara, Ohashi, Kawasumi, Terada, & Fujisawa, 2010). Unfortunately, this study used culture techniques to investigate the potential modulation on gut microbiota, which gives a limited description of microbial communities. Another human study showed that the whole apple consumption lowered fecal pH and resulted in differences in the denaturing gradient gel electrophoresis (DGGE) profile (Ravn-Haren et al., 2013). However, the specific modification of the gut microbiota population was not confirmed.

#### **1.3.3.5 Grapes**

Grapes are rich in bioactive polyphenolic compounds, including flavonoids, such as anthocyanins and flavan-3-ols, and non-flavonoids, such as stilbenes (Nash et al., 2018). Due to

the anti-inflammatory and antioxidant properties, grape consumption has a beneficial effect on intestinal or systemic inflammation including reducing adiposity and markers of inflammation as well as improving glucose tolerance (Collins et al., 2016; Van Hul et al., 2018). Unfortunately, a limited number of studies assess the impact of whole grapes on the gut microbiota structure and composition (**Table 1. 5**). The vast majority of grape-based studies focused on phenolic compound extracts from grape pomace, seed, or juice (Anonye, 2017; W. Liu et al., 2017; F. Lu et al., 2019; Van Hul et al., 2018). This review will address papers that used whole grapes to investigate the potential impact on modulating the gut microbiota profile.

The grape supplementation decreased the relative abundance of sulfidogenic bacteria *Desulfobacter* spp. and *Bilophila wadsworthia* in mice fed the high-fat diet compared to control groups (Baldwin et al., 2016). Hydrogen sulfide, produced by sulfidogenic bacteria, triggers pro-inflammatory pathways and hyperproliferation and may cause colorectal cancer and other intestinal disorders, including inflammatory bowel disease and irritable bowel syndrome (Feng, Stams, De Vos, & Sánchez-Andrea, 2017; Z. Feng et al., 2017). The results also showed a suggestive increase in the abundance of *A. muciniphila*, a mucin-degrading bacterium, in mice fed the high-fat diet compared to control groups. Additionally, *Allobaculum* abundance was increased in mice fed the high-fat diet with the grape supplementation compared to the high-fat diet control group (Baldwin et al., 2016). A follow-up study also investigated the impact of grape intake on composition gut microbiota in mice fed a high-fat diet. Unlike the previous study, this study used a higher level of dietary fat and a longer intervention time-period, 16 vs. 10 weeks. In this study, the whole-grape supplementation did not have the same positive outcomes in the gut microbiota modulation in mice as in the previous study (Collins et al., 2016). More recently, an *in vivo* study showed that the supplementation of whole grapes increased the relative abundance

of various taxa belonging to the Lachnospiraceae family in mice under healthy and diseased conditions (Yiying Zhao, Nakatsu, & Jiang, 2019).

A randomized, crossover, controlled intervention study showed that red wine and de-alcoholized red wine significantly changes the gut microbiota composition in healthy volunteers. The relative abundance of *Bifidobacterium*, *Enterococcus*, *Eggerthella lenta*, the *Blautia coccooides–Eubacterium rectale* group, and the phylum Fusobacteria were significantly increased after the red wine and de-alcoholized red wine compared to the baseline (Queipo-Ortuño et al., 2012). The *Blautia coccooides-Eubacterium rectale* group, *Enterococcus* spp., and *Bifidobacterium* spp. are generally considered to be beneficial due to their anti-inflammatory and neoplastic properties (Dueñas et al., 2015). Another randomized, crossover, controlled intervention study evaluated the modulation of red wine and de-alcoholized red wine on gut microbiota profiles in obese individuals diagnosed with metabolic syndrome (MetS) (Moreno-Indias et al., 2016). The red wine and de-alcoholized red wine intake significantly increased the abundance of gut mucosal barrier protectors, such as *Bifidobacterium* spp. and *Lactobacillus* spp. Furthermore, a positive impact of the red wine and de-alcoholized red wine intake on the growth of the butyrate producers, namely the *Blautia coccooides–Eubacterium rectale* group, *Faecalibacterium prausnitzii*, and *Roseburia* was observed. The red wine and de-alcoholized red wine also showed an inhibitory effect on the *Clostridium histolyticum* group, pathogens associated with the progression of colon cancer, and the onset of IBD (Moreno-Indias et al., 2016).



#### 1.3.3.6 Kiwifruits

Kiwifruits are rich in nutrients, such as dietary fiber and phytochemicals, which have prebiotic effects on intestinal bacteria (Leontowicz et al., 2016). Kiwifruits have shown to modify gut microbial ecology (**Table 1. 6**). Green and gold kiwifruits increased the relative abundances of *Bifidobacterium* spp. and *Bacteroides* spp. *in vitro* (Blatchford et al., 2015). Another study showed that green and gold kiwifruits increased the total intestinal bacteria composition in rats compared to the control diet. Both green and gold kiwifruit significantly increased the abundance of Lachnospiraceae, a family that contains members among the primary producers of SCFAs and other members found to be associated with diseases (Paturi, Butts, Bentley-Hewitt, & Ansell, 2014; Vacca et al., 2020). The relative abundance of the *Bacteroides*–*Prevotella*–*Porphyromonas* group and *Enterococcus* spp. were induced with gold kiwifruit supplementation (Paturi et al., 2014). In a pig model, dietary treatment with green kiwifruits significantly increased the number of total bacteria with a significant increase in the abundance of *Bacteroides* as well as a significant decrease of *E. coli* abundance. The ratio of *Lactobacillus*/Enterobacteria was significantly increased after kiwifruit supplementation compared to the control (K. S. Han, Balan, Molist Gasa, & Boland, 2011). Therefore, kiwifruits may play a role in enhancing the relative abundance of beneficial bacteria such as *Bifidobacterium* and *Lactobacillus* as well as inhibiting the growth of opportunistic pathogens such as *E. coli* and *Enterobacteria*.

#### 1.3.3.7 Oranges

The importance of oranges in promoting general health has long been established due to the richness in nutrients including vitamin C, dietary fiber, and other bioactive compounds such

as polyphenols. Flavonoids are among the most dominant polyphenols in oranges and orange juice (Stella, Ferrarezi, dos Santos, & Monteiro, 2011; Stinco et al., 2012). The bioactive components present in orange juice are associated with the gut microbiota metabolism (Laparra & Sanz, 2010; Pereira-Caro et al., 2015; Pereira-Caro et al., 2015). Numerous scientific papers have investigated the impact of orange juice in modulation gut microbiota ecology (**Table 1. 7**). An *in vitro* study reported that fresh orange juice induced the growth of *Bifidobacterium* spp., *Lactobacillus* spp., *Enterococcus* spp., and *Clostridium* spp. while the juice reduced the level of the *Enterobacteria*. Pasteurized orange juice showed a similar effect in gut microbiota by increasing the growth of *Lactobacillus* spp. and reducing *Enterobacteria* (Duque, Monteiro, Adorno, Sakamoto, & Sivieri, 2016). Two different orange juices from *Citrus sinensis* variety, cv. “Cara Cara” and cv. “Bahia”, influenced the composition of gut microbiota in humans in a short-term intervention. Cara Cara orange juice increased the relative abundance of Mogibacteriaceae and Tissierellaceae families while the relative abundance of Odoribacteraceae family and the *Odoribacter* genus were decreased. In the Bahia group, Mogibacteriaceae, Enterococcaceae, and Veillonellaceae families were increased, whereas the Ruminococcaceae family was decreased after the juice consumption. The Porphyromonadaceae, Odoribacteraceae, Christensenellaceae, Enterobacteriaceae, Lachnospiraceae, and Ruminococcaceae families and Parabacteroides and Butyrivibrio genera were increased in Cara Cara orange juice compared to the control group. The relative abundance of the Coriobacteriaceae family and *Adlercreutzia* genus, Enterococcaceae family and *Enterococcus* genus, Clostridiaceae family and *Clostridium* genus, Ruminococcaceae family and *Anaerotruncus* genus, and Pasteurellaceae family were significantly higher in Bahia orange juice compared to the control group (Brasili et al., 2019). A long-term intervention, eight weeks, of orange juice positively modulated the gut microbiota

population in healthy volunteers by significantly increasing the relative abundance of *Bifidobacterium* spp. and *Lactobacillus* spp. (Lima et al., 2019).

#### **1.3.3.8 Pomegranates**

Pomegranate fruits and products are widely consumed and recognized for their health benefits (Z. Li, Henning, et al., 2015). The health benefits are attributed to the presence of phenolic compounds, mainly ellagitannins (Henning et al., 2017; Z. Li, Henning, et al., 2015). Several studies have shown the impact of pomegranate fruits and products on the gut microbiota composition (**Table 1. 8**). The residual materials, which are left after the pomegranate juice production, are pressed and extracted with water to make the pomegranate by-product (POMx) which is commercially available as a dietary supplement (Bialonska et al., 2010). POMx was reported to increase the total bacteria and the level of *Bifidobacterium* spp. and *Lactobacillus* spp. *in vitro* (Bialonska et al., 2010). In agreement with the previous study, another *in vitro* study showed that the pomegranate (POM) juice and POMx increased the level of *Bifidobacterium* and *Lactobacillus*. *B. fragilis* group, *clostridia*, and Enterobacteriaceae was significantly inhibited by the POM juice and POMx in a dose–response manner (Z. Li, Summanen, et al., 2015).

In an *in vivo* study, an increase in the relative abundance of Ruminococcaceae, a family that has been reported to have anti-inflammatory activities (X. Zhang et al., 2012), was observed in rats supplemented with the pomegranate beverage compared to the control group (Kim et al., 2017). Moreover, pomegranate juice abolished the dysbiosis in gut microbiota caused by acrolein in apoE<sup>−/−</sup> mice (Rom et al., 2017).

POMx supplementation significantly increased the relative abundance of the commensal bacterium *A. muciniphila* in healthy volunteers (Henning et al., 2017). Furthermore, POMx

supplementation significantly increased the Actinobacteria population and decreased the Firmicutes population in healthy individuals. At the genus level, *Lactobacillus*, *Akkermansia*, *Prevotella*, *Butyrivibrio*, *Enterobacter*, *Escherichia*, *Serratia*, and *Veillonella* were increased after POMx intake, whereas *Collinsella* significantly decreased (Z. Li, Henning, et al., 2015).

#### **1.3.3.9 Avocados**

Avocados are rich in dietary fiber, monounsaturated fat, and a selection of essential phytochemicals, including lutein (Dreher & Davenport, 2013; Q.-Y. Lu et al., 2009). A few studies have investigated the potential impact of avocado administration on shaping the intestinal microbiota (**Table 1. 9**). A recent study showed that the Firmicutes/Bacteroidetes ratio was maintained by avocado consumption in healthy overweight and obese individuals. A significant increase was observed in the abundance of Veillonellaceae and Sutterellaceae with a suggestive increase in the abundance of Ruminococcaceae and its genera *Ruminococcus* and Prevotellaceae and its genera *Prevotella* after the avocado intervention. The genera *Bacteroides* and *Methanosphaera* were decreased in the avocado group with a significant increase in the abundance of *Dialister*, *Sutterella*, *Bilophila*, *Holdemanella*, *Herbaspirillum*, and *Acetivibrio*, of which several have the capability to ferment fiber (Bang et al., 2018; Foughse, Gänzle, Beattie, Vasanthan, & Zijlstra, 2017; Henning et al., 2019). A randomized controlled trial study showed that the avocado consumption significantly increased the relative abundances of *Lachnospira* and suggestively elevated the abundance of *Faecalibacterium* in overweight or obese adults (Thompson, Edwards, Reeser, Khan, & Holscher, 2019). *Lachnospira* and *Faecalibacterium* are thought to play a role in maintaining gut homeostasis and epithelial integrity due to their ability to produce butyrate (Ferrario et al., 2014; Vital, Howe, & Tiedje, 2014). *Lachnospira* belongs to Lachnospiraceae, which has been reported to restrain the growth of the opportunistic pathogen,

*Clostridium difficile* (Bhaduri, Dubey, & Basu, 2019; Reeves, Koenigsknecht, Bergin, & Young, 2012).

#### **1.3.3.10 Bitter melons**

Bitter melon fruit is an excellent source of physiologically active compounds, including dietary fiber and polyphenols, mainly catechin (Bai, Zhu, & Dong, 2016; Hwang, 2018; Ri, Magbagbeola, Akinwande, & Okunowo, 2010). A good number of studies reported the impact of the bitter melon fruit in hypoglycemic, hypolipidemic, and improving insulin resistance in the obese (Bao et al., 2013; Juan, Ying, & Ying, 2017; Xu et al., 2014; S. J. Yang et al., 2015; Zhu, Bai, Zhang, Xiao, & Dong, 2016). However, a few studies have investigated the potential impact on gut microbiota ecology, most of which were conducted with obesity-induced diets (Bai et al., 2016; Nerurkar, Orias, Soares, Kumar, & Nerurkar, 2019; Zhu et al., 2016) (**Table 1. 10**). In obese rats, the level of the families Desulfovibrionaceae and Enterobacteriaceae were reduced by the bitter melon supplemented diet compared to the high-fat diet, whereas the Odoribacteraceae, a butyrate producer, level was significantly increased after the dietary intervention (Bai et al., 2016). The association of Enterobacteriaceae and Desulfovibrionaceae families with a systemic inflammatory state have been reported (Hakansson & Molin, 2011; Zeng et al., 2017). At the genus level, the abundance of *Escherichia*, which contains opportunistic pathogens, was significantly decreased in the group supplemented with bitter melons compared to the high-fat diet group. This was coordinated with an increase in the abundance of the butyrate-producing genera *Faecalibacterium*, *Allobaculum*, *Butyricimonas*, and *Odoribacter* (Bai et al., 2016). A follow-up study showed that the bitter melon powder supplementation increased the Verrucomicrobia population in obese rats fed a high-fat diet. At the genus level, the bitter melon powder treatment significantly increased the relative abundance of *Blautia*, *Anaerotruncus*,

*Lactococcus*, *Allobaculum*, and *Oceanobacillus* and decreased *Prevotella* and *Anaeroplasma* (Bai, Zhu, & Dong, 2018). *Blautia* and *Allobaculum* are SCFAs producing-bacteria (Greetham et al., 2004; Park, Kim, Roh, & Bae, 2012). Bitter melon juice improved the ratio of Firmicutes/Bacteroidetes with a reduction in Actinobacteria abundance in mice fed a high-fat diet compared to the high-fat control group. Additionally, an increase in the abundance of Clostridiaceae, Lactobacillales, and *Lactobacillus* and a decrease in *Bacteroides* abundance was reported after bitter melon juice administration compared to the control group (Nerurkar et al., 2019). Lactobacillales is negatively correlated with colonic injury as well (Ritchie et al., 2015).

#### **1.3.3.11 Broccoli**

Broccoli is a good source of dietary fiber and phytochemicals, including glucosinolates and polyphenols (X. Liu et al., 2017). Glucosinolates are relatively non bioactive in their native state. However, the gut microbial residence can hydrolyze glucosinolates to form the bioactive glucosinolates derivatives, mainly isothiocyanates (Angelino et al., 2015; Tian, Liu, Lei, Zhang, & Shan, 2018). Several studies showed that broccoli consumption could alter the composition of gut microbiota (**Table 1. 11**). Cooked broccoli supplementation significantly increased the relative abundance of the genera *Akkermansia* and *Oscillospira* as well as the family Ruminococcaceae. The broccoli consumption significantly decreased the relative abundance of *Blautia*, *Clostridium*, and *Dorea* in rats (X. Liu et al., 2017). A controlled feeding and randomized crossover study showed that daily consumption of cooked broccoli in human volunteers increased the relative abundance of Bacteroidetes and its genus *Bacteroides* and decreased the relative abundance in Firmicutes (Kaczmarek et al., 2019). An *in vivo* study reported that the broccoli supplementation decreased the abundance of *C. perfringens* and *E. coli*, bacteria that are known to be associated with IBD, in *mdr1a*<sup>-/-</sup> mice, a model for IBD.

The broccoli-supplemented diet also lowered the abundance of *Enterococcus* spp., *Lactobacillus* spp., and *F. prausnitzii* compared to the control diet (Paturi et al., 2012).

#### **1.3.3.12 Mushrooms**

Mushrooms contain a high content of bioactive components, including dietary fiber and polyphenols (Amirullah, Abidin, & Abdullah, 2018; Sánchez, 2017). Beside the anti-inflammation and antioxidant effects, whole mushroom supplementation has been reported to influence the gut microbial ecosystem in animal models (Iuchi et al., 2015; S. P. Therkelsen, G. Hetland, T. Lyberg, I. Lygren, & E. Johnson, 2016; Stig Palm Therkelsen, Geir Hetland, Torstein Lyberg, Idar Lygren, & Egil Johnson, 2016; Vetvicka & Vetvickova, 2014) (**Table 1. 11**). White button mushrooms increased the Bacteroidetes population and decreased Firmicutes in mice. At the class level, the mushroom supplementation increased the abundance of Mollicutes, Proteobacteria sub-classes Delta, Gamma, and Epsilon, and Verrucomicrobia and decreased the abundance of Erysipelotrichi, Deferribacteres, as well as the potentially pathogenic bacteria Clostridia (Varshney et al., 2013). The dietary supplementation of white button mushrooms altered the composition of gut microbiota in pigs (Solano-Aguilar et al., 2018). A mushroom treatment enriched the abundance of the family Ruminococcaceae and its genera *Oscillibacter* and *Butyricicoccus* and the family Lachnospiraceae and its genera *Fusicatenibacter*, *Robinsoniella*, and *Eisenbergiella*, which have a role in plant fibers degradation and SCFAs production (Chassard, Delmas, Robert, Lawson, & Bernalier-Donadille, 2012; Meehan & Beiko, 2014). In addition, the white button supplementation increased the abundance of *Fusicatenibacter*, *Eisenbergiella*, *Lachnobacterium*, and *Robinsoniella* which were found to be associated with many health outcomes, including lowering levels of cholesterol, correcting the dysbiosis caused by obesity, and improving outcomes for patients with primary biliary cirrhosis

(Lv et al., 2016; Prieto et al., 2018; Togo et al., 2016). Feeding pigs with mushrooms reduced the relative abundance of *Bifidobacterium* and increased the relative abundance of *Prevotella*, which was found to be associated with improving glucose metabolism (Kovatcheva-Datchary et al., 2015). In contrast, the *Armillariella tabescens* administration promoted *Lactobacillus* spp. and *Bifidobacterium* spp. and inhibited *E.coli* growth (Chen et al., 2017). A variety of mushrooms increased the Firmicutes and Actinobacteria population and decreased the population of Bacteroidetes and Proteobacteria in mice fed a high-fat diet compared to the high-fat diet group. Furthermore, there was a tendency toward an increased relative abundance of SCFAs-producing bacteria, *Allobaculum*, *Bifidobacterium*, and *Ruminococcus*, as well as lactic acid-producing bacteria, *Lactobacillus*, *Lactococcus*, and *Streptococcus*, in mice fed mushrooms compared to the high-fat diet group. Additionally, *Adlercreutzia* and *Sutterella* showed a suggestive increase after the mushrooms intake, while a suggestive decrease was observed in *Bacteroides*, *Prevotella*, *Mucispirillum*, *Dorea*, *Roseburia*, *Anaerotruncus*, *Oscillospira*, *Escherichiam*, and *Akkermansia* (Shimizu, Mori, Ouchi, Kushida, & Tsuduki, 2018). More recently, *Pleurotus eryngii* mushrooms significantly increased the richness and diversity of gut microbiota and partially reversed the dysbiosis induced by dextran-sodium-sulfate (DSS)-treatment in mice (Q. Hu et al., 2019). *P. eryngii* administration significantly increased the relative abundance of S24-7 and Odoribacteraceae at the family level, and *Adlercreutzia*, *Akkermanisa*, *Lactobacillus*, *Anaerostipes*, and *Allobaculum* at the genus level compared to the DSS group. *Adlercreutzia* reduction was observed in patients with IBD (Shaw et al., 2016). *P. eryngii* supplementation also reduced the abundance of Actinobacteria, Mollicutes, Desulfovibrionaceae, Enterococcaceae, *Turicibacter*, *Dorea*, rc4-4, *Bacteroides*, and *Prvotella* compared to the DSS group. It has been reported that the family Desulfovibrionacea, a sulfate-reducing bacteria, is a potential endotoxin



producer and might be associated with the low grade and chronic inflammation (Xiao et al., 2014).

#### ***1.3.3.13 Jerusalem artichokes***

Jerusalem artichokes, also called sunchoke, sunroot, or earth apple, are rich in dietary fiber, mainly Inulin-type fructans. The inulin content in Jerusalem artichokes ranges from 7 to 30% for fresh weight or from 60 to 85.5% for dry weight (Aduldecha et al., 2016; Samal, Chaturvedi, & Pattanaik, 2017). A few studies have investigated the potential effect of Jerusalem artichoke on gut microbiota composition (**Table 1. 11**). Results from *an in vivo* study showed that Jerusalem artichoke supplementation increased the level of *Lactobacillus* spp. and *Bifidobacterium* spp. in the cecal, colonic, and rectal digesta in healthy rats. The Jerusalem artichoke supplementation decreased the Coliforms population in rats (Samal et al., 2017). More studies, especially *in vivo* studies, need to be done to assess the potential impact of Jerusalem artichoke on the gut microbiota composition and diversity.

### **1.4 Conclusion**

Fruit and vegetable consumption profoundly modifies and shapes the gut microbiota ecosystem due to the intense content of dietary fibers and bioactive compounds, including polyphenols. Several health outcomes have been linked to reshaping the gut microbiota and their associated functions by fruit and vegetable supplementation, including promoting the host's general health, reducing the risk of several diseases and metabolic disorders. Phytochemicals and fibers are the compounds found in fruits and vegetables that impact the intestinal microbiota composition and diversity. A majority of nutritional and microbiological researches have focused on investigating the effect of extracted compounds on gut microbiota profiles instead of the

whole foods. Berries have been more commonly used to investigate the impact of whole foods on gut microbiota. More studies of whole fruits and vegetables, and especially in combination, are needed due to potential synergies between compounds and interactions with the whole fruits or vegetables matrices.

**Table 1. 1:** Modulation of the gut microbiota by berries.

**HTS:** High throughput sequencing; **qPCR:** Quantitative polymerase chain reaction; **DGGE:** Denaturing gradient gel electrophoresis; **FISH:** Fluorescent in situ hybridization; **SHIME:** Simulator of the human intestinal microbial ecosystem; **PCR:** polymerase chain reaction.

Berries	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Strawberry	Animal study (diabetic mice)	2.35% of dried strawberry powder	10 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↓ Verrucomicrobia <u>Genus level:</u> ↑ <i>Bifidobacterium</i> ↓ <i>Bacteroides</i> and <i>Akkermansia</i> ↔ <i>Dehalobacterium</i> , <i>Dorea</i> , <i>Lactobacillus</i> , and <i>Turicibacter</i>	(Petersen et al., 2019)
Strawberry	Animal study (colitic mice)	0, 2.5, or 5% (w/w) freeze-dried strawberry powder	37 days	HTS (Illumina Miseq)	<u>Genus level:</u> ↑ <i>Bifidobacterium</i> and <i>Lactobacillus</i> ↓ <i>Akkermansia</i> , <i>Dorea</i> , <i>Bilophila</i> , and <i>Bacteroides</i>	(Y. Han et al., 2019)
Cranberry	<i>In vitro</i> (Dynamic Fermentation Models, SHIME)	1 mg/mL of whole cranberry powder	every 24 hours for 5 days	Ion Torrent PGM	<u>Family level:</u> ↑ Porphyromonadaceae and Bacteroidaceae ↓ Enterobacteriaceae ↔ Ruminococcaceae and Lachnospiraceae	(O'Connor et al., 2019)
Cranberry	Animal study (colitic mice)	1.5% (w/w) freeze-dried whole cranberry powder	37 days	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Verrucomicrobia and Proteobacteria ↓ Actinobacteria <u>Genus level:</u> ↑ <i>Bifidobacterium</i> and <i>Lactobacillus</i> ↓ <i>Bilophila</i> , <i>Sutterella</i> , and <i>Akkermansia</i>	(Cai et al., 2019)
Cranberry	Human study (healthy individuals)	30 g of dried cranberry powder	5 days	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Bacteroidetes ↓ Firmicutes <u>Genus level:</u> ↑ <i>Lachnospira</i> and <i>Anaerostipes</i> ↓ <i>Oribacterium</i>	(Rodríguez-Morató et al., 2018)
Blueberry	Animal study (mice, inflammation bowel disease model)	10% of dried blueberry powder	21 weeks	qPCR	↔ <i>Bacteroides-Prevotella</i> - <i>Porphyromonas</i> , <i>Bifidobacterium</i> , <i>Bacteroides vulgatus</i> , and <i>Faecalibacterium prausnitzii</i> ↓ <i>Clostridium perfringens</i> , <i>Enterococcus</i> , <i>Lactobacillus</i> , and <i>Escherichia coli</i>	(Paturi et al., 2012)

**Table 1. 1 (Cont.)**

Berries	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Blueberry	Animal study (healthy mice)	5% of dried blueberry powder	4 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Bacteroidetes and Tenericutes ↓ Firmicutes <u>Family level:</u> ↓ Deferribacteres	(Wankhade et al., 2019)
Blueberry	Animal study (obese mice)	4% of freeze-dried whole blueberry powder	8 weeks	HTS (Illumina Hiseq)	↑ <i>Turicibacter</i> sp. H121	(Morissette et al., 2020)
Blueberry	Animal study (healthy rats)	8% of dried blueberry powder	6 weeks	HTS (Illumina Hiseq)	<u>Phylum level:</u> ↑ Actinobacteria <u>order level:</u> ↑ Actinomycetale <u>Family level:</u> ↑ Bifidobacteriaceae and Coriobacteriaceae <u>Genus level:</u> ↓ <i>Lactobacillus</i> and <i>Enterococcus</i>	(Lacombe et al., 2013)
wild blueberry ( <i>Vaccinium angustifolium</i> )	Human study (healthy male)	25 g of wild blueberry powder in 250 mL of water	6 weeks	qPCR	↑ <i>Bifidobacterium</i> , <i>Lactobacillus acidophilus</i> ↔ <i>Bacteroides</i> , <i>Prevotella</i> , <i>Enterococcus</i> , and <i>Clostridium coccoides</i>	(Vendrame et al., 2011)
wild blueberry ( <i>Vaccinium angustifolium</i> )	Human study (healthy male)	25 g of wild blueberry powder in 250 mL of water	6 weeks	qPCR	↑ <i>Bifidobacterium longum</i> subsp. <i>Infantis</i> ↔ <i>Bifidobacterium adolescentis</i> , <i>B. longum</i> subsp. <i>longum</i> , <i>B. catenulatum</i> , <i>B. breve</i> , and <i>B. bifidum</i>	(Guglielmetti et al., 2013)
Black raspberries	Animal study (healthy rats)	5% of dried black raspberries powder	6 weeks	Pyrosequencing	↑ <i>Akkermansia</i> , <i>Desulfovibrio</i> , and <i>Anaerostipes</i>	(Pan et al., 2017)
Black raspberries	Animal study (specific-pathogen free mice)	10% of dried black raspberries powder	7 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Bacteroidetes and Verrucomicrobia ↓ Firmicutes <u>Species level:</u> ↑ <i>Akkermansia muciniphila</i>	(Tu et al., 2018)
Black raspberries	Animal study (healthy mice)	10% of dried black raspberries powder	6 weeks	HTS (Illumina Miseq)	↑ Bacteroidetes, <i>Barnesiella</i> , and <i>Turicibacter</i> ↓ Firmicutes, <i>Clostridium</i> , and <i>Lactobacillus</i>	(Gu et al., 2019)
Red raspberries	Animal study (obese diabetic <i>db/db</i> mice)	5.3% of dried red raspberry	8 weeks	qPCR	<u>Order level:</u> ↓ Bacteroidales <u>Family level:</u> ↑ Lachnospiraceae ↓ Enterobacteriaceae and S24-7 <u>Genus level:</u> ↓ <i>Lactobacillus</i> and <i>Odoribacter</i>	(Garcia-Mazcorro, Pedreschi, et al., 2018)

**Table 1. 1 (Cont.)**

Berries	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Strawberry, bilberry, red raspberry, cloudberry, blackcurrant, and sea buckthorn berry	<i>In vitro</i> (batch-culture fermentation)	2 and 10 mg ml <sup>-1</sup> of dried berry powder	24 hours	Culture methods	↓ <i>Salmonella enterica</i> , and <i>Staphylococcus aureus</i>	(Puuponen-Pimiä et al., 2005)
Sea buckthorn berry	<i>In vitro</i> (batch-culture fermentation)	250 mg lyophilized fraction of small intestine digested berries	72 hours	PCR-DGGE	↑ <i>Bifidobacteria</i> , <i>Bacteroides/Prevotella</i> group, and Lactic acid bacteria	(Attri et al., 2018)
Sea buckthorn berry	<i>In vitro</i> (batch-culture fermentation)	10% of berry juice	1 week (continuous gut model)	PCR-DGGE	↑ <i>Bifidobacteria</i> , <i>Lactobacilli</i> , and <i>Bacteroides/Prevotella</i>	(Attri & Goel, 2018)
Juçara ( <i>Euterpe edulis</i> )	<i>In vitro</i> (batch-culture fermentation)	1% of dried juçara pulp powder	24 hours	FISH	↑ <i>Bifidobacterium</i> , <i>Clostridium clusters XIVa</i> , <i>Bacteroides</i> , <i>Eubacterium rectale</i> , and <i>Prevotella</i>	(Guergoletto et al., 2016)
Juçara ( <i>Euterpe edulis</i> )	Animal study (rats fed a high-fat diet)	0.5 and 0.25% of dried berry powder	7 days	qPCR	↑ <i>Bifidobacterium</i>	(Jamar et al., 2018)
<i>Lonicera cerula</i> L. berry	Animal study (mice fed a high-fat diet)	1% of dried <i>Lonicera cerula</i> L. berry powder	45 Days	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Bacteroidetes, Parabacteroides, and Verrucomicrobia ↓ Firmicutes <u>Genus level:</u> ↑ <i>Akkermansia</i> , <i>Bacteroides</i> , and <i>Parabacteroides</i> ↓ <i>Staphylococcus</i> , <i>Lactobacillus</i> , <i>Oscillospira</i> , and <i>Ruminococcus</i>	(S. Wu et al., 2018)
Goji berry ( <i>Lycium barbarum</i> )	Animal study (colitis mice model)	1% of dried Goji berry powder	10 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Actinobacteria <u>Family level:</u> ↑ Lachnospiraceae and Ruminococcaceae <u>Genus level:</u> ↑ <i>Bifidobacterium</i> , <i>Roseburia</i> , and <i>Faecalibacterium prausnitzii</i>	(Kang et al., 2018)

**Table 1. 1 (Cont.)**

Berries	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Lingonberry	Animal study (mice fed a high-fat diet)	44% of dried Lingonberry powder	8 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Bacteroidetes ↓ Firmicutes <u>Genus level:</u> ↑ <i>Bacteroides</i> , <i>Clostridium</i> , and <i>Parabacteroides</i> ↓ <i>Mucispirillum</i> and <i>Oscillospira</i> <u>Species level:</u> ↑ <i>Akkermansia muciniphila</i> , <i>Blautia producta</i> , <i>Clostridium difficile</i> , and <i>Eubacterium dolichum</i>	(Matziouridou et al., 2016)
A berry mixture (blueberries, blackberries, raspberries, Portuguese crowberry, and strawberry tree fruit)	Animal study (rats fed a salt-induced hypertension diet)	2 g of lyophilized berry mixture	9 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Bacteroidetes ↓ Firmicutes and Proteobacteria <u>Family level:</u> ↑ Bacteroidaceae, S24-7, and Ruminococcaceae ↓ Erysipelotrichaceae	(Gomes et al., 2019)
<i>Schisandra chinensis</i> fruit	Human study (obese women)	100 ml, twice a day	12 weeks	qPCR	<u>Phylum level:</u> ↑ Bacteroidetes ↓ Firmicutes <u>Genus level:</u> ↑ <i>Bifidobacterium</i> , <i>Akkermansia</i> , <i>Roseburia</i> , <i>Prevotella</i> , and <i>Bacteroides</i>	(Song et al., 2015)

**Table 1. 2:** Modulation of the gut microbiota by cherries.

**HTS:** High throughput sequencing; **qPCR:** Quantitative polymerase chain reaction.

Cherries	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Tart Cherry	<i>In vitro</i> (batch-culture fermentation)	5 ml of concentrated tart cheery or 5g of tart cherry powder	48 Hours	Culture methods	↑ <i>Bacteroides</i> , <i>Veillonella</i> , <i>Bilophila</i> , <i>Escherichia</i> , <i>Clostridium</i> XIVa, and Enterobacteriaceae	(Mayta-Apaza et al., 2018)

**Table 1. 2 (Cont.)**

Cherries	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Tart Cherry	Human study (healthy individuals)	8 oz. of tart cherry juice daily	5 days	HTS (Illumina Miseq)	<u>Individuals with initial high-<i>Bacteroides</i> level:</u> ↑ <i>Lactobacillus</i> , <i>Lachnospiraceae</i> , <i>Ruminococcus</i> , <i>Streptococcus</i> , and <i>Collinsella</i> ↓ <i>Bifidobacterium</i> , <i>Bacteroides</i> , <i>Parabacteroides</i> , and <i>Alistipes</i> <u>Individuals with initial low-<i>Bacteroides</i> level:</u> ↑ <i>Bifidobacterium</i> , <i>Bacteroides</i> , and <i>Lactobacillus</i> (numerical) ↓ <i>Lachnospiraceae</i> , <i>Collinsella</i> , <i>Ruminococcus</i> , <i>Streptococcus</i> , <i>Dialister</i> , <i>Blautia</i> , and <i>Roseburia</i>	(Mayta-Apaza et al., 2018)
Dark sweet cherry ( <i>Prunus avium</i> )	Animal study (obese diabetic <i>db/db</i> mice)	10% of dark sweet cherry powder	12 weeks	qPCR	↑ <i>Akkermansia</i> spp. and fiber-degraders (the S24-7 family) ↓ <i>Lactobacillus</i> and Enterobacteriaceae	(García-Mazcorro, Lage, et al., 2018)
Sweet and tart cherries juices	Animal study (Healthy mice)	Increasing concentration of the juices (1/20 v/v, 1/15 v/v, 1/10 v/v, 1/7 v/v, and 1/4 v/v) every 5 days	25 days	HTS (Illumina Miseq)	↑ <i>Barnesiella</i> and <i>Akkermansia</i> ↓ <i>Bacteroides</i>	(Al Othaim et al., 2020)

**Table 1. 3: Modulation of the gut microbiota by mangos.**

**HTS:** High throughput sequencing; **qPCR:** Quantitative polymerase chain reaction.

Mangos	Study type	Dosage	Treatment duration	Analytical methods	Changes in gut microbiota	Ref.
Mango	Animal study (mice fed a high-fat diet)	1% or 10% of dried mango	12 weeks	HTS (Illumina Miseq)	↑ <i>Bifidobacteria</i> , <i>Akkermansia</i> , <i>Aldercreutzia</i> , and <i>Ruminococcus</i> ↓ <i>Bacteroides</i> and <i>Parabacteroides</i>	(Ojo et al., 2016)
Mango	Human study (healthy lean and obese individuals)	400g of dried mango daily	6 weeks	qPCR	↑ <i>Lactococcus lactis</i> ↓ <i>Clostridium leptum</i> and <i>Bacteroides thetaiotaomicron</i>	(Barnes et al., 2019)

**Table 1. 4: Modulation of the gut microbiota by apples.**

**HTS:** High throughput sequencing; **DGGE:** Denaturing gradient gel electrophoresis.

Apples	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Apple	Animal study (healthy rat)	10 g/day	4 weeks	PCR-DGGE	↓ <i>Bacteroides</i>	(Licht et al., 2010)

**Table 1. 4 (Cont.)**

Apples	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Apple varieties (Renetta Canada, Golden Delicious, and Pink Lady)	<i>In vitro</i> (batch-culture fermentation)	2g of apple, a final concentration of 1% (w/v)	24 hours	HTS (Illumina Miseq) and FISH	<u>Phylum level:</u> ↑ Actinobacteria (with all apple varieties) ↑ Proteobacteria (with Renetta Canada and Golden Delicious) ↓ Bacteroidetes (with Renetta Canada and Golden Delicious) ↔ Firmicutes (with all apple varieties) <u>Genus level:</u> ↑ <i>Bifidobacterium</i> spp. (with Renetta Canada and Golden Delicious) ↑ <i>Faecalibacterium prausnitzii</i> (with Renetta Canada)	(Koutsos et al., 2017)
Apple	Animal study (rat fed a high-fat or low-fat diet)	5% of freeze dried apple	6 weeks	HTS (Illumina Miseq)	<u>Phylum level (high-fat diet group):</u> ↑ Bacteroidetes ↓ Firmicutes <u>Other levels (high-fat diet group):</u> ↑ Clostridiales ↓ Bacteroidaceae <u>Genus level (low-fat diet group):</u> ↑ <i>Blautia</i>	(Garcia-Mazcorro et al., 2019)
Apple	Animal study (mice fed a high-fat or low-fat diet)	40% of freeze dried whole RedDel apple	4 weeks	HTS (Illumina Miseq)	↑ <i>Akkermansia</i>	(Elkhoui et al., 2019)
Apple (apple peel)	Animal study (mice fed a high-fat or low-fat diet)	20% of freeze dried RedDel apple peel	4 weeks	HTS (Illumina Miseq)	↑ <i>Bacteroides</i>	(Elkhoui et al., 2019)
Apple	Human study (healthy individuals)	Two apples daily	2 weeks	Culture methods	↑ <i>Bifidobacteria</i> , <i>Lactobacillus</i> , <i>Streptococcus</i> , and <i>Enterococcus</i> . ↓ Enterobacteriaceae, Lecithinase-positive clostridia, including <i>C. perfringens</i> , and <i>Pseudomona</i>	(Shinohara et al., 2010)

**Table 1. 5:** Modulation of the gut microbiota by grapes.

**HTS:** High throughput sequencing; **DGGE:** Denaturing gradient gel electrophoresis; **PCR:** polymerase chain reaction.

Grapes	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Grape	Animal study (mice following a high fat diet)	3% or 5% powdered grapes	11 weeks	HTS (Illumina Miseq)	↑ <i>Akkermansia muciniphila</i> and <i>Allobaculum</i> ↓ <i>Desulfovibrio</i> spp. and <i>Bilophila wadsworthia</i>	(Baldwin et al., 2016)



**Table 1. 5 (Cont.)**

Grapes	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Grape	Animal study (mice fed a high-fat diet)	5% powdered grapes	16 weeks	HTS (Illumina Miseq)	No significant impact was reported	(Collins et al., 2016)
Red wine and de-alcoholized red wine	Human study (healthy individuals)	272 mL of red wine or de-alcoholized red wine daily	20 days	PCR-DGGE	<u>Phylum level:</u> ↑ Fusobacteria <u>Genus level:</u> ↑ <i>Bifidobacterium</i> and <i>Enterococcus</i> <u>Species level:</u> ↑ <i>Eggerthella lenta</i> , <i>Blautia coccoides</i> – <i>Eubacterium rectale</i> group	(Queipo-Ortuño et al., 2012)
Red wine and de-alcoholized red wine	Human study (obese individuals)	272 mL of red wine or de-alcoholized red wine daily	30 days	PCR-DGGE	<u>Phylum level:</u> ↑ Fusobacteria and Bacteroidetes ↓ Firmicutes <u>Other levels:</u> ↑ <i>Lactobacillus</i> , <i>Bifidobacterium</i> , <i>Blautia coccoides</i> – <i>Eubacterium rectale</i> group, <i>Faecalibacterium prausnitzii</i> , <i>Roseburia</i> , <i>Eggerthella lenta</i> , and <i>Prevotella</i> ↓ <i>Clostridium</i> , <i>Clostridium histolyticum</i> group, <i>Bacteroides</i> , <i>Escherichia coli</i> and <i>Enterobacter cloacae</i>	(Moreno-Indias et al., 2016)

**Table 1. 6:** Modulation of the gut microbiota by kiwifruits.

**qPCR:** Quantitative polymerase chain reaction; **DGGE:** Denaturing gradient gel electrophoresis;

**PCR:** polymerase chain reaction.

Kiwifruits	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Kiwifruit (green and gold kiwifruit)	<i>In vitro</i> (batch-culture fermentation)	1% of green or gold kiwifruit	48 hours	qPCR	↑ <i>Bifidobacterium</i> spp. and <i>Bacteroides</i> spp.	(Blatchford et al., 2015)
Kiwifruit (green kiwifruit)	Animal study (healthy rats)	10% of green kiwifruit	4 weeks	PCR-DGGE	↑ Lachnospiraceae	(Paturi et al., 2014)
Kiwifruit (green kiwifruit)	Animal study (healthy pigs)	25% of dried green kiwi	14 days	PCR-DGGE	↑ <i>Bacteroides</i> ↓ Enterobacteria and <i>Escherichia coli</i>	(K. S. Han et al., 2011)

**Table 1. 7:** Modulation of the gut microbiota by oranges.

**DGGE:** Denaturing gradient gel electrophoresis; **SHIME:** Simulator of the human intestinal microbial ecosystem.

Oranges	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Orange juice	<i>In vitro</i> (Dynamic Fermentation Models, SHIME)	105 ml of fresh orange juice	14 days	PCR-DGGE	↑ <i>Bifidobacterium</i> spp., <i>Lactobacillus</i> spp., <i>Enterococcus</i> spp., and <i>Clostridium</i> spp. ↓ <i>Enterobacteria</i>	(Duqu e et al., 2016)
	<i>In vitro</i> (Dynamic Fermentation Models, SHIME)	105 ml of pasteurized orange juice	14 days	PCR-DGGE	↑ <i>Lactobacillus</i> spp. ↓ <i>Enterobacteria</i>	
Orange juice (Cara Cara)	Human study (healthy individuals)	500 mL of orange juice	7 days	HTS	<u>Family level:</u> ↑ Mogibacteriaceae, Tissierellaceae Porphyromonadaceae, Christensenellaceae, Enterobacteriaceae, Lachnospiraceae, and Ruminococcaceae ↓ Odoribacteraceae <u>Genus level:</u> ↑ <i>Parabacteroides</i> and <i>Butyricimonas</i> ↓ <i>Odoribacter</i>	(Brasili et al., 2019)
Orange juice (Bahia)	Human study (healthy individuals)	500 mL of orange juice	7 days	HTS	<u>Family level:</u> ↑ Mogibacteriaceae, Enterococcaceae, Veillonellaceae, Coriobacteriaceae, Enterococcaceae, Clostridiaceae, and Pasteurellaceae ↓ Ruminococcaceae <u>Genus level:</u> ↑ <i>Adlercreutzia</i> , <i>Enterococcus</i> , <i>Clostridium</i> , and <i>Anaerotruncus</i>	
Orange juice	Human study (healthy individuals)	300 mL of orange juice	8 weeks	PCR-DGGE	↑ <i>Bifidobacterium</i> spp. and <i>Lactobacillus</i> spp.	(Lima et al., 2019)

**Table 1. 8:** Modulation of the gut microbiota by pomegranates.

**HTS:** High throughput sequencing; **qPCR:** Quantitative polymerase chain reaction; **FISH:** Fluorescent in situ hybridization.

Pomegranates	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Pomegranate (POMx)	<i>In vitro</i> (batch-culture fermentation)	1.5 mL of POMx	0, 5, 10, 24 and 48 hours	FISH	↑ <i>Bifidobacterium</i> spp. and <i>Lactobacillus</i> spp.	(Bialowska et al., 2010)
Pomegranate beverage	Animal study (healthy rats)	<i>ad libitum</i>	2 weeks	qPCR	↑ Ruminococcaceae	(Kim et al., 2017)

**Table 1. 8**

Pomegranates	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Pomegranate (POMx)	Human study (healthy individuals)	1000 mg of POMx	4 weeks	qPCR	↑ <i>Akkermansia muciniphila</i>	(Hennig et al., 2017)
Pomegranate (POMx)	Human study (healthy individuals)	1000 mg of POMx	4 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Actinobacteria ↓ Firmicutes <u>Genus level:</u> ↑ <i>Lactobacillus</i> , <i>Akkermansia</i> , <i>Prevotella</i> , <i>Butyrivibrio</i> , <i>Enterobacter</i> , <i>Escherichia</i> , <i>Serratia</i> , and <i>Veillonella</i> ↓ <i>Collinsella</i>	(Z. Li, Hennig, et al., 2015)

**Table 1. 9:** Modulation of the gut microbiota by avocados.**HTS:** High throughput sequencing.

Avocados	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Avocado (Hass Avocado)	Human study (healthy overweight/obese individuals)	1 Hass avocado daily	12 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Bacteroidetes ↓ Firmicutes <u>Family level:</u> ↑ Veillonellaceae, Sutterellaceae, Ruminococcaceae, and Prevotellaceae <u>Genus level:</u> ↑ <i>Ruminococcus</i> , <i>Prevotella</i> , <i>Dialister</i> , <i>Sutterella</i> , <i>Bilophila</i> , <i>Holdemanella</i> , <i>Herbaspirillum</i> , and <i>Acetivibrio</i> ↓ <i>Bacteroides</i> and <i>Methanospaera</i>	(Hennig et al., 2019)
Avocado (Hass Avocado)	Human study (healthy overweight/obese individuals)	1 Hass avocado daily	12 weeks	HTS (Illumina Miseq)	↑ <i>Lachnospira</i> and <i>Faecalibacterium</i>	(Thompson et al., 2019)

**Table 1. 10:** Modulation of the gut microbiota by bitter melons.**HTS:** High throughput sequencing.

Bitter melons	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Bitter melon ( <i>Momordica charantia</i> L.)	Animal study (obese rats)	300 mg of bitter melon powder /kg	8 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Odoribacteraceae ↓ Desulfovibrionaceae and Enterobacteriaceae <u>Genus level:</u> ↑ <i>Faecalibacterium</i> , <i>Allobaculum</i> , <i>Butyricimonas</i> , and <i>Odoribacter</i> ↓ <i>Escherichia</i>	(Bai et al., 2016)

**Table 1. 10**

Bitter melons	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Bitter melon	Animal study (obese rats)	400 mg of bitter melon powder /kg	8 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Verrucomicrobia <u>Genus level:</u> ↑ <i>Blautia</i> , <i>Anaerotruncus</i> , <i>Lactococcus</i> , <i>Allobaculum</i> , and <i>Oceanobacillus</i> ↓ <i>Prevotella</i> , <i>Anaeroplasma</i>	(Bai et al., 2018)
Bitter melon (dried juice)	Animal study (mice fed a high-fat diet)	1.5% of freeze-dried bitter melon juice	16 weeks	HTS	<u>Phylum level:</u> ↑ Bacteroidetes ↓ Firmicutes ↓ Actinobacteria <u>Other levels:</u> ↑ <i>Lactobacillus</i> , Lactobacillales, and Clostridiaceae ↓ <i>Bacteroides</i>	(Nerurkar et al., 2019)

**Table 1. 11:** Modulation of the gut microbiota by vegetables.

**HTS:** High throughput sequencing; **qPCR:** Quantitative polymerase chain reaction.

Vegetables	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Broccoli	Animal study (mice, inflammation bowel disease model)	10% of dried broccoli powder)	21 weeks	qPCR	↔ <i>Bacteroides-Prevotella-Porphyromonas</i> , <i>Bifidobacterium</i> , and <i>Bacteroides vulgatus</i> ↓ <i>Clostridium perfringens</i> , <i>Enterococcus</i> , <i>Lactobacillus</i> , <i>Escherichia coli</i> , and <i>Faecalibacterium prausnitzii</i>	(Paturi et al., 2012)
Broccoli	Animal study (healthy rats)	10% of dried cooked broccoli powder)	4, 7, or 14 days	HTS (Illumina Miseq)	↑ <i>Akkermansia</i> , <i>Oscillospira</i> , and Ruminococcaceae ↓ <i>Blautia</i> , <i>Clostridium</i> , and <i>Dorea</i>	(X. Liu et al., 2017)
Broccoli	Human study (healthy individuals)	200 g of cooked broccoli daily	15 days	qPCR	<u>Phylum level:</u> ↑ Bacteroidetes ↓ Firmicutes <u>Genus level:</u> ↑ <i>Bacteroides</i>	(Kaczmarek et al., 2019)
Mushrooms ( <i>Agaricus bisporus</i> , commonly called white button mushrooms)	Animal study (healthy rats)	1% of dried mushroom	6 weeks	454 Titanium sequencer	<u>Phylum level:</u> ↑ Bacteroidetes, Mollicutes, Proteobacteria sub-classes Delta, Gamma, and Epsilon, and Verrucomicrobia ↓ Firmicutes, Erysipelotrichi, Deferribacteres, and Clostridia	(Varshney et al., 2013)

**Table 1. 11 (Cont.)**

Vegetables	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Mushrooms (white button mushrooms)	Animal study (healthy pigs)	75 g or 150 g of dried mushrooms	6 weeks	Multi-Tag Sequencing	<u>Family level:</u> ↑ Ruminococcaceae, Lachnospiraceae, Veillonellaceae, Succinivibrionaceae, and Clostridiaceae <u>Genus level:</u> ↑ <i>Prevotella</i> , <i>Oscillibacter</i> , <i>Butyrivibrio</i> , <i>Fusicatenibacter</i> , <i>Robinsoniella</i> , <i>Eisenbergiella</i> , <i>Fusicatenibacter</i> , <i>Eisenbergiella</i> , <i>Lachnobacterium</i> , <i>Robinsoniella</i> , <i>Schwartzia</i> , <i>Selenomonas</i> , <i>Anerovibrio</i> , <i>Succinivibrio</i> , <i>Sarcina</i> , and <i>Clostridium sensu stricto</i> ↓ <i>Bifidobacterium</i>	(Solan o-Aguila r et al., 2018)
Mushrooms ( <i>Armillariella tabescens</i> )	Animal study (early-weaned pigs)	0.1, 0.3, or 0.9% of dried <i>Armillariella tabescens</i>	30 days	qPCR	↑ <i>Lactobacillus</i> spp. and <i>Bifidobacterium</i> spp. ↓ <i>E. coli</i>	(Chen et al., 2017)
Mushrooms ( <i>Bunashimeji</i> , <i>Shiitake</i> , <i>Maitake</i> , and <i>Eringi</i> )	Animal study (mice fed a high-fat diet)	0.5% or 3% of dried mushrooms	4 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↑ Firmicutes and Actinobacteria ↓ Bacteroidetes and Proteobacteria <u>Genus level:</u> ↑ <i>Allobaculum</i> , <i>Bifidobacterium</i> , <i>Ruminococcus</i> , <i>Lactobacillus</i> , <i>Lactococcus</i> , <i>Streptococcus</i> , <i>Adlercreutzia</i> , and <i>Sutterella</i> ↓ <i>Bacteroides</i> , <i>Prevotella</i> , <i>Mucispirillum</i> , <i>Dorea</i> , <i>Roseburia</i> , <i>Anaerotruncus</i> , <i>Oscillospira</i> , <i>Escherichia</i> , and <i>Akkermansia</i>	(Shimizu et al., 2018)

**Table 1. 11 (Cont.)**

Vegetables	Study type	Dosage	Treatment duration	Analytical method	Changes in gut microbiota	Ref.
Mushrooms ( <i>Pleurotus eryngii</i> )	Animal study (IBD mice model)	1.5% or 3% of <i>Pleurotus eryngii</i> mushroom powder	6 weeks	HTS (Illumina Miseq)	<u>Phylum level:</u> ↓ Actinobacteria <u>Class level:</u> ↓ Mollicutes <u>Family level:</u> ↑ S24-7 and Odoribacteraceae ↓ Desulfovibrionaceae and Enterococcaceae <u>Genus level:</u> ↑ <i>Adlercreutzia</i> , <i>Akkermanisa</i> , <i>Lactobacillus</i> , <i>Anaerostipes</i> , and <i>Allobaculum</i> ↓ <i>Turicibacter</i> , <i>Dorea</i> , <i>rc4-4</i> , <i>Bacteroides</i> , and <i>Prvotella</i>	(Q. Hu et al., 2019)
Jerusalem artichoke	Animal study (healthy rats)	2, 4, or 6% of Jerusalem artichoke powder	12 weeks	Culture method	↑ <i>Lactobacillus</i> spp. and <i>Bifidobacterium</i> spp. ↓ Coliforms	(Sama l et al., 2017)

## References

- 2015–2020 Dietary Guidelines. (2015). Retrieved from <https://health.gov/our-work/food-nutrition/2015-2020-dietary-guidelines>
- Aduldecha, C., Kaewpradit, W., Vorasoot, N., Puangbut, D., Jogloy, S., & Patanothai, A. (2016). Effects of water regimes on inulin content and inulin yield of Jerusalem artichoke genotypes with different levels of drought tolerance. *Turkish Journal of Agriculture and Forestry*, 40(3), 335-343.
- Aguirre, M., & Venema, K. (2015). Does the gut microbiota contribute to obesity? Going beyond the gut feeling. *Microorganisms*, 3(2), 213-235.
- Ajila, C. M., Naidu, K. A., Bhat, S. G., & Rao, U. J. S. P. (2007). Bioactive compounds and antioxidant potential of mango peel extract. *Food Chemistry*, 105(3), 982-988.
- Al Othaim, A., Marasini, D., & Carbonero, F. (2020). Impact of increasing concentration of tart and sweet cherries juices concentrates on healthy mice gut microbiota. *Food Frontiers*, 1(3), 224-233. doi:10.1002/fft2.46
- Alhinai, E. A., Walton, G. E., & Commane, D. M. (2019). The Role of the Gut Microbiota in Colorectal Cancer Causation. *International Journal of Molecular Sciences*, 20(21), 5295.
- Amirullah, N. A., Abidin, N. Z., & Abdullah, N. (2018). The potential applications of mushrooms against some facets of atherosclerosis: A review. *Food Research International*, 105, 517-536.
- Angelino, D., Dosz, E. B., Sun, J., Hoeflinger, J. L., Van Tassell, M. L., Chen, P., . . . Jeffery, E. H. (2015). Myrosinase-dependent and-independent formation and control of isothiocyanate products of glucosinolate hydrolysis. *Frontiers in plant science*, 6, 831.
- Anonye, B. O. (2017). Commentary: dietary polyphenols promote growth of the gut bacterium *Akkermansia muciniphila* and attenuate high-fat diet-induced metabolic syndrome. *Frontiers in immunology*, 8, 850.
- Aoki, R., Kamikado, K., Suda, W., Takii, H., Mikami, Y., Suganuma, N., . . . Koga, Y. (2017). A proliferative probiotic *Bifidobacterium* strain in the gut ameliorates progression of metabolic disorders via microbiota modulation and acetate elevation. *Scientific reports*, 7, 43522.
- Attri, S., & Goel, G. (2018). Influence of polyphenol rich seabuckthorn berries juice on release of polyphenols and colonic microbiota on exposure to simulated human digestion model. *Food Research International*, 111, 314-323.
- Attri, S., Sharma, K., Raigond, P., & Goel, G. (2018). Colonic fermentation of polyphenolics from Sea buckthorn (*Hippophae rhamnoides*) berries: Assessment of effects on microbial diversity by Principal Component Analysis. *Food Research International*, 105, 324-332.

- Bäckhed, F., Ley, R. E., Sonnenburg, J. L., Peterson, D. A., & Gordon, J. I. (2005). Host-bacterial mutualism in the human intestine. *Science*, 307(5717), 1915-1920.
- Bai, J., Zhu, Y., & Dong, Y. (2016). Response of gut microbiota and inflammatory status to bitter melon (*Momordica charantia* L.) in high fat diet induced obese rats. *Journal of Ethnopharmacology*, 194, 717-726.
- Bai, J., Zhu, Y., & Dong, Y. (2018). Modulation of gut microbiota and gut-generated metabolites by bitter melon results in improvement in the metabolic status in high fat diet-induced obese rats. *Journal of Functional Foods*, 41, 127-134.
- Balamurugan, R., Rajendiran, E., George, S., Samuel, G. V., & Ramakrishna, B. S. (2008). Real-time polymerase chain reaction quantification of specific butyrate-producing bacteria, *Desulfovibrio* and *Enterococcus faecalis* in the feces of patients with colorectal cancer. *Journal of gastroenterology and hepatology*, 23(8pt1), 1298-1303.
- Baldwin, J., Collins, B., Wolf, P. G., Martinez, K., Shen, W., Chuang, C.-C., . . . Chang, E. (2016). Table grape consumption reduces adiposity and markers of hepatic lipogenesis and alters gut microbiota in butter fat-fed mice. *The Journal of nutritional biochemistry*, 27, 123-135.
- Balish, E., & Warner, T. (2002). *Enterococcus faecalis* induces inflammatory bowel disease in interleukin-10 knockout mice. *The American journal of pathology*, 160(6), 2253-2257.
- Bang, S.-J., Kim, G., Lim, M. Y., Song, E.-J., Jung, D.-H., Kum, J.-S., . . . Seo, D.-H. (2018). The influence of in vitro pectin fermentation on the human fecal microbiome. *Amb Express*, 8(1), 1-9.
- Bao, B., Chen, Y.-G., Zhang, L., Xu, Y. L. N., Wang, X., Liu, J., & Qu, W. (2013). *Momordica charantia* (Bitter Melon) reduces obesity-associated macrophage and mast cell infiltration as well as inflammatory cytokine expression in adipose tissues. *PloS one*, 8(12).
- Barlow, G. M., Yu, A., & Mathur, R. (2015). Role of the gut microbiome in obesity and diabetes mellitus. *Nutrition in clinical practice*, 30(6), 787-797.
- Barnes, R. C., Kim, H., Fang, C., Bennett, W., Nemec, M., Sirven, M. A., . . . Mertens-Talcott, S. U. (2019). Body Mass Index as a Determinant of Systemic Exposure to Gallotannin Metabolites during 6-Week Consumption of Mango (*Mangifera indica* L.) and Modulation of Intestinal Microbiota in Lean and Obese Individuals. *Molecular nutrition & food research*, 63(2), 1800512.
- Berkow, S. E., & Barnard, N. (2006). Vegetarian diets and weight status. *Nutrition reviews*, 64(4), 175-188.
- Bhaduri, A., Dubey, A. K., & Basu, A. (2019). Method for increasing abundance of lachnospira. In: Google Patents.



- Bialonska, D., Ramnani, P., Kasimsetty, S. G., Muntha, K. R., Gibson, G. R., & Ferreira, D. (2010). The influence of pomegranate by-product and punicalagins on selected groups of human intestinal microbiota. *International journal of food microbiology*, 140(2-3), 175-182.
- Biragyn, A., & Ferrucci, L. (2018). Gut dysbiosis: a potential link between increased cancer risk in ageing and inflammaging. *The Lancet Oncology*, 19(6), e295-e304.
- Blatchford, P., Stoklosinski, H., Walton, G., Swann, J., Gibson, G., Gearry, R., & Ansell, J. (2015). Kiwifruit fermentation drives positive gut microbial and metabolic changes irrespective of initial microbiota composition. *Bioactive Carbohydrates and Dietary Fibre*, 6(1), 37-45.
- Bradbury, K. E., Crowe, F. L., Appleby, P. N., Schmidt, J. A., Travis, R. C., & Key, T. J. (2014). Serum concentrations of cholesterol, apolipoprotein AI and apolipoprotein B in a total of 1694 meat-eaters, fish-eaters, vegetarians and vegans. *European Journal of Clinical Nutrition*, 68(2), 178-183.
- Brasili, E., Hassimotto, N. M. A., Del Chierico, F., Marini, F., Quagliariello, A., Sciubba, F., . . . Lajolo, F. (2019). Daily consumption of orange juice from *Citrus sinensis* L. Osbeck cv. Cara Cara and cv. Bahia differently affects gut microbiota profiling as unveiled by an integrated meta-omics approach. *Journal of agricultural and food chemistry*, 67(5), 1381-1391.
- Budak, N. H. (2017). Bioactive components of *Prunus avium* L. black gold (red cherry) and *Prunus avium* L. stark gold (white cherry) juices, wines and vinegars. *Journal of food science and technology*, 54(1), 62-70.
- Cai, X., Han, Y., Gu, M., Song, M., Wu, X., Li, Z., . . . Xiao, H. (2019). Dietary cranberry suppressed colonic inflammation and alleviated gut microbiota dysbiosis in dextran sodium sulfate-treated mice. *Food & Function*, 10(10), 6331-6341.
- Cancer. (2018). Retrieved from [https://www.who.int/health-topics/cancer#tab=tab\\_1](https://www.who.int/health-topics/cancer#tab=tab_1)
- Cani, P. D., & de Vos, W. M. (2017). Next-generation beneficial microbes: the case of *Akkermansia muciniphila*. *Frontiers in microbiology*, 8, 1765.
- Cardiovascular diseases (CVDs). (2017). Retrieved from [https://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
- Cardona, F., Andrés-Lacueva, C., Tulipani, S., Tinahones, F. J., & Queipo-Ortuño, M. I. (2013). Benefits of polyphenols on gut microbiota and implications in human health. *The Journal of nutritional biochemistry*, 24(8), 1415-1422.
- Cassidy, A., & Minihaue, A.-M. (2017). The role of metabolism (and the microbiome) in defining the clinical efficacy of dietary flavonoids. *The American journal of clinical nutrition*, 105(1), 10-22.

- Centers for Disease Control and, P. (2020). National diabetes statistics report, 2020. *Atlanta, GA: Centers for Disease Control and Prevention, US Department of Health and Human Services.*
- Chassard, C., Delmas, E., Robert, C., Lawson, P. A., & Bernalier-Donadille, A. (2012). *Ruminococcus champanellensis* sp. nov., a cellulose-degrading bacterium from human gut microbiota. *International Journal of Systematic and Evolutionary Microbiology*, 62(1), 138-143.
- Chen, W. B., Cheng, M. J., Tian, Y. B., Wang, Q. H., Wang, B., Li, M. J., & Fang, R. J. (2017). Effects of *Armillariella tabescens* mycelia on the growth performance and intestinal immune response and microflora of early-weaned pigs. *Animal Science Journal*, 88(9), 1388-1397.
- Chiu, T. H. T., Pan, W.-H., Lin, M.-N., & Lin, C.-L. (2018). Vegetarian diet, change in dietary patterns, and diabetes risk: a prospective study. *Nutrition & diabetes*, 8(1), 1-9.
- Cho, N. H., Shaw, J. E., Karuranga, S., Huang, Y., da Rocha Fernandes, J. D., Ohlrogge, A. W., & Malanda, B. (2018). IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes research and clinical practice*, 138, 271-281.
- Chung, W. S. F., Meijerink, M., Zeuner, B., Holck, J., Louis, P., Meyer, A. S., . . . Duncan, S. H. (2017). Prebiotic potential of pectin and pectic oligosaccharides to promote anti-inflammatory commensal bacteria in the human colon. *FEMS microbiology ecology*, 93(11), fix127.
- Cirstea, M., Radisavljevic, N., & Finlay, B. B. (2018). Good bug, bad bug: breaking through microbial stereotypes. *Cell host & microbe*, 23(1), 10-13.
- Collins, B., Hoffman, J., Martinez, K., Grace, M., Lila, M. A., Cockrell, C., . . . Zhong, W. (2016). A polyphenol-rich fraction obtained from table grapes decreases adiposity, insulin resistance and markers of inflammation and impacts gut microbiota in high-fat-fed mice. *The Journal of nutritional biochemistry*, 31, 150-165.
- Costa, C., Tsatsakis, A., Mamoulakis, C., Teodoro, M., Briguglio, G., Caruso, E., . . . Kouretas, D. (2017). Current evidence on the effect of dietary polyphenols intake on chronic diseases. *Food and Chemical Toxicology*, 110, 286-299.
- Crascì, L., Lauro, M. R., Puglisi, G., & Panico, A. (2018). Natural antioxidant polyphenols on inflammation management: Anti-glycation activity vs metalloproteinases inhibition. *Critical Reviews in Food Science and Nutrition*, 58(6), 893-904.
- Cui, J., Lian, Y., Zhao, C., Du, H., Han, Y., Gao, W., . . . Zheng, J. (2019). Dietary Fibers from Fruits and Vegetables and Their Health Benefits via Modulation of Gut Microbiota. *Comprehensive Reviews in Food Science and Food Safety*, 18(5), 1514-1532.
- Dao, M. C., Everard, A., Aron-Wisnewsky, J., Sokolovska, N., Prifti, E., Verger, E. O., . . . Hoyles, L. (2016). *Akkermansia muciniphila* and improved metabolic health during a

- dietary intervention in obesity: relationship with gut microbiome richness and ecology. *Gut*, 65(3), 426-436.
- De Palma, G., Nadal, I., Medina, M., Donat, E., Ribes-Koninckx, C., Calabuig, M., & Sanz, Y. (2010). Intestinal dysbiosis and reduced immunoglobulin-coated bacteria associated with coeliac disease in children. *BMC microbiology*, 10(1), 1-7.
- Del Chierico, F., Nobili, V., Vernocchi, P., Russo, A., De Stefanis, C., Gnani, D., . . . Capuani, G. (2017). Gut microbiota profiling of pediatric nonalcoholic fatty liver disease and obese patients unveiled by an integrated meta-omics-based approach. *Hepatology*, 65(2), 451-464.
- Denis, M.-C., Roy, D., Yeganeh, P. R., Desjardins, Y., Varin, T., Haddad, N., . . . Furtos, A. (2016). Apple peel polyphenols: a key player in the prevention and treatment of experimental inflammatory bowel disease. *Clinical Science*, 130(23), 2217-2237.
- Diabetes. (2020). Retrieved from <https://www.who.int/news-room/fact-sheets/detail/diabetes>
- Dinh, D. M., Volpe, G. E., Duffalo, C., Bhalchandra, S., Tai, A. K., Kane, A. V., . . . Ward, H. D. (2015). Intestinal microbiota, microbial translocation, and systemic inflammation in chronic HIV infection. *The Journal of Infectious Diseases*, 211(1), 19-27.
- Dinu, M., Abbate, R., Gensini, G. F., Casini, A., & Sofi, F. (2017). Vegetarian, vegan diets and multiple health outcomes: a systematic review with meta-analysis of observational studies. *Critical Reviews in Food Science and Nutrition*, 57(17), 3640-3649.
- Dreher, M. L., & Davenport, A. J. (2013). Hass avocado composition and potential health effects. *Critical Reviews in Food Science and Nutrition*, 53(7), 738-750.
- Dueñas, M., Muñoz-González, I., Cueva, C., Jiménez-Girón, A., Sánchez-Patán, F., Santos-Buelga, C., . . . Bartolomé, B. (2015). A survey of modulation of gut microbiota by dietary polyphenols. *BioMed research international*, 2015.
- Duque, A. L. R. F., Monteiro, M., Adorno, M. A. T., Sakamoto, I. K., & Sivieri, K. (2016). An exploratory study on the influence of orange juice on gut microbiota using a dynamic colonic model. *Food Research International*, 84, 160-169.
- Eckburg, P. B., Bik, E. M., Bernstein, C. N., Purdom, E., Dethlefsen, L., Sargent, M., . . . Relman, D. A. (2005). Diversity of the human intestinal microbial flora. *Science*, 308(5728), 1635-1638.
- Elkahoui, S., Levin, C. E., Bartley, G. E., Yokoyama, W., & Friedman, M. (2019). Levels of Fecal Procyanidins and Changes in Microbiota and Metabolism in Mice Fed a High-Fat Diet Supplemented with Apple Peel. *Journal of agricultural and food chemistry*, 67(37), 10352-10360.

- Evans, S. F., Meister, M., Mahmood, M., Eldoumi, H., Peterson, S., Perkins-Veazie, P., . . . Lucas, E. A. (2014). Mango supplementation improves blood glucose in obese individuals. *Nutrition and metabolic insights*, 7, NMI-S17028.
- Everard, A., Belzer, C., Geurts, L., Ouwerkerk, J. P., Druart, C., Bindels, L. B., . . . Delzenne, N. M. (2013). Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity. *Proceedings of the National Academy of Sciences*, 110(22), 9066-9071.
- Falk, A., Olsson, C., Ahrné, S., Molin, G., Adawi, D., & Jeppsson, B. (2007). Ileal pelvic pouch microbiota from two former ulcerative colitis patients, analysed by DNA-based methods, were unstable over time and showed the presence of *Clostridium perfringens*. *Scandinavian journal of gastroenterology*, 42(8), 973-985.
- Feng, Y., Stams, A. J. M., De Vos, W. M., & Sánchez-Andrea, I. (2017). Enrichment of sulfidogenic bacteria from the human intestinal tract. *FEMS Microbiology Letters*, 364(4), fnx028.
- Feng, Z., Long, W., Hao, B., Ding, D., Ma, X., Zhao, L., & Pang, X. (2017). A human stool-derived *Bilophila wadsworthia* strain caused systemic inflammation in specific-pathogen-free mice. *Gut pathogens*, 9(1), 1-10.
- Fernandes, J., Su, W., Rahat-Rozenbloom, S., Wolever, T. M. S., & Comelli, E. M. (2014). Adiposity, gut microbiota and faecal short chain fatty acids are linked in adult humans. *Nutrition & diabetes*, 4(6), e121-e121.
- Ferrario, C., Taverniti, V., Milani, C., Fiore, W., Laureati, M., De Noni, I., . . . Guglielmetti, S. (2014). Modulation of fecal Clostridiales bacteria and butyrate by probiotic intervention with *Lactobacillus paracasei* DG varies among healthy adults. *The Journal of nutrition*, 144(11), 1787-1796.
- Food demand Survey. (2018). Retrieved from [http://www.agecon.okstate.edu/agecon\\_research.asp](http://www.agecon.okstate.edu/agecon_research.asp)
- Fouhse, J. M., Gänzle, M. G., Beattie, A. D., Vasanthan, T., & Zijlstra, R. T. (2017). Whole-grain starch and fiber composition modifies ileal flow of nutrients and nutrient availability in the hindgut, shifting fecal microbial profiles in pigs. *The Journal of nutrition*, 147(11), 2031-2040.
- Furet, J.-P., Kong, L.-C., Tap, J., Poitou, C., Basdevant, A., Bouilliot, J.-L., . . . Henegar, C. (2010). Differential adaptation of human gut microbiota to bariatric surgery-induced weight loss: links with metabolic and low-grade inflammation markers. *Diabetes*, 59(12), 3049-3057.
- Gagnière, J., Raisch, J., Veziant, J., Barnich, N., Bonnet, R., Buc, E., . . . Bonnet, M. (2016). Gut microbiota imbalance and colorectal cancer. *World Journal of Gastroenterology*, 22(2), 501.

- Garcia-Mazcorro, J. F., Lage, N. N., Mertens-Talcott, S., Talcott, S., Chew, B., Dowd, S. E., . . . Noratto, G. D. (2018). Effect of dark sweet cherry powder consumption on the gut microbiota, short-chain fatty acids, and biomarkers of gut health in obese db/db mice. *PeerJ*, 6, e4195.
- Garcia-Mazcorro, J. F., Pedreschi, R., Chew, B., Dowd, S. E., Kawa, J. R., & Noratto, G. (2018). Dietary supplementation with raspberry extracts modifies the fecal microbiota in obese diabetic db/db mice. *J. Microbiol. Biotechnol*, 28(8), 1247-1259.
- Garcia-Mazcorro, J. F., Pedreschi, R., Yuan, J., Kawa, J. R., Chew, B., Dowd, S. E., & Noratto, G. (2019). Apple consumption is associated with a distinctive microbiota, proteomics and metabolomics profile in the gut of Dawley Sprague rats fed a high-fat diet. *PloS one*, 14(3).
- Gentile, C. L., & Weir, T. L. (2018). The gut microbiota at the intersection of diet and human health. *Science*, 362(6416), 776-780.
- Gomes, A., Oudot, C., Macià, A., Foito, A., Carregosa, D., Stewart, D., . . . Brenner, C. (2019). Berry-enriched diet in salt-sensitive hypertensive rats: metabolic fate of (poly) phenols and the role of gut microbiota. *Nutrients*, 11(11), 2634.
- Gondi, M., Basha, S. A., Bhaskar, J. J., Salimath, P. V., & Prasada Rao, U. J. S. (2015). Anti-diabetic effect of dietary mango (*Mangifera indica* L.) peel in streptozotocin-induced diabetic rats. *Journal of the Science of Food and Agriculture*, 95(5), 991-999.
- Greetham, H. L., Gibson, G. R., Giffard, C., Hippe, H., Merkhoffer, B., Steiner, U., . . . Collins, M. D. (2004). *Allobaculum stercoricanis* gen. nov., sp. nov., isolated from canine feces. *Anaerobe*, 10(5), 301-307.
- Gu, J., Thomas-Ahner, J. M., Riedl, K. M., Bailey, M. T., Vodovotz, Y., Schwartz, S. J., & Clinton, S. K. (2019). Dietary black raspberries impact the colonic microbiome and phytochemical metabolites in mice. *Molecular nutrition & food research*, 63(8), 1800636.
- Guergoletto, K. B., Costabile, A., Flores, G., Garcia, S., & Gibson, G. R. (2016). In vitro fermentation of juçara pulp (*Euterpe edulis*) by human colonic microbiota. *Food Chemistry*, 196, 251-258.
- Guglielmetti, S., Fracassetti, D., Taverniti, V., Del Bo', C., Vendrame, S., Klimis-Zacas, D., . . . Porrini, M. (2013). Differential modulation of human intestinal bifidobacterium populations after consumption of a wild blueberry (*Vaccinium angustifolium*) drink. *Journal of agricultural and food chemistry*, 61(34), 8134-8140.
- Hakansson, A., & Molin, G. (2011). Gut microbiota and inflammation. *Nutrients*, 3(6), 637-682.
- Hales, C. M., Carroll, M. D., Fryar, C. D., & Ogden, C. L. (2020). Prevalence of obesity and severe obesity among adults: United States, 2017–2018.

- Han, K. S., Balan, P., Molist Gasa, F., & Boland, M. (2011). Green kiwifruit modulates the colonic microbiota in growing pigs. *Letters in applied microbiology*, 52(4), 379-385.
- Han, Y., Song, M., Gu, M., Ren, D., Zhu, X., Cao, X., . . . Yuan, B. (2019). Dietary intake of whole strawberry inhibited colonic inflammation in dextran-sulfate-sodium-treated mice via restoring immune homeostasis and alleviating gut microbiota dysbiosis. *Journal of agricultural and food chemistry*, 67(33), 9168-9177.
- Heart Disease and Stroke Statistics-2019 At-a-Glance. (2019). Retrieved from <https://healthmetrics.heart.org/wp-content/uploads/2019/02/At-A-Glance-Heart-Disease-and-Stroke-Statistics-%E2%80%93-2019.pdf>
- Heintz-Buschart, A., Pandey, U., Wicke, T., Sixel-Döring, F., Janzen, A., Sittig-Wiegand, E., . . . Wilmes, P. (2018). The nasal and gut microbiome in Parkinson's disease and idiopathic rapid eye movement sleep behavior disorder. *Movement disorders*, 33(1), 88-98.
- Henning, S. M., Summanen, P. H., Lee, R.-P., Yang, J., Finegold, S. M., Heber, D., & Li, Z. (2017). Pomegranate ellagitannins stimulate the growth of *Akkermansia muciniphila* in vivo. *Anaerobe*, 43, 56-60.
- Henning, S. M., Yang, J., Woo, S. L., Lee, R.-P., Huang, J., Rasmussen, A., . . . Tseng, C.-H. (2019). Hass avocado inclusion in a weight-loss diet supported weight loss and altered gut microbiota: a 12-week randomized, parallel-controlled trial. *Current developments in nutrition*, 3(8), nzz068.
- Hever, J., & Cronise, R. J. (2017). Plant-based nutrition for healthcare professionals: implementing diet as a primary modality in the prevention and treatment of chronic disease. *Journal of geriatric cardiology: JGC*, 14(5), 355.
- Hiippala, K., Jouhten, H., Ronkainen, A., Hartikainen, A., Kainulainen, V., Jalanka, J., & Satokari, R. (2018). The potential of gut commensals in reinforcing intestinal barrier function and alleviating inflammation. *Nutrients*, 10(8), 988.
- Hill-Burns, E. M., Debelius, J. W., Morton, J. T., Wissemann, W. T., Lewis, M. R., Wallen, Z. D., . . . Zabetian, C. P. (2017). Parkinson's disease and Parkinson's disease medications have distinct signatures of the gut microbiome. *Movement disorders*, 32(5), 739-749.
- Hillman, E. T., Lu, H., Yao, T., & Nakatsu, C. H. (2017). Microbial ecology along the gastrointestinal tract. *Microbes and environments*, ME17017.
- Hjorth, M. F., Roager, H. M., Larsen, T. M., Poulsen, S. K., Licht, T. R., Bahl, M. I., . . . Astrup, A. (2018). Pre-treatment microbial Prevotella-to-Bacteroides ratio, determines body fat loss success during a 6-month randomized controlled diet intervention. *International journal of obesity*, 42(3), 580-583.
- How Many Adults in the U.S are Vegetarian and Vegan? (2016). Retrieved from [https://www.vrg.org/nutshell/Polls/2016\\_adults\\_veg.htm](https://www.vrg.org/nutshell/Polls/2016_adults_veg.htm)

- Hu, H.-J., Park, S.-G., Jang, H. B., Choi, M.-G., Park, K.-H., Kang, J. H., . . . Cho, S.-H. (2015). Obesity alters the microbial community profile in Korean adolescents. *PloS one*, 10(7).
- Hu, Q., Yuan, B., Wu, X., Du, H., Gu, M., Han, Y., . . . Xiao, H. (2019). Dietary intake of *Pleurotus eryngii* ameliorated dextran-sodium-sulfate-induced colitis in mice. *Molecular nutrition & food research*, 63(17), 1801265.
- Huang, R.-Y., Huang, C.-C., Hu, F. B., & Chavarro, J. E. (2016). Vegetarian diets and weight reduction: a meta-analysis of randomized controlled trials. *Journal of general internal medicine*, 31(1), 109-116.
- Huttenhower, C., Gevers, D., Knight, R., Abubucker, S., Badger, J. H., Chinwalla, A. T., . . . Fulton, R. S. (2012). Structure, function and diversity of the healthy human microbiome. *Nature*, 486(7402), 207.
- Hwang, E.-S. (2018). Comparison of antioxidant capacity and  $\alpha$ -glucosidase inhibitory activity between bitter melon (*Momordica charanti*) fruit and leaf extract. *Asian Pacific Journal of Tropical Biomedicine*, 8(4), 189.
- Iuchi, T., Hosaka, T., Shiroishi, M., Ono, H., Inukai, K., Sumita, T., . . . Awata, T. (2015). Influence of treatment with extracts of *Hypsizygus marmoreus* mushroom on body composition during obesity development in kk-ay mice. *Journal of nutritional science and vitaminology*, 61(1), 96-100.
- Jakobsdottir, G., Blanco, N., Xu, J., Ahrné, S., Molin, G., Sterner, O., & Nyman, M. (2013). Formation of short-chain fatty acids, excretion of anthocyanins, and microbial diversity in rats fed blackcurrants, blackberries, and raspberries. *Journal of nutrition and metabolism*, 2013.
- Jamar, G., Santamarina, A. B., Mennitti, L. V., de Cássia Cesar, H., Oyama, L. M., de Rosso, V. V., & Pisani, L. P. (2018). Bifidobacterium spp. reshaping in the gut microbiota by low dose of juçara supplementation and hypothalamic insulin resistance in Wistar rats. *Journal of Functional Foods*, 46, 212-219.
- Janda, J. M., & Abbott, S. L. (2007). 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. *Journal of Clinical Microbiology*, 45(9), 2761-2764.
- Jandhyala, S. M., Talukdar, R., Subramanyam, C., Vuyyuru, H., Sasikala, M., & Reddy, D. N. (2015). Role of the normal gut microbiota. *World journal of gastroenterology: WJG*, 21(29), 8787.
- Jayachandran, M., Chung, S. S. M., & Xu, B. (2019). A critical review of the relationship between dietary components, the gut microbe *Akkermansia muciniphila*, and human health. *Critical Reviews in Food Science and Nutrition*, 1-12.

- Juan, B. A. I., Ying, Z. H. U., & Ying, D. (2017). Bitter melon powder protects against obesity-associated fatty liver disease by improving colonic microenvironment in rats with high-fat diet-induced obesity. *Biomedical and environmental sciences*, 30(8), 611-615.
- Jumpertz, R., Le, D. S., Turnbaugh, P. J., Trinidad, C., Bogardus, C., Gordon, J. I., & Krakoff, J. (2011). Energy-balance studies reveal associations between gut microbes, caloric load, and nutrient absorption in humans. *The American journal of clinical nutrition*, 94(1), 58-65.
- Kaczmarek, J. L., Liu, X., Charron, C. S., Novotny, J. A., Jeffery, E. H., Seifried, H. E., . . . Holscher, H. D. (2019). Broccoli consumption affects the human gastrointestinal microbiota. *The Journal of nutritional biochemistry*, 63, 27-34.
- Kang, Y., Yang, G., Zhang, S., Ross, C. F., & Zhu, M. J. (2018). Goji Berry Modulates Gut Microbiota and Alleviates Colitis in IL-10-Deficient Mice. *Molecular nutrition & food research*, 62(22), 1800535.
- Karl, J. P., Hatch, A. M., Arcidiacono, S. M., Pearce, S. C., Pantoja-Feliciano, I. G., Doherty, L. A., & Soares, J. W. (2018). Effects of psychological, environmental and physical stressors on the gut microbiota. *Frontiers in microbiology*, 9, 2013.
- Kasai, C., Sugimoto, K., Moritani, I., Tanaka, J., Oya, Y., Inoue, H., . . . Takei, Y. (2015). Comparison of the gut microbiota composition between obese and non-obese individuals in a Japanese population, as analyzed by terminal restriction fragment length polymorphism and next-generation sequencing. *BMC gastroenterology*, 15(1), 100.
- Kawabata, K., Yoshioka, Y., & Terao, J. (2019). Role of intestinal microbiota in the bioavailability and physiological functions of dietary polyphenols. *Molecules*, 24(2), 370.
- Kim, H., Banerjee, N., Sirven, M. A., Minamoto, Y., Markel, M. E., Suchodolski, J. S., . . . Mertens-Talcott, S. U. (2017). Pomegranate polyphenolics reduce inflammation and ulceration in intestinal colitis—involvement of the miR-145/p70S6K1/HIF1 $\alpha$  axis in vivo and in vitro. *The Journal of nutritional biochemistry*, 43, 107-115.
- Kim, H., Caulfield, L. E., Garcia-Larsen, V., Steffen, L. M., Coresh, J., & Rebholz, C. M. (2019). Plant-Based Diets Are Associated With a Lower Risk of Incident Cardiovascular Disease, Cardiovascular Disease Mortality, and All-Cause Mortality in a General Population of Middle-Aged Adults. *Journal of the American Heart Association*, 8(16), e012865.
- Kim, H. I., Kim, J.-K., Kim, J.-Y., Jang, S.-E., Han, M. J., & Kim, D.-H. (2019). Lactobacillus plantarum LC27 and Bifidobacterium longum LC67 simultaneously alleviate high-fat diet-induced colitis, endotoxemia, liver steatosis, and obesity in mice. *Nutrition Research*, 67, 78-89.
- Kinger, M., Kumar, S., & Kumar, V. (2018). Some important dietary polyphenolic compounds: an anti-inflammatory and immunoregulatory perspective. *Mini reviews in medicinal chemistry*, 18(15), 1270-1282.



- Kobayashi, Y., Sugahara, H., Shimada, K., Mitsuyama, E., Kuhara, T., Yasuoka, A., . . . Xiao, J.-z. (2017). Therapeutic potential of *Bifidobacterium breve* strain A1 for preventing cognitive impairment in Alzheimer's disease. *Scientific reports*, 7(1), 1-10.
- Koh, A., De Vadder, F., Kovatcheva-Datchary, P., & Bäckhed, F. (2016). From dietary fiber to host physiology: short-chain fatty acids as key bacterial metabolites. *Cell*, 165(6), 1332-1345.
- Koliada, A., Syzenko, G., Moseiko, V., Budovska, L., Puchkov, K., Perederiy, V., . . . Tkach, S. (2017). Association between body mass index and Firmicutes/Bacteroidetes ratio in an adult Ukrainian population. *BMC microbiology*, 17(1), 120.
- Koutsos, A., Lima, M., Conterno, L., Gasperotti, M., Bianchi, M., Fava, F., . . . Tuohy, K. M. (2017). Effects of commercial apple varieties on human gut microbiota composition and metabolic output using an in vitro colonic model. *Nutrients*, 9(6), 533.
- Koutsos, A., Tuohy, K. M., & Lovegrove, J. A. (2015). Apples and cardiovascular health—is the gut microbiota a core consideration? *Nutrients*, 7(6), 3959-3998.
- Kovatcheva-Datchary, P., Nilsson, A., Akrami, R., Lee, Y. S., De Vadder, F., Arora, T., . . . Bäckhed, F. (2015). Dietary fiber-induced improvement in glucose metabolism is associated with increased abundance of *Prevotella*. *Cell metabolism*, 22(6), 971-982.
- Kowalska-Duplaga, K., Gosiewski, T., Kapusta, P., Sroka-Oleksiak, A., Wędrychowicz, A., Pieczarkowski, S., . . . Fyderek, K. (2019). Differences in the intestinal microbiome of healthy children and patients with newly diagnosed Crohn's disease. *Scientific reports*, 9(1), 1-11.
- Kumar Singh, A., Cabral, C., Kumar, R., Ganguly, R., Kumar Rana, H., Gupta, A., . . . Pandey, A. K. (2019). Beneficial effects of dietary polyphenols on gut microbiota and strategies to improve delivery efficiency. *Nutrients*, 11(9), 2216.
- Kumari, R., Ahuja, V., & Paul, J. (2013). Fluctuations in butyrate-producing bacteria in ulcerative colitis patients of North India. *World journal of gastroenterology: WJG*, 19(22), 3404.
- Lacombe, A., Li, R. W., Klimis-Zacas, D., Kristo, A. S., Tadepalli, S., Krauss, E., . . . Wu, V. C. H. (2013). Lowbush wild blueberries have the potential to modify gut microbiota and xenobiotic metabolism in the rat colon. *PloS one*, 8(6).
- Laparra, J. M., & Sanz, Y. (2010). Interactions of gut microbiota with functional food components and nutraceuticals. *Pharmacological research*, 61(3), 219-225.
- Lavefve, L., Howard, L. R., & Carbonero, F. (2020). Berry polyphenols metabolism and impact on human gut microbiota and health. *Food & Function*, 11(1), 45-65.

- Lazar, V., Ditu, L.-M., Pircalabioru, G. G., Gheorghe, I., Curutiu, C., Holban, A. M., . . . Chifiriuc, M. C. (2018). Aspects of gut microbiota and immune system interactions in infectious diseases, immunopathology, and cancer. *Frontiers in immunology*, 9, 1830.
- LeBlanc, J. G., Chain, F., Martín, R., Bermúdez-Humarán, L. G., Courau, S., & Langella, P. (2017). Beneficial effects on host energy metabolism of short-chain fatty acids and vitamins produced by commensal and probiotic bacteria. *Microbial cell factories*, 16(1), 79.
- Leontowicz, H., Leontowicz, M., Latocha, P., Jesion, I., Park, Y.-S., Katrich, E., . . . Gorinstein, S. (2016). Bioactivity and nutritional properties of hardy kiwi fruit *Actinidia arguta* in comparison with *Actinidia deliciosa* ‘Hayward’ and *Actinidia eriantha* ‘Bidan’. *Food Chemistry*, 196, 281-291.
- Li, M., Fan, Y., Zhang, X., Hou, W., & Tang, Z. (2014). Fruit and vegetable intake and risk of type 2 diabetes mellitus: meta-analysis of prospective cohort studies. *BMJ open*, 4(11), e005497.
- Li, Z., Henning, S. M., Lee, R.-P., Lu, Q.-Y., Summanen, P. H., Thames, G., . . . Finegold, S. M. (2015). Pomegranate extract induces ellagitannin metabolite formation and changes stool microbiota in healthy volunteers. *Food & Function*, 6(8), 2487-2495.
- Li, Z., Summanen, P. H., Komoriya, T., Henning, S. M., Lee, R.-P., Carlson, E., . . . Finegold, S. M. (2015). Pomegranate ellagitannins stimulate growth of gut bacteria in vitro: Implications for prebiotic and metabolic effects. *Anaerobe*, 34, 164-168.
- Licht, T. R., Hansen, M., Bergström, A., Poulsen, M., Krath, B. N., Markowski, J., . . . Wilcks, A. (2010). Effects of apples and specific apple components on the cecal environment of conventional rats: role of apple pectin. *BMC microbiology*, 10(1), 13.
- Lima, A. C. D., Cecatti, C., Fidélis, M. P., Adorno, M. A. T., Sakamoto, I. K., Cesar, T. B., & Sivieri, K. (2019). Effect of daily consumption of orange juice on the levels of blood glucose, lipids, and gut microbiota metabolites: Controlled clinical trials. *Journal of medicinal food*, 22(2), 202-210.
- Liu, J.-P., Zou, W.-L., Chen, S.-J., Wei, H.-Y., Yin, Y.-N., Zou, Y.-Y., & Lu, F.-G. (2016). Effects of different diets on intestinal microbiota and nonalcoholic fatty liver disease development. *World Journal of Gastroenterology*, 22(32), 7353.
- Liu, W., Zhao, S., Wang, J., Shi, J., Sun, Y., Wang, W., . . . Liu, R. (2017). Grape seed proanthocyanidin extract ameliorates inflammation and adiposity by modulating gut microbiota in high-fat diet mice. *Molecular nutrition & food research*, 61(9), 1601082.
- Liu, X., Wang, Y., Hoeflinger, J. L., Neme, B. P., Jeffery, E. H., & Miller, M. J. (2017). Dietary broccoli alters rat cecal microbiota to improve glucoraphanin hydrolysis to bioactive isothiocyanates. *Nutrients*, 9(3), 262.

- Logan, A. C., Jacka, F. N., & Prescott, S. L. (2016). Immune-microbiota interactions: dysbiosis as a global health issue. *Current allergy and asthma reports*, 16(2), 13.
- Lu, F., Li, Y., Zhou, B., Guo, Q., Chen, F., Hu, X., & Zhang, Y. (2019). Early-Life Supplementation of Grape Pomace Extracts Lastingly Promotes Polyphenol Metabolism and Optimizes Gut Microbiota. *Available at SSRN 3405555*.
- Lu, Q.-Y., Zhang, Y., Wang, Y., Wang, D., Lee, R.-p., Gao, K., . . . Heber, D. (2009). California Hass avocado: profiling of carotenoids, tocopherol, fatty acid, and fat content during maturation and from different growing areas. *Journal of agricultural and food chemistry*, 57(21), 10408-10413.
- Lv, L. X., Fang, D. Q., Shi, D., Chen, D. Y., Yan, R., Zhu, Y. X., . . . Wu, W. R. (2016). Alterations and correlations of the gut microbiome, metabolism and immunity in patients with primary biliary cirrhosis. *Environmental microbiology*, 18(7), 2272-2286.
- Majowicz, S. E., Musto, J., Scallan, E., Angulo, F. J., Kirk, M., O'Brien, S. J., . . . International Collaboration on Enteric Disease “Burden of Illness”, S. (2010). The global burden of nontyphoidal Salmonella gastroenteritis. *Clinical infectious diseases*, 50(6), 882-889.
- Martín, R., Chain, F., Miquel, S., Lu, J., Gratadoux, J.-J., Sokol, H., . . . Langella, P. (2014). The commensal bacterium *Faecalibacterium prausnitzii* is protective in DNBS-induced chronic moderate and severe colitis models. *Inflammatory bowel diseases*, 20(3), 417-430.
- Martín, R., Miquel, S., Benevides, L., Bridonneau, C., Robert, V., Hudault, S., . . . Chatel, J. M. (2017). Functional characterization of novel *Faecalibacterium prausnitzii* strains isolated from healthy volunteers: a step forward in the use of *F. prausnitzii* as a next-generation probiotic. *Frontiers in microbiology*, 8, 1226.
- Martínez, I., Wallace, G., Zhang, C., Legge, R., Benson, A. K., Carr, T. P., . . . Walter, J. (2009). Diet-induced metabolic improvements in a hamster model of hypercholesterolemia are strongly linked to alterations of the gut microbiota. *Applied and environmental microbiology*, 75(12), 4175-4184.
- Maruo, T., Sakamoto, M., Ito, C., Toda, T., & Benno, Y. (2008). *Adlercreutzia equolifaciens* gen. nov., sp. nov., an equol-producing bacterium isolated from human faeces, and emended description of the genus *Eggerthella*. *International Journal of Systematic and Evolutionary Microbiology*, 58(5), 1221-1227.
- Masibo, M., & He, Q. (2009). Mango bioactive compounds and related nutraceutical properties—a review. *Food Reviews International*, 25(4), 346-370.
- Masumoto, S., Terao, A., Yamamoto, Y., Mukai, T., Miura, T., & Shoji, T. (2016). Non-absorbable apple procyanidins prevent obesity associated with gut microbial and metabolomic changes. *Scientific reports*, 6, 31208.

- Mathur, R., & Barlow, G. M. (2015). Obesity and the microbiome. *Expert Review of Gastroenterology & Hepatology*, 9(8), 1087-1099.
- Matsumoto, S., Beeson, W. L., Shavlik, D. J., Siapco, G., Jaceldo-Siegl, K., Fraser, G., & Knutsen, S. F. (2019). Association between vegetarian diets and cardiovascular risk factors in non-Hispanic white participants of the Adventist Health Study-2. *Journal of nutritional science*, 8.
- Matziouridou, C., Marungruang, N., Nguyen, T. D., Nyman, M., & Fåk, F. (2016). Lingonberries reduce atherosclerosis in Apoe<sup>-/-</sup> mice in association with altered gut microbiota composition and improved lipid profile. *Molecular nutrition & food research*, 60(5), 1150-1160.
- Mayta-Apaza, A. C., Pottgen, E., De Bodt, J., Papp, N., Marasini, D., Howard, L., . . . Carbonero, F. (2018). Impact of tart cherries polyphenols on the human gut microbiota and phenolic metabolites in vitro and in vivo. *The Journal of nutritional biochemistry*, 59, 160-172.
- McMacken, M., & Shah, S. (2017). A plant-based diet for the prevention and treatment of type 2 diabetes. *Journal of geriatric cardiology: JGC*, 14(5), 342.
- Meehan, C. J., & Beiko, R. G. (2014). A phylogenomic view of ecological specialization in the Lachnospiraceae, a family of digestive tract-associated bacteria. *Genome biology and evolution*, 6(3), 703-713.
- Melina, V., Craig, W., & Levin, S. (2016). Position of the academy of nutrition and dietetics: vegetarian diets. *Journal of the Academy of Nutrition and Dietetics*, 116(12), 1970-1980.
- Mikulic-Petkovsek, M., Stampar, F., Veberic, R., & Sircelj, H. (2016). Wild Prunus fruit species as a rich source of bioactive compounds. *Journal of Food Science*, 81(8), C1928-C1937.
- Miquel, S., Leclerc, M., Martin, R., Chain, F., Lenoir, M., Raguideau, S., . . . Bowen, B. (2015). Identification of metabolic signatures linked to anti-inflammatory effects of Faecalibacterium prausnitzii. *MBio*, 6(2).
- Mizrahi-Man, O., Davenport, E. R., & Gilad, Y. (2013). Taxonomic classification of bacterial 16S rRNA genes using short sequencing reads: evaluation of effective study designs. *PloS one*, 8(1).
- Moreno-Indias, I., Sánchez-Alcoholado, L., Pérez-Martínez, P., Andrés-Lacueva, C., Cardona, F., Tinahones, F., & Queipo-Ortuño, M. I. (2016). Red wine polyphenols modulate fecal microbiota and reduce markers of the metabolic syndrome in obese patients. *Food & Function*, 7(4), 1775-1787.
- Morgan, X. C., & Huttenhower, C. (2012). Human microbiome analysis. *PLoS Comput Biol*, 8(12), e1002808.

- Morissette, A., Kropp, C., Songpadith, J.-P., Junges Moreira, R., Costa, J., Mariné Casadó, R., . . . Boutekrabt, L. (2020). Blueberry proanthocyanidins and anthocyanins improve metabolic health through a gut microbiota-dependent mechanism in diet-induced obese mice. *American Journal of Physiology-Endocrinology and Metabolism*.
- Nash, V., Ranadheera, C. S., Georgousopoulou, E. N., Mellor, D. D., Panagiotakos, D. B., McKune, A. J., . . . Naumovski, N. (2018). The effects of grape and red wine polyphenols on gut microbiota—A systematic review. *Food research international*, 113, 277-287.
- Neish, A. S. (2009). Microbes in gastrointestinal health and disease. *Gastroenterology*, 136(1), 65-80.
- Nerurkar, P. V., Orias, D., Soares, N., Kumar, M., & Nerurkar, V. R. (2019). Momordica charantia (bitter melon) modulates adipose tissue inflammasome gene expression and adipose-gut inflammatory cross talk in high-fat diet (HFD)-fed mice. *The Journal of nutritional biochemistry*, 68, 16-32.
- Neyrinck, A. M., Possemiers, S., Verstraete, W., De Backer, F., Cani, P. D., & Delzenne, N. M. (2012). Dietary modulation of clostridial cluster XIVa gut bacteria (Roseburia spp.) by chitin–glucan fiber improves host metabolic alterations induced by high-fat diet in mice. *The Journal of nutritional biochemistry*, 23(1), 51-59.
- Nile, S. H., & Park, S. W. (2014). Edible berries: Bioactive components and their effect on human health. *Nutrition*, 30(2), 134-144.
- Noratto, G. D., Bertoldi, M. C., Krennek, K., Talcott, S. T., Stringheta, P. C., & Mertens-Talcott, S. U. (2010). Anticarcinogenic effects of polyphenolics from mango (Mangifera indica) varieties. *Journal of agricultural and food chemistry*, 58(7), 4104-4112.
- O'Connor, K., Morissette, M., Strandwitz, P., Ghiglieri, M., Caboni, M., Liu, H., . . . Lewis, K. (2019). Cranberry extracts promote growth of Bacteroidaceae and decrease abundance of Enterobacteriaceae in a human gut simulator model. *PloS one*, 14(11), e0224836.
- Obesity and overweight. (2020). Retrieved from <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>
- Ojo, B., El-Rassi, G. D., Payton, M. E., Perkins-Veazie, P., Clarke, S., Smith, B. J., & Lucas, E. A. (2016). Mango supplementation modulates gut microbial dysbiosis and short-chain fatty acid production independent of body weight reduction in C57BL/6 mice fed a high-fat diet. *The Journal of nutrition*, 146(8), 1483-1491.
- Okumura, R., & Takeda, K. (2017). Roles of intestinal epithelial cells in the maintenance of gut homeostasis. *Experimental & molecular medicine*, 49(5), e338-e338.
- Okumura, R., & Takeda, K. (2018). Maintenance of intestinal homeostasis by mucosal barriers. *Inflammation and regeneration*, 38(1), 5.

- Orlich, M. J., Singh, P. N., Sabaté, J., Fan, J., Sveen, L., Bennett, H., . . . Butler, T. L. (2015). Vegetarian dietary patterns and the risk of colorectal cancers. *JAMA internal medicine*, 175(5), 767-776.
- Ostfeld, R. J. (2017). Definition of a plant-based diet and overview of this special issue. *Journal of geriatric cardiology: JGC*, 14(5), 315.
- Pan, P., Lam, V., Salzman, N., Huang, Y.-W., Yu, J., Zhang, J., & Wang, L.-S. (2017). Black raspberries and their anthocyanin and fiber fractions alter the composition and diversity of gut microbiota in F-344 rats. *Nutrition and cancer*, 69(6), 943-951.
- Papier, K., Appleby, P. N., Fensom, G. K., Knuppel, A., Perez-Cornago, A., Schmidt, J. A., . . . Key, T. J. (2019). Vegetarian diets and risk of hospitalisation or death with diabetes in British adults: results from the EPIC-Oxford study. *Nutrition & diabetes*, 9(1), 1-8.
- Papuc, C., Goran, G. V., Predescu, C. N., Nicorescu, V., & Stefan, G. (2017). Plant polyphenols as antioxidant and antibacterial agents for shelf-life extension of meat and meat products: classification, structures, sources, and action mechanisms. *Comprehensive Reviews in Food Science and Food Safety*, 16(6), 1243-1268.
- Park, S.-K., Kim, M.-S., Roh, S. W., & Bae, J.-W. (2012). *Blautia stercoris* sp. nov., isolated from human faeces. *International Journal of Systematic and Evolutionary Microbiology*, 62(4), 776-779.
- Paturi, G., Butts, C. A., Bentley-Hewitt, K. L., & Ansell, J. (2014). Influence of green and gold kiwifruit on indices of large bowel function in healthy rats. *Journal of Food Science*, 79(8), H1611-H1620.
- Paturi, G., Mandimika, T., Butts, C. A., Zhu, S., Roy, N. C., McNabb, W. C., & Ansell, J. (2012). Influence of dietary blueberry and broccoli on cecal microbiota activity and colon morphology in *mdr1a*<sup>-/-</sup> mice, a model of inflammatory bowel diseases. *Nutrition*, 28(3), 324-330.
- Pereira-Caro, G., Oliver, C. M., Weerakkody, R., Singh, T., Conlon, M., Borges, G., . . . Crozier, A. (2015). Chronic administration of a microencapsulated probiotic enhances the bioavailability of orange juice flavanones in humans. *Free Radical Biology and Medicine*, 84, 206-214.
- Pereira-Caro, G., Borges, G., Ky, I., Ribas, A., Calani, L., Del Rio, D., . . . Crozier, A. (2015). In vitro colonic catabolism of orange juice (poly) phenols. *Molecular nutrition & food research*, 59(3), 465-475.
- Peron, G., Hidalgo-Liberona, N., González-Domínguez, R., Garcia-Aloy, M., Guglielmetti, S., Bernardi, S., . . . Riso, P. (2019). Exploring the molecular pathways behind the effects of nutrients and dietary polyphenols on gut microbiota and intestinal permeability: a perspective on the potential of metabolomics and future clinical applications. *Journal of agricultural and food chemistry*.

- Petersen, C., Wankhade, U. D., Bharat, D., Wong, K., Mueller, J. E., Chintapalli, S. V., . . . Symons, J. D. (2019). Dietary supplementation with strawberry induces marked changes in the composition and functional potential of the gut microbiome in diabetic mice. *The Journal of nutritional biochemistry*, 66, 63-69.
- Plovier, H., Everard, A., Druart, C., Depommier, C., Van Hul, M., Geurts, L., . . . Lichtenstein, L. (2017). A purified membrane protein from *Akkermansia muciniphila* or the pasteurized bacterium improves metabolism in obese and diabetic mice. *Nature medicine*, 23(1), 107.
- Poretzky, R., Rodriguez-R, L. M., Luo, C., Tsementzi, D., & Konstantinidis, K. T. (2014). Strengths and limitations of 16S rRNA gene amplicon sequencing in revealing temporal microbial community dynamics. *PloS one*, 9(4).
- Prieto, I., Hidalgo, M., Segarra, A. B., Martínez-Rodríguez, A. M., Cobo, A., Ramirez, M., . . . Martinez-Canamero, M. (2018). Influence of a diet enriched with virgin olive oil or butter on mouse gut microbiota and its correlation to physiological and biochemical parameters related to metabolic syndrome. *PloS one*, 13(1).
- Puupponen-Pimiä, R., Nohynek, L., Hartmann-Schmidlin, S., Kähkönen, M., Heinonen, M., Määttä-Riihinen, K., & Oksman-Caldentey, K. M. (2005). Berry phenolics selectively inhibit the growth of intestinal pathogens. *Journal of applied microbiology*, 98(4), 991-1000.
- Puupponen-Pimiä, R., Seppänen-Laakso, T., Kankainen, M., Maukonen, J., Törrönen, R., Kolehmainen, M., . . . Aura, A. M. (2013). Effects of ellagitannin-rich berries on blood lipids, gut microbiota, and urolithin production in human subjects with symptoms of metabolic syndrome. *Molecular nutrition & food research*, 57(12), 2258-2263.
- Queipo-Ortuño, M. I., Boto-Ordóñez, M., Murri, M., Gomez-Zumaquero, J. M., Clemente-Postigo, M., Estruch, R., . . . Tinahones, F. J. (2012). Influence of red wine polyphenols and ethanol on the gut microbiota ecology and biochemical biomarkers. *The American journal of clinical nutrition*, 95(6), 1323-1334.
- Quévrain, E., Maubert, M. A., Michon, C., Chain, F., Marquant, R., Tailhades, J., . . . Pigneur, B. (2016). Identification of an anti-inflammatory protein from *Faecalibacterium prausnitzii*, a commensal bacterium deficient in Crohn's disease. *Gut*, 65(3), 415-425.
- Quigley, E. M. M. (2013). Gut bacteria in health and disease. *Gastroenterology & hepatology*, 9(9), 560.
- Rahman, S. U., Li, Y., Huang, Y., Zhu, L., Feng, S., Wu, J., & Wang, X. (2018). Treatment of inflammatory bowel disease via green tea polyphenols: Possible application and protective approaches. *Inflammopharmacology*, 26(2), 319-330.
- Rajilić-Stojanović, M., & de Vos, W. M. (2014). The first 1000 cultured species of the human gastrointestinal microbiota. *FEMS microbiology reviews*, 38(5), 996-1047.

- Rajilić–Stojanović, M., Biagi, E., Heilig, H. G. H. J., Kajander, K., Kekkonen, R. A., Tims, S., & de Vos, W. M. (2011). Global and deep molecular analysis of microbiota signatures in fecal samples from patients with irritable bowel syndrome. *Gastroenterology*, 141(5), 1792-1801.
- Ravn-Haren, G., Dragsted, L. O., Buch-Andersen, T., Jensen, E. N., Jensen, R. I., Nemeth-Balogh, M., . . . Licht, T. R. (2013). Intake of whole apples or clear apple juice has contrasting effects on plasma lipids in healthy volunteers. *European journal of nutrition*, 52(8), 1875-1889.
- Redondo, D., Arias, E., Oria, R., & Venturini, M. E. (2017). Thinned stone fruits are a source of polyphenols and antioxidant compounds. *Journal of the Science of Food and Agriculture*, 97(3), 902-910.
- Reeves, A. E., Koenigsknecht, M. J., Bergin, I. L., & Young, V. B. (2012). Suppression of *Clostridium difficile* in the gastrointestinal tracts of germfree mice inoculated with a murine isolate from the family Lachnospiraceae. *Infection and immunity*, 80(11), 3786-3794.
- Ri, B., Magbagbeola, O. A., Akinwande, A. I., & Okunowo, O. W. (2010). Nutritional and chemical evaluation of *Momordica charantia*. *Journal of Medicinal Plants Research*, 4(21), 2189-2193.
- Ritchie, L. E., Sturino, J. M., Carroll, R. J., Rooney, L. W., Azcarate-Peril, M. A., & Turner, N. D. (2015). Polyphenol-rich sorghum brans alter colon microbiota and impact species diversity and species richness after multiple bouts of dextran sodium sulfate-induced colitis. *FEMS microbiology ecology*, 91(3).
- Rivera-Chávez, F., Zhang, L. F., Faber, F., Lopez, C. A., Byndloss, M. X., Olsan, E. E., . . . Winter, S. E. (2016). Depletion of butyrate-producing *Clostridia* from the gut microbiota drives an aerobic luminal expansion of *Salmonella*. *Cell host & microbe*, 19(4), 443-454.
- Rizzo, N. S., Sabaté, J., Jaceldo-Siegl, K., & Fraser, G. E. (2011). Vegetarian dietary patterns are associated with a lower risk of metabolic syndrome: the adventist health study 2. *Diabetes care*, 34(5), 1225-1227.
- Rodríguez-Morató, J., Matthan, N. R., Liu, J., de la Torre, R., & Chen, C. Y. O. (2018). Cranberries attenuate animal-based diet-induced changes in microbiota composition and functionality: a randomized crossover controlled feeding trial. *The Journal of nutritional biochemistry*, 62, 76-86.
- Rom, O., Korach-Rechtman, H., Hayek, T., Danin-Poleg, Y., Bar, H., Kashi, Y., & Aviram, M. (2017). Acrolein increases macrophage atherogenicity in association with gut microbiota remodeling in atherosclerotic mice: protective role for the polyphenol-rich pomegranate juice. *Archives of toxicology*, 91(4), 1709-1725.



- Rosell, M., Appleby, P., Spencer, E., & Key, T. (2006). Weight gain over 5 years in 21 966 meat-eating, fish-eating, vegetarian, and vegan men and women in EPIC-Oxford. *International journal of obesity*, 30(9), 1389-1396.
- Rossi, O., Khan, M. T., Schwarzer, M., Hudcovic, T., Srutkova, D., Duncan, S. H., . . . Samsom, J. N. (2015). Faecalibacterium prausnitzii strain HTF-F and its extracellular polymeric matrix attenuate clinical parameters in DSS-induced colitis. *PloS one*, 10(4), e0123013.
- Rossi, O., Van Berkel, L. A., Chain, F., Khan, M. T., Taverne, N., Sokol, H., . . . Langella, P. (2016). Faecalibacterium prausnitzii A2-165 has a high capacity to induce IL-10 in human and murine dendritic cells and modulates T cell responses. *Scientific reports*, 6, 18507.
- Round, J. L., & Mazmanian, S. K. (2009). The gut microbiota shapes intestinal immune responses during health and disease. *Nature reviews immunology*, 9(5), 313-323.
- Round, J. L., & Mazmanian, S. K. (2010). Inducible Foxp3<sup>+</sup> regulatory T-cell development by a commensal bacterium of the intestinal microbiota. *Proceedings of the National Academy of Sciences*, 107(27), 12204-12209.
- Rowland, I., Gibson, G., Heinken, A., Scott, K., Swann, J., Thiele, I., & Tuohy, K. (2018). Gut microbiota functions: metabolism of nutrients and other food components. *European journal of nutrition*, 57(1), 1-24.
- Sakamoto, M., Lan, P. T. N., & Benno, Y. (2007). Barnesiella viscericola gen. nov., sp. nov., a novel member of the family Porphyromonadaceae isolated from chicken caecum. *International Journal of Systematic and Evolutionary Microbiology*, 57(2), 342-346.
- Sakamoto, M., Takagaki, A., Matsumoto, K., Kato, Y., Goto, K., & Benno, Y. (2009). Butyricimonas synergistica gen. nov., sp. nov. and Butyricimonas virosa sp. nov., butyric acid-producing bacteria in the family 'Porphyromonadaceae' isolated from rat faeces. *International Journal of Systematic and Evolutionary Microbiology*, 59(7), 1748-1753.
- Samal, L., Chaturvedi, V. B., & Pattanaik, A. K. (2017). Effects of dietary supplementation with Jerusalem artichoke (Helianthus tuberosus L.) tubers on growth performance, nutrient digestibility, activity and composition of large intestinal microbiota in rats. *J Anim Feed Sci*, 26(1), 50-58.
- Sánchez, C. (2017). Bioactives from mushroom and their application. In *Food bioactives* (pp. 23-57): Springer.
- Santisteban, M. M., Qi, Y., Zubcevic, J., Kim, S., Yang, T., Shenoy, V., . . . Rubiano, A. (2017). Hypertension-linked pathophysiological alterations in the gut. *Circulation research*, 120(2), 312-323.
- Sarrabayrouse, G., Bossard, C., Chauvin, J.-M., Jarry, A., Meurette, G., Quévrain, E., . . . Labarrière, N. (2014). CD4CD8 $\alpha\alpha$  lymphocytes, a novel human regulatory T cell subset

- induced by colonic bacteria and deficient in patients with inflammatory bowel disease. *PLoS Biol*, 12(4), e1001833.
- Satija, A., Bhupathiraju, S. N., Rimm, E. B., Spiegelman, D., Chiuve, S. E., Borgi, L., . . . Hu, F. B. (2016). Plant-based dietary patterns and incidence of type 2 diabetes in US men and women: results from three prospective cohort studies. *PLoS medicine*, 13(6).
- Savage, D. C. (1977). Microbial ecology of the gastrointestinal tract. *Annual review of microbiology*, 31(1), 107-133.
- Schippa, S., & Conte, M. P. (2014). Dysbiotic events in gut microbiota: impact on human health. *Nutrients*, 6(12), 5786-5805.
- Sender, R., Fuchs, S., & Milo, R. (2016). Revised estimates for the number of human and bacteria cells in the body. *PLoS biology*, 14(8), e1002533.
- Sharoba, A. M., Farrag, M. A., & Abd El-Salam, A. M. (2013). Utilization of some fruits and vegetables waste as a source of dietary fiber and its effect on the cake making and its quality attributes. *Journal of Agroalimentary Processes and Technologies*, 19(4), 429-444.
- Shaw, K. A., Bertha, M., Hofmekler, T., Chopra, P., Vatanen, T., Srivatsa, A., . . . Zwick, M. E. (2016). Dysbiosis, inflammation, and response to treatment: a longitudinal study of pediatric subjects with newly diagnosed inflammatory bowel disease. *Genome medicine*, 8(1), 75.
- Shimizu, T., Mori, K., Ouchi, K., Kushida, M., & Tsuduki, T. (2018). Effects of dietary intake of Japanese mushrooms on visceral fat accumulation and gut microbiota in mice. *Nutrients*, 10(5), 610.
- Shinohara, K., Ohashi, Y., Kawasumi, K., Terada, A., & Fujisawa, T. (2010). Effect of apple intake on fecal microbiota and metabolites in humans. *Anaerobe*, 16(5), 510-515.
- Siegel, R. L., Miller, K. D., & Jemal, A. (2020). Cancer statistics, 2020. *CA: A Cancer Journal for Clinicians*, 70(1), 7-30.
- Sokol, H., Pigneur, B., Watterlot, L., Lakhdari, O., Bermúdez-Humarán, L. G., Gratadoux, J.-J., . . . Corthier, G. (2008). Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proceedings of the National Academy of Sciences*, 105(43), 16731-16736.
- Solano-Aguilar, G. I., Jang, S., Lakshman, S., Gupta, R., Beshah, E., Sikaroodi, M., . . . Urban, J. F. (2018). The effect of dietary mushroom Agaricus bisporus on intestinal microbiota composition and host immunological function. *Nutrients*, 10(11), 1721.
- Song, M.-y., Wang, J.-h., Eom, T., & Kim, H. (2015). Schisandra chinensis fruit modulates the gut microbiota composition in association with metabolic markers in obese women: a randomized, double-blind placebo-controlled study. *Nutrition Research*, 35(8), 655-663.

- Sonnenburg, E. D., Smits, S. A., Tikhonov, M., Higginbottom, S. K., Wingreen, N. S., & Sonnenburg, J. L. (2016). Diet-induced extinctions in the gut microbiota compound over generations. *Nature*, 529(7585), 212-215.
- Stella, S. P., Ferrarezi, A. C., dos Santos, K. O., & Monteiro, M. (2011). Antioxidant activity of commercial ready-to-drink orange juice and nectar. *Journal of food science*, 76(3), C392-C397.
- Stinco, C. M., Fernández-Vázquez, R., Escudero-Gilete, M. L., Heredia, F. J., Meléndez-Martínez, A. J., & Vicario, I. M. (2012). Effect of orange juice's processing on the color, particle size, and bioaccessibility of carotenoids. *Journal of agricultural and food chemistry*, 60(6), 1447-1455.
- Taing, M.-W., Pierson, J.-T., Hoang, V. L. T., Shaw, P. N., Dietzgen, R. G., Gidley, M. J., . . . Monteith, G. R. (2012). Mango fruit peel and flesh extracts affect adipogenesis in 3T3-L1 cells. *Food & Function*, 3(8), 828-836.
- Tamanai-Shacoori, Z., Smida, I., Bousarghin, L., Loreal, O., Meuric, V., Fong, S. B., . . . Jolivet-Gougeon, A. (2017). Roseburia spp.: a marker of health? *Future Microbiology*, 12(2), 157-170.
- Tannock, G. W. (2010). The bowel microbiota and inflammatory bowel diseases. *International journal of inflammation*, 2010.
- Tantamango-Bartley, Y., Jaceldo-Siegl, K., Fan, J., & Fraser, G. (2013). Vegetarian diets and the incidence of cancer in a low-risk population. *Cancer Epidemiology and Prevention Biomarkers*, 22(2), 286-294.
- Tantamango-Bartley, Y., Knutsen, S. F., Knutsen, R., Jacobsen, B. K., Fan, J., Beeson, W. L., . . . Penniecook, J. (2016). Are strict vegetarians protected against prostate cancer? *The American journal of clinical nutrition*, 103(1), 153-160.
- Therkelsen, S. P., Hetland, G., Lyberg, T., Lygren, I., & Johnson, E. (2016). Cytokine Levels After Consumption of a Medicinal Agaricus blazei Murill-Based Mushroom Extract, AndoSan™, in Patients with Crohn's Disease and Ulcerative Colitis in a Randomized Single-Blinded Placebo-Controlled Study. *Scandinavian journal of immunology*, 84(6), 323-331.
- Therkelsen, S. P., Hetland, G., Lyberg, T., Lygren, I., & Johnson, E. (2016). Effect of the medicinal Agaricus blazei murill-based mushroom extract, andosantm, on symptoms, fatigue and quality of life in patients with crohn's disease in a randomized single-blinded placebo controlled study. *PloS one*, 11(7).
- Thevaranjan, N., Puchta, A., Schulz, C., Naidoo, A., Szamosi, J. C., Verschoor, C. P., . . . Foley, K. P. (2017). Age-associated microbial dysbiosis promotes intestinal permeability, systemic inflammation, and macrophage dysfunction. *Cell host & microbe*, 21(4), 455-466.

- Thompson, S., Edwards, C., Reeser, G., Khan, N., & Holscher, H. (2019). Effects of avocado consumption on gastrointestinal microbial metabolite concentrations and taxa abundances: a randomized, controlled trial (OR23-07-19). *Current developments in nutrition*, 3(Supplement\_1), nzz040-OR023.
- Thursby, E., & Juge, N. (2017). Introduction to the human gut microbiota. *Biochemical Journal*, 474(11), 1823-1836.
- Tian, S., Liu, X., Lei, P., Zhang, X., & Shan, Y. (2018). Microbiota: a mediator to transform glucosinolate precursors in cruciferous vegetables to the active isothiocyanates. *Journal of the Science of Food and Agriculture*, 98(4), 1255-1260.
- Tims, S., Derom, C., Jonkers, D. M., Vlietinck, R., Saris, W. H., Kleerebezem, M., . . . Zoetendal, E. G. (2013). Microbiota conservation and BMI signatures in adult monozygotic twins. *The ISME journal*, 7(4), 707-717.
- Togo, A. H., Khelaifia, S., Bittar, F., Maraninchi, M., Raoult, D., & Million, M. (2016). 'Eisenbergiella massiliensis', a new species isolated from human stool collected after bariatric surgery.
- Tomova, A., Bukovsky, I., Rembert, E., Yonas, W., Alwarith, J., Barnard, N. D., & Kahleova, H. (2019). The effects of vegetarian and vegan diets on gut microbiota. *Frontiers in nutrition*, 6, 47.
- Tonstad, S., Stewart, K., Oda, K., Batech, M., Herring, R. P., & Fraser, G. E. (2013). Vegetarian diets and incidence of diabetes in the Adventist Health Study-2. *Nutrition, Metabolism and Cardiovascular Diseases*, 23(4), 292-299.
- Tu, P., Bian, X., Chi, L., Gao, B., Ru, H., Knobloch, T. J., . . . Lu, K. (2018). Characterization of the functional changes in mouse gut microbiome associated with increased Akkermansia muciniphila population modulated by dietary black raspberries. *ACS omega*, 3(9), 10927-10937.
- Turnbaugh, P. J., Ley, R. E., Mahowald, M. A., Magrini, V., Mardis, E. R., & Gordon, J. I. (2006). An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature*, 444(7122), 1027.
- Tuso, P. J., Ismail, M. H., Ha, B. P., & Bartolotto, C. (2013). Nutritional update for physicians: plant-based diets. *The Permanente Journal*, 17(2), 61.
- Ubeda, C., Djukovic, A., & Isaac, S. (2017). Roles of the intestinal microbiota in pathogen protection. *Clinical & translational immunology*, 6(2), e128.
- Vacca, M., Celano, G., Calabrese, F. M., Portincasa, P., Gobbetti, M., & De Angelis, M. (2020). The Controversial Role of Human Gut Lachnospiraceae. *Microorganisms*, 8(4), 573.
- Van Hul, M., Geurts, L., Plovier, H., Druart, C., Everard, A., Ståhlman, M., . . . Delzenne, N. M. (2018). Reduced obesity, diabetes, and steatosis upon cinnamon and grape pomace are

- associated with changes in gut microbiota and markers of gut barrier. *American Journal of Physiology-Endocrinology and Metabolism*, 314(4), E334-E352.
- Varshney, J., Ooi, J. H., Jayarao, B. M., Albert, I., Fisher, J., Smith, R. L., . . . Cantorna, M. T. (2013). White button mushrooms increase microbial diversity and accelerate the resolution of *Citrobacter rodentium* infection in mice. *The Journal of nutrition*, 143(4), 526-532.
- Vendrame, S., Guglielmetti, S., Riso, P., Arioli, S., Klimis-Zacas, D., & Porrini, M. (2011). Six-week consumption of a wild blueberry powder drink increases bifidobacteria in the human gut. *Journal of agricultural and food chemistry*, 59(24), 12815-12820.
- Vetvicka, V., & Vetvickova, J. (2014). Immune-enhancing effects of Maitake (*Grifola frondosa*) and Shiitake (*Lentinula edodes*) extracts. *Annals of translational medicine*, 2(2).
- Vital, M., Howe, A. C., & Tiedje, J. M. (2014). Revealing the bacterial butyrate synthesis pathways by analyzing (meta) genomic data. *MBio*, 5(2), e00889-00814.
- Vogt, N. M., Kerby, R. L., Dill-McFarland, K. A., Harding, S. J., Merluzzi, A. P., Johnson, S. C., . . . Blennow, K. (2017). Gut microbiome alterations in Alzheimer's disease. *Scientific reports*, 7(1), 1-11.
- Walters, W. A., Xu, Z., & Knight, R. (2014). Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS letters*, 588(22), 4223-4233.
- Wang, F., Zheng, J., Yang, B., Jiang, J., Fu, Y., & Li, D. (2015). Effects of vegetarian diets on blood lipids: a systematic review and meta-analysis of randomized controlled trials. *Journal of the American Heart Association*, 4(10), e002408.
- Wankhade, U. D., Zhong, Y., Lazarenko, O. P., Chintapalli, S. V., Piccolo, B. D., Chen, J.-R., & Shankar, K. (2019). Sex-specific changes in gut microbiome composition following blueberry consumption in C57BL/6J mice. *Nutrients*, 11(2), 313.
- Williams, B. A., Grant, L. J., Gidley, M. J., & Mikkelsen, D. (2017). Gut fermentation of dietary fibres: physico-chemistry of plant cell walls and implications for health. *International Journal of Molecular Sciences*, 18(10), 2203.
- Wright, N., Wilson, L., Smith, M., Duncan, B., & McHugh, P. (2017). The BROAD study: A randomised controlled trial using a whole food plant-based diet in the community for obesity, ischaemic heart disease or diabetes. *Nutrition & diabetes*, 7(3), e256.
- Wu, H.-J., & Wu, E. (2012). The role of gut microbiota in immune homeostasis and autoimmunity. *Gut microbes*, 3(1), 4-14.
- Wu, S., Hu, R., Nakano, H., Chen, K., Liu, M., He, X., . . . Hou, D.-X. (2018). Modulation of gut microbiota by *Lonicera caerulea* L. Berry polyphenols in a mouse model of fatty liver induced by high fat diet. *Molecules*, 23(12), 3213.

- Wylie, K. M., Truty, R. M., Sharpton, T. J., Mihindukulasuriya, K. A., Zhou, Y., Gao, H., . . . Pollard, K. S. (2012). Novel bacterial taxa in the human microbiome. *PloS one*, 7(6), e35294.
- Xiao, S., Fei, N., Pang, X., Shen, J., Wang, L., Zhang, B., . . . Li, M. (2014). A gut microbiota-targeted dietary intervention for amelioration of chronic inflammation underlying metabolic syndrome. *FEMS microbiology ecology*, 87(2), 357-367.
- Xu, J., Cao, K., Li, Y., Zou, X., Chen, C., Szeto, I. M.-Y., . . . Wang, J. (2014). Bitter gourd inhibits the development of obesity-associated fatty liver in C57BL/6 mice fed a high-fat diet. *The Journal of nutrition*, 144(4), 475-483.
- Yan, Z., Yang, F., Hong, Z., Wang, S., Jinjuan, Z., Han, B., . . . Yang, Q. (2019). Blueberry Attenuates Liver Fibrosis, Protects Intestinal Epithelial Barrier, and Maintains Gut Microbiota Homeostasis. *Canadian Journal of Gastroenterology and Hepatology*, 2019.
- Yang, I., Eibach, D., Kops, F., Brenneke, B., Woltemate, S., Schulze, J., . . . Fox, J. G. (2013). Intestinal microbiota composition of interleukin-10 deficient C57BL/6J mice and susceptibility to *Helicobacter hepaticus*-induced colitis. *PloS one*, 8(8), e70783.
- Yang, J. Y., Lee, Y. S., Kim, Y., Lee, S. H., Ryu, S., Fukuda, S., . . . Kim, M. S. (2017). Gut commensal *Bacteroides acidifaciens* prevents obesity and improves insulin sensitivity in mice. *Mucosal immunology*, 10(1), 104-116.
- Yang, Q., Liang, Q., Balakrishnan, B., Belobrajdic, D. P., Feng, Q.-J., & Zhang, W. (2020). Role of dietary nutrients in the modulation of gut microbiota: A narrative review. *Nutrients*, 12(2), 381.
- Yang, S. J., Choi, J. M., Park, S. E., Rhee, E. J., Lee, W. Y., Oh, K. W., . . . Park, C.-Y. (2015). Preventive effects of bitter melon (*Momordica charantia*) against insulin resistance and diabetes are associated with the inhibition of NF- $\kappa$ B and JNK pathways in high-fat-fed OLETF rats. *The Journal of nutritional biochemistry*, 26(3), 234-240.
- Yokoyama, Y., Barnard, N. D., Levin, S. M., & Watanabe, M. (2014). Vegetarian diets and glycemic control in diabetes: a systematic review and meta-analysis. *Cardiovascular diagnosis and therapy*, 4(5), 373.
- Yokoyama, Y., Nishimura, K., Barnard, N. D., Takegami, M., Watanabe, M., Sekikawa, A., . . . Miyamoto, Y. (2014). Vegetarian diets and blood pressure: a meta-analysis. *JAMA internal medicine*, 174(4), 577-587.
- Zeng, M. Y., Inohara, N., & Nuñez, G. (2017). Mechanisms of inflammation-driven bacterial dysbiosis in the gut. *Mucosal immunology*, 10(1), 18-26.
- Zhang, M., Zhou, Q., Dorfman, R. G., Huang, X., Fan, T., Zhang, H., . . . Yu, C. (2016). Butyrate inhibits interleukin-17 and generates Tregs to ameliorate colorectal colitis in rats. *BMC gastroenterology*, 16(1), 1-9.

- Zhang, X., Zhao, Y., Zhang, M., Pang, X., Xu, J., Kang, C., . . . Zhang, Y. (2012). Structural changes of gut microbiota during berberine-mediated prevention of obesity and insulin resistance in high-fat diet-fed rats. *PloS one*, 7(8), e42529.
- Zhang, Y.-J., Li, S., Gan, R.-Y., Zhou, T., Xu, D.-P., & Li, H.-B. (2015). Impacts of gut bacteria on human health and diseases. *International Journal of Molecular Sciences*, 16(4), 7493-7519.
- Zhao, Y., Nakatsu, C., & Jiang, Q. (2019). A Grape Powder Attenuates Colonic Inflammation, Inhibits Colitis-associated Colorectal Cancer Development and Causes Favorable Changes of Gut Microbiota in Mice (OR04-07-19). *Current developments in nutrition*, 3(Supplement\_1), nzz030-OR004.
- Zhao, Y., Wu, J., Li, J. V., Zhou, N.-Y., Tang, H., & Wang, Y. (2013). Gut microbiota composition modifies fecal metabolic profiles in mice. *Journal of proteome research*, 12(6), 2987-2999.
- Zhong, Y., Nyman, M., & Fåk, F. (2015). Modulation of gut microbiota in rats fed high-fat diets by processing whole-grain barley to barley malt. *Molecular nutrition & food research*, 59(10), 2066-2076.
- Zhu, Y., Bai, J., Zhang, Y., Xiao, X., & Dong, Y. (2016). Effects of bitter melon (*Momordica charantia* L.) on the gut microbiota in high fat diet and low dose streptozocin-induced rats. *International Journal of Food Sciences and Nutrition*, 67(6), 686-695.

## CHAPTER 2:

### Impact of Increasing Concentration of Tart and Sweet Cherries Juices Concentrates on Healthy Mice Gut Microbiota

Ayoub Al Othaim<sup>1,2</sup>, Daya Marasini<sup>3,†</sup>, and Franck Carbonero<sup>1,3,4\*</sup>

<sup>1</sup>Cell and Molecular Biology Program; University of Arkansas, Fayetteville, AR, 72701 USA; franck.carbonero@wsu.edu

<sup>2</sup>Department of Medical Laboratories, College of Applied Medical Sciences, Majmaah University, Al-Majmaah, 11952, Saudi Arabia

<sup>3</sup>Department of Food Science; University of Arkansas, Fayetteville, AR, 72704 USA

<sup>4</sup>Department of Nutrition and Exercise Physiology, Elson Floyd School of Medicine, Washington State University-Spokane, 412 East Spokane Falls Boulevard, Spokane, WA, 99202 USA

\* Correspondence: franck.carbonero@wsu.edu

† Current affiliation: IHRC INC, Atlanta, GA, United States of America.

#### Abstract

Cherries are an excellent source of dietary polyphenols and are marketed as sports and health dietary supplements. However, there is still limited direct evidence for their purported nutritional and health benefits. Gut microbiota modulation ought to be tested since most polyphenols reach the colon where they undergo microbial metabolization to become bioactive and bioavailable metabolites. In the present study, varying dilutions of three concentrate cherry juices, Montmorency tart cherry juice, Balaton tart cherry juice, and sweet cherry juice, were examined to determine their potential effect on the murine gut microbiota composition. Forty-five mice were randomly assigned to three different groups. Each group received an increased concentration of an assigned juice (1/20 v/v, 1/15 v/v, 1/10 v/v, 1/7 v/v, and 1/4 v/v) every five days for 25 days. Fecal samples were collected after each administration for microbiota characterization by high throughput 16S rRNA gene sequencing using Illumina Miseq. The



results revealed consistent gut microbiota modulation based on the concentration regardless of the juice type, but not in a true dose-dependent manner. The two median and the two higher concentrations were significantly different from each other and from the baseline and lowest concentration. Increasing cherry consumption consistently resulted in a significant increase of the relative abundance of *Barnesiella* and *Akkermansia*, whereas *Bacteroides* abundance was negatively correlated with the concentration of the juice. Overall, we demonstrate that cherries induce an overall beneficial modulation of the murine gut microbiota, and that amounts of fruits consumed need to be considered to devise appropriate health and nutrition studies.

**Keywords:** Tart cherries, Sweet cherries, Gut microbiota, Polyphenols, Dietary supplement.

## 2.1 Introduction

Cherry fruits belong to the genus *Prunus* from the family Rosaceae. The most common types of cherries consumed by humans are *Prunus cerasus*, designated as sour or tart cherries, and *Prunus avium*, designated as sweet cherries (Kelley, Adkins, & Laugero, 2018; Mayta-Apaza, Marasini, & Carbonero, 2017). Cherry production has been steadily increasing worldwide in the last few decades (Cásedas, Les, Gómez-Serranillos, Smith, & López, 2016), with the average world cherry production 27% higher from 2010 to 2014 than it was from 1995 to 1999 according to data from Food and Agriculture Organization (FAO). Furthermore, data from The United States Department of Agriculture (USDA) showed that the average cherry production in the United States, the second leading global producers and suppliers of cherries, during 2014 to 2016 was nearly double the average production during the 1990s (Perez, Ferreira, & Minor, 2017), and in 2019, the US total cherries production was more than one billion pounds ("Crop Production," 2019). This dramatic growth of cherries production is probably due to the

increasing awareness among food industries and consumers about the purported health-promoting properties of cherries (Blando & Oomah, 2019; Kelley et al., 2018; Mayta-Apaza et al., 2017).

Cherries are nutrient-dense foods and an excellent source of bioactive components including phytochemicals, mainly polyphenols, with relatively low caloric and sugar content (Kelley et al., 2018; Khoo, Clausen, Pedersen, & Larsen, 2011; McCune, Kubota, Stendell-Hollis, & Thomson, 2010; Wojdyło, Nowicka, Laskowski, & Oszmiański, 2014). These properties give cherries unique organoleptic characteristics and desirable nutritional value leading to consumer attractiveness and cash crop status (Bak et al., 2010; P. G. Bell, McHugh, Stevenson, & Howatson, 2014; Mayta-Apaza et al., 2017). A growing body of research indicates that polyphenols have antioxidant and anti-inflammatory properties and may exhibit a wide range of potential health benefits through the regulation of metabolism, chronic disease, weight, and cell proliferation (Cory, Passarelli, Szeto, Tamez, & Mattei, 2018; Levers et al., 2016; Tomás-Barberán, Selma, & Espín, 2016). Generally, both sweet and tart cherries have higher concentrations of total phenolic compounds (Commisso et al., 2017; Ferretti, Bacchetti, Belleggia, & Neri, 2010; McCune et al., 2010). In fact, the total phenolic content in tart cherries is roughly twice higher than sweet cherry, while sweet cherries contain more anthocyanins (Kim, Heo, Kim, Yang, & Lee, 2005; McCune et al., 2010). Cherries are rich in phenolic compounds, mainly, anthocyanins, flavonoids (anthocyanins, flavan-3-ols and flavonols) which are the major class of polyphenolics in tart cherry, and phenolic acids (hydroxycinnamic acids or Hydroxycinnamates) which are the major class of polyphenolics in sweet cherry (Abrankó & Szilvássy, 2015; Chaovanalikit & Wrolstad, 2004; Damar & Ekşi, 2012; Gonçalves, Bento, Jesus, Alves, & Silva, 2018; Mayta-Apaza et al., 2017; Wojdyło et al., 2014). Several studies

have reported potential health benefits of cherry consumption which includes promoting cardiovascular health and reducing the risk of diabetes, obesity, and metabolic syndrome (Blando & Oomah, 2019; Chai, Davis, Wright, Kuczmarski, & Zhang, 2018; Csiki et al., 2015; Juhasz et al., 2013; Lachin, 2014; Nemes et al., 2019). Furthermore, the use of tart cherry juice has been rapidly increasing in sports medicine due to its beneficial impact on improving muscle function and reducing muscle soreness, muscle damage, oxidative stress, and inflammation in athletes (Gao & Chilibeck, 2020; Vitale, Hueglin, & Broad, 2017)

Polyphenols, in general, are relatively large molecules with high molecular weights (Costa et al., 2017; Peron et al., 2019; Santhakumar, Battino, & Alvarez-Suarez, 2018). As a consequence, the absorption of these compounds is relatively low in the upper intestinal tract (Espín, González-Sarriás, & Tomás-Barberán, 2017). The vast majority of polyphenols persist into the large intestine where they undergo biotransformation by bacterial members of gut microbiota, mainly in the colon (Edwards et al., 2017; Ozdal et al., 2016; Peron et al., 2019). The complex interplay between gut microbiota and food indigestible nutrients and phytochemicals, particularly polyphenols, largely contributes to the host's overall health (Kalt et al., 2020). There is a reciprocal relationship between gut microbiota and polyphenols (Gowd, Karim, Shishir, Xie, & Chen, 2019). The gut microbial residence actively drives the metabolism of polyphenols and modulates the biological activity of polyphenols by converting them into phenolic metabolites (Cardona, Andrés-Lacueva, Tulipani, Tinahones, & Queipo-Ortuño, 2013; Kumar Singh et al., 2019). On the other hand, polyphenols and their metabolites can impact the intestinal ecosystem and modulate gut microbiota composition (Braune, Engst, & Blaut, 2016; Gowd et al., 2019; Marín, Miguélez, Villar, & Lombó, 2015; Mayta-Apaza et al., 2018; Nash et al., 2018). However, the knowledge on how dietary polyphenols influence the gut microbiota and its

metabolic potential is still scarce with an emphasis on specific berries and grapes (Espín et al., 2017; Lavefve, Howard, & Carbonero, 2020). In addition to the prevailing focus on specific food products or extracts, there is still limited knowledge about the impact of the concentration or amounts of polyphenols on the gut microbiota. Often, amounts of food used are selected arbitrarily and would be unrealistically high or low concerning a typical human dietary intake. Tart cherry juice, for instance, has been used in most research studies, mainly in sports medicine, in a wide range of dosage and concentrations, mostly from 1oz of concentrated juice to 12oz of non-concentrated juice, with dilution ratio ranges from 1/3 v/v to 1/8 v/v of concentrated juice in water, twice a day. (Phillip G. Bell, Stevenson, Davison, & Howatson, 2016; Phillip G. Bell, Walshe, Davison, Stevenson, & Howatson, 2014; Phillip G. Bell, Walshe, Davison, Stevenson, & Howatson, 2015; Bowtell, Sumners, Dyer, Fox, & Mileva, 2011; Howatson et al., 2012; Kent et al., 2017; Lynn et al., 2014; Mayta-Apaza et al., 2018). That range of tart cherry juice dose seems efficient in accelerating strength recovery after exercise and decreasing blood markers of inflammation and oxidative stress in athletes, but some concentrations may only be realistic for consumption as a dietary supplement. In particular, the optimal bioactive dose to modulate the gut microbiome with a desired response is most often unclear because of the lack of studies including a dose-dependent experimental design. To the best of our knowledge, only two human studies have investigated the potential impact of tart cherry juice on modifying the composition of gut microbiota (Lear et al., 2019; Mayta-Apaza et al., 2018). Choosing the ideal dosage is quite challenging as the phenolic content depends on many factors including product preparation, processing, and shelf-life (Vitale et al., 2017). Therefore, the goal of this study was to investigate the impact of varying amounts of concentrated tart and sweet cherries juices on the murine gut microbiota composition.

## **2.2 Materials and Methods**

### **2.2.1 Animals**

The animal study was conducted at the Animal House Facility of the University of Arkansas after receiving approval from the Institutional Animal Care and Use Committee (IACUC). Forty-five (45) male mice (*Mus musculus* strain C57BL/6J) aged ten weeks were purchased from The Jackson Laboratory (Bar Harbor, ME, USA). Mice were housed in stainless steel cages under standard pathogen-free conditions at a temperature of 22 °C with 50% humidity and exposed to a 12 h light/12 h dark cycle. Before the dietary intervention, mice were provided with standard Teklad chow pellets (Envigo, Madison, WI) and water *ad libitum* for one week.

### **2.2.2 Treatments**

Commercially available pure cherry juices concentrates were used in this study; (i) Montmorency tart cherry juice (MTC) (King Orchards, Central Lake, MI), (ii) Balaton tart cherry juice (BTC) (Royal Farms, Inc., Ellsworth, MI), and (iii) sweet cherry juice (SC) (Plentiful Planet Brands, Boulder, CO).

### **2.2.3 Experiment intervention**

Mice were randomly assigned into three different groups ( $n = 15$  per treatment group), and each major group was split into five sub-groups of three. The groups had been assigned based on the type of juice they received. In this study, group (1) received (MTC), group (2) received (BTC), and group (3) received (SC). The experiment was performed for a duration of 23 days. Animals had *ad libitum* access to chow pellets throughout the experiment. Furthermore, groups received an increased concentration of their assigned juices added to their drinking water

(1/20 v/v, 1/15 v/v, 1/10 v/v, and 1/7 v/v) every five days; except the final intervention where the animals received the highest juice concentration (1/4 v/v) for only three days (**Figure 2. 1**). At the end of each period, mice were transferred from stainless steel cages into metabolic cages (Tecniplast Cdd, 170013) for 6 hours in order to collect fecal samples at day 0, 5, 10, 15, 20, and 23, where day 0 samples represented the baseline. Samples were stored at  $-20^{\circ}\text{C}$  until used for microbiome analysis.

#### **2.2.4 DNA extraction**

A cell lysis step was performed initially to ensure full disruption of microbial cells. Briefly, 220 mg of fecal content were weighed and transformed to 2 ml tube containing 0.1 g of 0.1 mm diameter and 0.1 g of 0.5 mm diameter zirconia-silica beads (BioSpec Products, Bartlesville, OK, United States), and 1 mL of InhibitEX Buffer (from the QIAamp® Fast DNA Stool Mini Kit) was added to the tubes. The tubes were then subjected to a bead-beater for 60 seconds at maximum speed using a FastPrep®-24 bead-beater (MP Biomedicals, Santa Ana, CA, United States) and then heated to  $70^{\circ}\text{C}$  for 5 min in a Thermomixer (Eppendorf, Montesson, France). Microbial DNA was then extracted using a commercial QIAamp DNA stool mini kit (QIAGEN, Valencia, CA) following the manufacturer's protocol.

The NanoDrop™ 1000 Spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA, United States) was used to determine the DNA concentration. Extracted DNA was then visualized following electrophoresis in a 2% agarose gel and 1X TAE (Tris-acetate ethylenediaminetetraacetic acid) buffer (AMRESCO®, Cleveland, OH, United States).

### **2.2.5 Universal polymerase chain reaction**

16S ribosomal RNA gene was amplified by PCR (95 °C for 3 min, followed by 35 cycles at 98 °C for 30 s, 55 °C for 30 s, and 72 °C for 1 min and a final extension at 72 °C for 5 min) using primers (8F 5' - AGAGTTTGATCCTGGCTCAG - 3' and 1541R 5' - AAGGAGGTGATCCAGCCGCA - 3'). PCR reactions were performed in 25 µL mixture containing 3 µL of DNA template; 1 µL of each universal primer (8F and 1541R); 12.5 µL of GoTaq® Green Master Mix (Promega™ Corporation, Wisconsin, USA); and 7.5 µL of Nuclease-Free Water. Agarose gel electrophoresis was then performed to check the success of the amplification by using 1 µL of SYBR Safe DNA Gel Stain (Thermo Fisher Scientific, USA) fluorescent dye.

### **2.2.6 Libraries Preparation and Sequencing**

Index PCR was performed by targeting the V4 region of the bacterial 16S rRNA gene using the dual-index strategy for primer design as previously described (Kozich, Westcott, Baxter, Highlander, & Schloss, 2013). Briefly, dual-index PCR was performed in 25 µL reaction components with 3 µL of DNA template, 2.5 µL of AccuPrime™ PCR buffer II, 0.5 µL of each index primers, 0.1 µL of AccuPrime™ Taq DNA Polymerase (Invitrogen, USA), and 18.4 µL of Nuclease-Free Water following the manufacture's protocol. The PCR consisted of initial denaturation at 95 °C for 3 min, followed by 30 cycles of denaturation at 95 °C for 30 sec, primer annealing at 55 °C for 30 sec, and an extension at 72 °C for 1 min, with a final extension step at 72 °C for 5 min. After successful amplification, the correct size amplicons were subjected to purification and normalization by using SequelPrep kits (Invitrogen, USA) following the

manufacturer's protocol. Amplicon concentrations were measured by Agilent bioanalyzer after pooling the library.

For a very accurate concentration, the amplicons library was quantified by qPCR using the same primers and PerfeCta NGS library quantification kits (Quanta Biosciences, USA) following manufacturer instructions. The library was then denatured and diluted to appropriate concentrations and sequenced on an Illumina MiSeq.

The sequencing reads were downloaded from the Illumina Basespace server in Fastq files format. The sequences were demultiplexed in read 1 and read 2 with approximately 250bp in length. Operational taxonomic units (OTUs) with 97% of identity were determined and classified against the SILVA database as a reference. The Mothur v.1.41.1 pipeline was used for further analysis (Schloss et al., 2009). Non-metric multidimensional scaling (NMDS) plots based on the Bray-Curtis index were obtained in PAST 3.18 software (Hammer, Harper, & Ryan, 2001).

### **2.2.7 Statistical analysis**

Statistical analysis including *t*-test, ANOVA followed by Tukey post hoc, Mann-Whitney U test, and Kruskal-Wallis were performed to detect significant differences between groups (by convention, differences were considered significant when  $P < 0.05$ ). Non-Metric Multidimensional scaling (NMDS) based on count distance metrics (Bray-Curtis similarity index; ANOSIM:  $P < 0.05$ ) were considered to be significant similarities between groups using Past4 software.



## 2.3 Results and Discussion

### 2.3.1 Nutritional analyses

It was observed that the addition of juice to drinking water resulted in increased consumption, but mainly for the median concentration (1/15 and 1/10 v/v; **Figure 2. 2 A**). This would be ascribed to the sweetness of the juices making the water more palatable, as well as a source of calories. Notably, at higher concentration, palatability decreased markedly; and it was noticed that mice started spitting out the water. Such concentrations are tolerable by human consumers when drinking cherry juice as a supplement. We elected to provide juices in drinking water to limit the stress on the animals, but these observations showed that there are limits of such an approach depending on the food tested, and tart cherries, in particular, being astringent with a strong flavor (Clausen, Pedersen, Bertram, & Kidmose, 2011).

The type of cherry juice consumed significantly affected mice weight trajectories, regardless of the concentration in drinking water (**Figure 2. 2 B**). Specifically, the Balaton and sweet cherries juices resulted in significant weight gain, whereas the Montmorency juice resulted in stable weight. Montmorency cherries are known to have lower sugar content, which would directly explain this observation, also supporting their potential as a low-calorie dietary supplement (Blando & Oomah, 2019; Wang et al., 1997).

### 2.3.2 Gut microbiota modulation

Without considering the concentration, Bray-Curtis dissimilarity-based non-metric multidimensional scaling (NMDS) revealed no significant clustering of the gut microbiota profile based on juice type (**Figure 2. 3 A**). This is somewhat unexpected given the difference in sugars, carbohydrates, and possibly phytochemicals in the different juices. However, it is clear

from our observation that the gut microbiota modulation was almost entirely the same with each cherry type. While caution should be used with animal data, this observation suggests that other cherries than the Montmorency variety may provide similar benefits, at least from the gut microbiota perspective.

Regardless of juice type, there was significant clustering based on concentration. Specifically, the two median concentrations (1/15 and 1/10 v/v) and the two higher (1/7 and 1/4 v/v) were significantly different from each other and from the baseline and lowest concentration (**Figure 2. 3 B, C, D**). It is interesting to note that this response does not fulfill the definition of a true dose-dependent response which had been suggested by previous reports (Li et al., 2015) based on comparing low and high doses only. It can thus be postulated that dietary modulation of the gut microbiota probably follows a multimodal response to any nutrient/food dose, and that the amounts used for nutrition and microbiome research need to be tested and validated.

Diversity indices were influenced differently by the concentration of the different juices. MTC induced a consistent decline in diversity regardless of concentration (**Figure 2. 4 A and B**). BTC resulted in fluctuating diversity, but a significant decline for the two highest concentrations (**Figure 2. 4 C and D**). In stark contrast, SC brought diversity down at the median concentrations, especially the 1/10 v/v (**Figure 2. 4 E and F**). Interestingly, these results were inconsistent with another cherry-based study result which showed that obese diabetic mice that were fed the dark sweet cherry supplementation had the highest diversity compared to obese and lean controls (Garcia-Mazcorro et al., 2018).

In this experiment, the mice gut microbiota were initially dominated by relatively similar relative abundance of Bacteroidetes and Firmicutes, followed by Verrucomicrobia,

Proteobacteria, Actinobacteria, and Tenericutes, in line with commonly described murine gut microbiota profiles (Hugenholtz & de Vos, 2018; Nagpal et al., 2018; Nguyen, Vieira-Silva, Liston, & Raes, 2015). While the different juices exerted various dynamics of the phyla, Firmicutes relative abundances were consistently and significantly decreased and replaced by Verrucomicrobia with increasing concentration of the juices, while Bacteroidetes stayed relatively stable (**Figure 2. 5 A and B**). The negative correlation between the abundance of Firmicutes and Verrucomicrobia was also reported in healthy mice supplemented black raspberry as well as in high-fat fed mice consuming *Lonicera caerulea* L. berry for seven weeks and 45 days, respectively (Tu et al., 2018; Wu et al., 2018). Blueberry consumption for four weeks also significantly decreased Firmicutes population in mice (Wankhade et al., 2019).

While Bacteroidetes were stable, there were still significant changes among the genera, with a decrease in *Bacteroides* and an increase in *Barnesiella* (**Figure 2. 6 A and B**). The decrease in *Bacteroides* is consistent with what we observed in human subjects who had initially high *Bacteroides* levels after tart cherry consumption (Mayta-Apaza et al., 2018). The increase in *Barnesiella* was particularly marked at the highest concentrations (from non-detectable to 5-10%). This increase by cherry consumption may be one explanation for tart cherry benefits for exercise recovery, as *Barnesiella* has been reported to increase with exercise and being possibly beneficial (Fielding et al., 2019; Liu et al., 2017).

*Lactobacillus* has been reported, with wide variation, to respond significantly to berries and other polyphenol-rich fruit consumption (Han & Xiao, 2020; Lavefve et al., 2020). In this study, the increase of *Lactobacillus* was observed only for specific and different concentrations for each different juice (**Figure 2. 6 C**). While these erratic dynamics are difficult to interpret,

they may also be a good illustration of the reason for the large variation in *Lactobacillus* bloom in response to polyphenol-rich fruits.

The clear and consistent impact of the cherry consumption was a steep increase of *Akkermansia* (**Figure 2. 6 D**). The increase was specifically marked at higher concentrations, with the BTC even inducing the changes at the median concentrations. In agreement with our findings, the *Akkermansia* spp. relative abundance was increased in obese diabetic mice fed dark sweet cherry powder compared to obese and lean control groups (Garcia-Mazcorro et al., 2018). *Akkermansia muciniphila* is atype species of the genus typically detected in intestinal samples (Derrien, Belzer, & de Vos, 2017; Everard et al., 2013; van Passel et al., 2011). This genus/species has been increasingly reported as highly responsive to polyphenol-rich food consumption in murine models (Anhê et al., 2015; Tu et al., 2018; Wu et al., 2018; Zhang et al., 2017). However, such a response has not been commonly described, at least not as remarkable, in human subjects, which needs to be accounted when discussing such a bloom. *Akkermansia* has also recently been associated with exercise and exercise recovery (Munukka et al., 2018), and this effect of cherry juices is again noteworthy in that light.


## 2.4 Conclusion

We report that cherry juices concentrations in normal drinking water impacted weight gain and palatability in mice, which is important to consider for future animal studies. The impact of cherry juices on gut microbiota was notably influenced by the variation in the concentrations of the juices, but not significantly according to the type of cherry. Increasing concentration of juices was positively correlated with some bacterial genera, including *Akkermansia* and *Barnesiella*, whereas the opposite response was observed for *Bacteroides*



population. These consistent trends can be considered positive, and specifically *Akkermansia* and *Barnesiella* have recently been suggested to play roles in improved exercise recovery, which would make sense knowing that cherries as dietary supplements have been shown to provide such benefits.

## **Acknowledgments**


This research project was funded by a research grant from the Cherry Marketing Institute. Ayoub Al Othaim received a full graduate student stipend from Majmaah University.



**Figure 2. 1:** Study design and intervention.



**Figure 2. 2:** (A) Palatability of cherries juices concentrates as represented by daily water consumption. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 15$ ). Different letters indicate significant difference ( $P < 0.05$ ) between concentrations for (MTC) group. (B) The impact of cherries juices consumption on weight trajectories for 23 days. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 15$ ). Different letters indicate significant difference ( $P < 0.05$ ) while (N.S.) indicates no significant difference.



**Figure 2. 3:** Cherries type (A) have less impact on gut microbiota than concentration and amounts consumed (B, C, and D). Gut microbiota data are shown on non-metric multidirectional scaling (NMDS) (Bray-Curtis similarity index; ANOSIM: ( $P < 0.05$ )).









Figure 2. 3 (Cont.)



**Figure 2. 4:** The impact of concentration of cherries juices on gut microbiota diversity. **(A)** represents the impact of varying concentration of (MTC) on the number of taxa. **(B)** represents the impact of varying concentration of (MTC) on Shannon\_H and Evenness\_e<sup>H/S</sup> indexes. **(C)** represents the impact of varying concentration of (BTC) on the number of taxa. **(D)** represents the impact of varying concentration of (BTC) on Shannon\_H and Evenness\_e<sup>H/S</sup> indexes. **(E)** represents the impact of varying concentrations of (SC) on the number of taxa. **(F)** represents the impact of varying concentration of (SC) on Shannon\_H and Evenness\_e<sup>H/S</sup> indexes. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 15$ ). Different letters indicate significant difference ( $P < 0.05$ ) while (N.S.) indicates no significant difference.



**Figure 2. 5:** Impact of varying concentrations of cherries juices on the phylum level. Increasing concentration of cherries juices result in a decrease of Firmicutes (**A**) and an increase of Verrucomicrobia (**B**), with a slight variation by cherries type. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 15$ ). Different letters indicate significant difference ( $P < 0.05$ ).



**Figure 2. 6:** Impact of varying concentrations of cherries juices on the genus level. Cherries consumption consistently resulted in decreased (A) *Bacteroides* and increased (B) *Barnesiella* and (D) *Akkermansia* in healthy mice. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 15$ ). Different letters indicate significant difference ( $P < 0.05$ ) while (N.S.) indicates no significant difference.

## References

- Abrankó, L., & Szilvássy, B. (2015). Mass spectrometric profiling of flavonoid glycoconjugates possessing isomeric aglycones. *Journal of Mass Spectrometry*, 50(1), 71-80. Doi:10.1002/jms.3474
- Anhê, F. F., Roy, D., Pilon, G., Dudonné, S., Matamoros, S., Varin, T. V., . . . Levy, E. (2015). A polyphenol-rich cranberry extract protects from diet-induced obesity, insulin resistance and intestinal inflammation in association with increased *Akkermansia* spp. Population in the gut microbiota of mice. *Gut*, 64(6), 872-883.
- Bak, I., Lekli, I., Juhasz, B., Varga, E., Varga, B., Gesztelyi, R., . . . Tosaki, A. (2010). Isolation and analysis of bioactive constituents of sour cherry (*Prunus cerasus*) seed kernel: an emerging functional food. *Journal of medicinal food*, 13(4), 905-910.
- Bell, P. G., McHugh, M. P., Stevenson, E., & Howatson, G. (2014). The role of cherries in exercise and health. *Scandinavian Journal of Medicine & Science in Sports*, 24(3), 477-490.
- Bell, P. G., Stevenson, E., Davison, G. W., & Howatson, G. (2016). The effects of montmorency tart cherry concentrate supplementation on recovery following prolonged, intermittent exercise. *Nutrients*, 8(7), 441.
- Bell, P. G., Walshe, I. H., Davison, G. W., Stevenson, E., & Howatson, G. (2014). Montmorency cherries reduce the oxidative stress and inflammatory responses to repeated days high-intensity stochastic cycling. *Nutrients*, 6(2), 829-843.
- Bell, P. G., Walshe, I. H., Davison, G. W., Stevenson, E. J., & Howatson, G. (2015). Recovery facilitation with Montmorency cherries following high-intensity, metabolically challenging exercise. *Applied Physiology, Nutrition, and Metabolism*, 40(4), 414-423.
- Blando, F., & Oomah, B. D. (2019). Sweet and sour cherries: Origin, distribution, nutritional composition and health benefits. *Trends in food science & technology*.
- Bowtell, J. L., Sumners, D. P., Dyer, A., Fox, P., & Mileva, K. N. (2011). Montmorency cherry juice reduces muscle damage caused by intensive strength exercise. *Medicine & Science in Sports & Exercise*, 43(8), 1544-1551.
- Braune, A., Engst, W., & Blaut, M. (2016). Identification and functional expression of genes encoding flavonoid O- and C-glycosidases in intestinal bacteria. *Environmental microbiology*, 18(7), 2117-2129. Doi:10.1111/1462-2920.12864
- Cardona, F., Andrés-Lacueva, C., Tulipani, S., Tinahones, F. J., & Queipo-Ortuño, M. I. (2013). Benefits of polyphenols on gut microbiota and implications in human health. *The Journal of nutritional biochemistry*, 24(8), 1415-1422. Doi:10.1016/j.jnutbio.2013.05.001

- Cásedas, G., Les, F., Gómez-Serranillos, M. P., Smith, C., & López, V. (2016). Bioactive and functional properties of sour cherry juice (*Prunus cerasus*). *Food & Function*, 7(11), 4675-4682.
- Chai, S. C., Davis, K., Wright, R. S., Kuczmarski, M. F., & Zhang, Z. (2018). Impact of tart cherry juice on systolic blood pressure and low-density lipoprotein cholesterol in older adults: a randomized controlled trial. *Food & Function*, 9(6), 3185-3194.
- Chaovanalikit, A., & Wrolstad, R. E. (2004). Anthocyanin and Polyphenolic Composition of Fresh and Processed Cherries. *Journal of Food Science*, 69(1), FCT73-FCT83. Doi:10.1111/j.1365-2621.2004.tb17859.x
- Clausen, M. R., Pedersen, B. H., Bertram, H. C., & Kidmose, U. (2011). Quality of sour cherry juice of different clones and cultivars (*Prunus cerasus* L.) determined by a combined sensory and NMR spectroscopic approach. *Journal of agricultural and food chemistry*, 59(22), 12124-12130.
- Commisso, M., Bianconi, M., Di Carlo, F., Poletti, S., Bulgarini, A., Munari, F., . . . Avesani, L. (2017). Multi-approach metabolomics analysis and artificial simplified phytocomplexes reveal cultivar-dependent synergy between polyphenols and ascorbic acid in fruits of the sweet cherry (*Prunus avium* L.). *PloS one*, 12(7).
- Cory, H., Passarelli, S., Szeto, J., Tamez, M., & Mattei, J. (2018). The Role of Polyphenols in Human Health and Food Systems: A Mini-Review. *Frontiers in nutrition*, 5, 87. Doi:10.3389/fnut.2018.00087
- Costa, C., Tsatsakis, A., Mamoulakis, C., Teodoro, M., Briguglio, G., Caruso, E., . . . Kouretas, D. (2017). Current evidence on the effect of dietary polyphenols intake on chronic diseases. *Food and Chemical Toxicology*, 110, 286-299.
- Crop Production. (2019). In: United States Department of Agriculture (USDA)
- Csiki, Z., Papp-Bata, A., Czompa, A., Nagy, A., Bak, I., Lekli, I., . . . Tosaki, A. (2015). Orally Delivered Sour Cherry Seed Extract (SCSE) Affects Cardiovascular and Hematological Parameters in Humans: TOXICOLOGICAL EVALUATION OF SOUR CHERRY SEED EXTRACT. *Phytotherapy Research*, 29(3), 444-449. Doi:10.1002/ptr.5273
- Damar, İ., & Ekşi, A. (2012). Antioxidant capacity and anthocyanin profile of sour cherry (*Prunus cerasus* L.) juice. *Food chemistry*, 135(4), 2910-2914.
- Derrien, M., Belzer, C., & de Vos, W. M. (2017). Akkermansia muciniphila and its role in regulating host functions. *Microbial pathogenesis*, 106, 171-181.
- Edwards, C. A., Havlik, J., Cong, W., Mullen, W., Preston, T., Morrison, D. J., & Combet, E. (2017). Polyphenols and health: Interactions between fibre, plant polyphenols and the gut microbiota. *Nutrition bulletin*, 42(4), 356-360.

- Espín, J. C., González-Sarrías, A., & Tomás-Barberán, F. A. (2017). The gut microbiota: A key factor in the therapeutic effects of (poly) phenols. *Biochemical pharmacology*, 139, 82-93.
- Everard, A., Belzer, C., Geurts, L., Ouwerkerk, J. P., Druart, C., Bindels, L. B., . . . Delzenne, N. M. (2013). Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity. *Proceedings of the National Academy of Sciences*, 110(22), 9066-9071.
- Ferretti, G., Bacchetti, T., Belleggia, A., & Neri, D. (2010). Cherry antioxidants: from farm to table. *Molecules (Basel, Switzerland)*, 15(10), 6993-7005.  
Doi:10.3390/molecules15106993
- Fielding, R. A., Reeves, A. R., Jasuja, R., Liu, C., Barrett, B. B., & Lustgarten, M. S. (2019). Muscle strength is increased in mice that are colonized with microbiota from high-functioning older adults. *Experimental gerontology*, 127, 110722.
- Gao, R., & Chilibeck, P. D. (2020). Effect of Tart Cherry Concentrate on Endurance Exercise Performance: A Meta-analysis. *Journal of the American College of Nutrition*, 1-8.
- Garcia-Mazcorro, J. F., Lage, N. N., Mertens-Talcott, S., Talcott, S., Chew, B., Dowd, S. E., . . . Noratto, G. D. (2018). Effect of dark sweet cherry powder consumption on the gut microbiota, short-chain fatty acids, and biomarkers of gut health in obese db/db mice. *PeerJ*, 6, e4195.
- Gonçalves, A. C., Bento, C., Jesus, F., Alves, G., & Silva, L. R. (2018). Sweet Cherry Phenolic Compounds: Identification, Characterization, and Health Benefits. In *Studies in Natural Products Chemistry* (Vol. 59, pp. 31-78): Elsevier.
- Gowd, V., Karim, N., Shishir, M. R. I., Xie, L., & Chen, W. (2019). Dietary polyphenols to combat the metabolic diseases via altering gut microbiota. *Trends in Food Science & Technology*.
- Hammer, Ø., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia electronica*, 4(1), 9.
- Han, Y., & Xiao, H. (2020). Whole Food–Based Approaches to Modulating Gut Microbiota and Associated Diseases. *Annual Review of Food Science and Technology*, 11, 119-143.
- Howatson, G., Bell, P. G., Tallent, J., Middleton, B., McHugh, M. P., & Ellis, J. (2012). Effect of tart cherry juice (*Prunus cerasus*) on melatonin levels and enhanced sleep quality. *European journal of nutrition*, 51(8), 909-916.
- Hughenholz, F., & de Vos, W. M. (2018). Mouse models for human intestinal microbiota research: a critical evaluation. *Cellular and Molecular Life Sciences*, 75(1), 149-160.

- Juhasz, B., Kertész, A., Balla, J., Balla, G., Szabo, Z., Bombicz, M., . . . D Haines, D. (2013). Cardioprotective effects of sour cherry seed extract (SCSE) on the hypercholesterolemic rabbit heart. *Current Pharmaceutical Design*, 19(39), 6896-6905.
- Kalt, W., Cassidy, A., Howard, L. R., Krikorian, R., Stull, A. J., Tremblay, F., & Zamora-Ros, R. (2020). Recent research on the health benefits of blueberries and their anthocyanins. *Advances in nutrition*, 11(2), 224-236.
- Kelley, D. S., Adkins, Y., & Laugero, K. D. (2018). A Review of the Health Benefits of Cherries. *Nutrients*, 10(3), 368. Doi:10.3390/nu10030368
- Kent, K., Charlton, K., Roodenrys, S., Batterham, M., Potter, J., Traynor, V., . . . Richards, R. (2017). Consumption of anthocyanin-rich cherry juice for 12 weeks improves memory and cognition in older adults with mild-to-moderate dementia. *European journal of nutrition*, 56(1), 333-341.
- Khoo, G. M., Clausen, M. R., Pedersen, B. H., & Larsen, E. (2011). Bioactivity and total phenolic content of 34 sour cherry cultivars. *Journal of Food Composition and Analysis*, 24(6), 772-776. Doi:10.1016/j.jfca.2011.03.004
- Kim, D.-O., Heo, H. J., Kim, Y. J., Yang, H. S., & Lee, C. Y. (2005). Sweet and Sour Cherry Phenolics and Their Protective Effects on Neuronal Cells. *Journal of agricultural and food chemistry*, 53(26), 9921-9927. Doi:10.1021/jf0518599
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Applied and Environmental Microbiology*, 79(17), 5112-5120. Doi:10.1128/AEM.01043-13
- Kumar Singh, A., Cabral, C., Kumar, R., Ganguly, R., Kumar Rana, H., Gupta, A., . . . Pandey, A. K. (2019). Beneficial effects of dietary polyphenols on gut microbiota and strategies to improve delivery efficiency. *Nutrients*, 11(9), 2216.
- Lachin, T. (2014). Effect of antioxidant extract from cherries on diabetes. *Recent patents on endocrine, metabolic & immune drug discovery*, 8(1), 67-74.
- Lavefve, L., Howard, L. R., & Carbonero, F. (2020). Berry polyphenols metabolism and impact on human gut microbiota and health. *Food & Function*, 11(1), 45-65.
- Lear, R., O'Leary, M., O'Brien Andersen, L., Holt, C. C., Stensvold, C. R., van der Giezen, M., & Bowtell, J. L. (2019). Tart cherry concentrate does not alter the gut microbiome, glycaemic control or systemic inflammation in a middle-aged population. *Nutrients*, 11(5), 1063.
- Levers, K., Dalton, R., Galvan, E., O'Connor, A., Goodenough, C., Simbo, S., . . . Riechman, S. (2016). Effects of powdered Montmorency tart cherry supplementation on acute endurance exercise performance in aerobically trained individuals. *Journal of the International Society of Sports Nutrition*, 13(1), 22.



- Li, Z., Summanen, P. H., Komoriya, T., Henning, S. M., Lee, R.-P., Carlson, E., . . . Finegold, S. M. (2015). Pomegranate ellagitannins stimulate growth of gut bacteria in vitro: Implications for prebiotic and metabolic effects. *Anaerobe*, 34, 164-168.
- Liu, Z., Liu, H.-Y., Zhou, H., Zhan, Q., Lai, W., Zeng, Q., . . . Xu, D. (2017). Moderate-intensity exercise affects gut microbiome composition and influences cardiac function in myocardial infarction mice. *Frontiers in microbiology*, 8, 1687.
- Lynn, A., Mathew, S., Moore, C. T., Russell, J., Robinson, E., Soumpasi, V., & Barker, M. E. (2014). Effect of a tart cherry juice supplement on arterial stiffness and inflammation in healthy adults: a randomised controlled trial. *Plant foods for human nutrition*, 69(2), 122-127.
- Marín, L., Miguélez, E. M., Villar, C. J., & Lombó, F. (2015). Bioavailability of dietary polyphenols and gut microbiota metabolism: antimicrobial properties. *BioMed research international*, 2015.
- Mayta-Apaza, A. C., Marasini, D., & Carbonero, F. (2017). Tart Cherries and Health: Current Knowledge and Need for a Better Understanding of the Fate of Phytochemicals in Human Gastrointestinal Tract. *Crit. Rev. Food Sci. Nutr*, 28, 1-13.
- Mayta-Apaza, A. C., Pottgen, E., De Bodt, J., Papp, N., Marasini, D., Howard, L., . . . Carbonero, F. (2018). Impact of tart cherries polyphenols on the human gut microbiota and phenolic metabolites in vitro and in vivo. *The Journal of nutritional biochemistry*, 59, 160-172.
- McCune, L. M., Kubota, C., Stendell-Hollis, N. R., & Thomson, C. A. (2010). Cherries and Health: A Review. *Critical Reviews in Food Science and Nutrition*, 51(1), 1-12. Doi:10.1080/10408390903001719
- Munukka, E., Ahtiainen, J. P., Puigbó, P., Jalkanen, S., Pahkala, K., Keskitalo, A., . . . Elo, L. (2018). Six-week endurance exercise alters gut metagenome that is not reflected in systemic metabolism in over-weight women. *Frontiers in microbiology*, 9, 2323.
- Nagpal, R., Wang, S., Solberg Woods, L. C., Seshie, O., Chung, S. T., Shively, C. A., . . . Yadav, H. (2018). Comparative microbiome signatures and short-chain fatty acids in mouse, rat, non-human primate, and human feces. *Frontiers in microbiology*, 9, 2897.
- Nash, V., Ranadheera, C. S., Georgousopoulou, E. N., Mellor, D. D., Panagiotakos, D. B., McKune, A. J., . . . Naumovski, N. (2018). The effects of grape and red wine polyphenols on gut microbiota—A systematic review. *Food research international*, 113, 277-287.
- Nemes, A., Homoki, J. R., Kiss, R., Hegedűs, C., Kovács, D., Peitl, B., . . . Remenyik, J. (2019). Effect of Anthocyanin-Rich Tart Cherry Extract on Inflammatory Mediators and Adipokines Involved in Type 2 Diabetes in a High Fat Diet Induced Obesity Mouse Model. *Nutrients*, 11(9), 1966. Doi:10.3390/nu11091966

- Nguyen, T. L. A., Vieira-Silva, S., Liston, A., & Raes, J. (2015). How informative is the mouse for human gut microbiota research? *Disease models & mechanisms*, 8(1), 1-16.
- Ozdal, T., Sela, D. A., Xiao, J., Boyacioglu, D., Chen, F., & Capanoglu, E. (2016). The reciprocal interactions between polyphenols and gut microbiota and effects on bioaccessibility. *Nutrients*, 8(2), 78.
- Perez, A., Ferreira, G., & Minor, T. (2017). Fruit and Tree Nuts Outlook. Retrieved from <https://www.ers.usda.gov/webdocs/outlooks/85287/fts-365.pdf?v=9364.6>
- Peron, G., Hidalgo-Liberona, N., González-Domínguez, R., Garcia-Aloy, M., Guglielmetti, S., Bernardi, S., . . . Riso, P. (2019). Exploring the molecular pathways behind the effects of nutrients and dietary polyphenols on gut microbiota and intestinal permeability: a perspective on the potential of metabolomics and future clinical applications. *Journal of agricultural and food chemistry*.
- Santhakumar, A. B., Battino, M., & Alvarez-Suarez, J. M. (2018). Dietary polyphenols: Structures, bioavailability and protective effects against atherosclerosis. *Food and Chemical Toxicology*, 113, 49-65.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., . . . Robinson, C. J. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology*, 75(23), 7537-7541.
- Tomás-Barberán, F. A., Selma, M. V., & Espín, J. C. (2016). Interactions of gut microbiota with dietary polyphenols and consequences to human health. *Current opinion in clinical nutrition and metabolic care*, 19(6), 471-476.
- Tu, P., Bian, X., Chi, L., Gao, B., Ru, H., Knobloch, T. J., . . . Lu, K. (2018). Characterization of the functional changes in mouse gut microbiome associated with increased *Akkermansia muciniphila* population modulated by dietary black raspberries. *ACS omega*, 3(9), 10927-10937.
- Van Passel, M. W. J., Kant, R., Zoetendal, E. G., Plugge, C. M., Derrien, M., Malfatti, S. A., . . . de Vos, W. M. (2011). The genome of *Akkermansia muciniphila*, a dedicated intestinal mucin degrader, and its use in exploring intestinal metagenomes. *PloS one*, 6(3).
- Vitale, K. C., Hueglin, S., & Broad, E. (2017). Tart cherry juice in athletes: a literature review and commentary. *Current sports medicine reports*, 16(4), 230-239.
- Wang, H., Nair, M. G., Iezzoni, A. F., Strasburg, G. M., Booren, A. M., & Gray, J. I. (1997). Quantification and characterization of anthocyanins in Balaton tart cherries. *Journal of agricultural and food chemistry*, 45(7), 2556-2560.
- Wankhade, U. D., Zhong, Y., Lazarenko, O. P., Chintapalli, S. V., Piccolo, B. D., Chen, J.-R., & Shankar, K. (2019). Sex-specific changes in gut microbiome composition following blueberry consumption in C57BL/6J mice. *Nutrients*, 11(2), 313.

- Wojdyło, A., Nowicka, P., Laskowski, P., & Oszmiański, J. (2014). Evaluation of Sour Cherry (*Prunus cerasus* L.) Fruits for Their Polyphenol Content, Antioxidant Properties, and Nutritional Components. *Journal of Agricultural and Food Chemistry*, 62(51), 12332-12345. Doi:10.1021/jf504023z
- Wu, S., Hu, R., Nakano, H., Chen, K., Liu, M., He, X., . . . Hou, D.-X. (2018). Modulation of gut microbiota by *Lonicera caerulea* L. Berry polyphenols in a mouse model of fatty liver induced by high fat diet. *Molecules*, 23(12), 3213.
- Zhang, Z., Wu, X., Cao, S., Cromie, M., Shen, Y., Feng, Y., . . . Li, L. (2017). Chlorogenic acid ameliorates experimental colitis by promoting growth of *Akkermansia* in mice. *Nutrients*, 9(7), 677.

## CHAPTER 3:

### Impact of Cranberry Juice Consumption on Gut and Vaginal Microbiota in Postmenopausal Women

Ayoub Al Othaim<sup>1,2</sup>, Daya Marasini<sup>3,†</sup>, and Franck Carbonero<sup>1,3,4,5\*</sup>

<sup>1</sup>Cell and Molecular Biology Program; University of Arkansas, Fayetteville, AR, 72701 USA; franck.carbonero@wsu.edu

<sup>2</sup>Department of Medical Laboratories, College of Applied Medical Sciences, Majmaah University, Al-Majmaah, 11952, Saudi Arabia

<sup>3</sup>Department of Food Science; University of Arkansas, Fayetteville, AR, 72704 USA

<sup>4</sup>Department of Food Science, Washington State University

<sup>5</sup>Department of Nutrition and Exercise Physiology, Elson Floyd School of Medicine, Washington State University-Spokane, 412 East Spokane Falls Boulevard, Spokane, WA, 99202 USA

\* Correspondence: franck.carbonero@wsu.edu

† Current affiliation: IHRC INC, Atlanta, GA, United States of America.

#### Abstract

Cranberries have long been purported to provide protection against urinary tract infections (UTIs). There is a line of evidence suggesting that causal pathogens might be seeded from the bacteria reservoirs in the intestinal and vaginal tracts. We tested the hypothesis whether cranberry intake would reshape bacteria taxa in the gut, as well as the vaginal ecosystem. A total of 25 postmenopausal women were enrolled into a randomized, double-blind, placebo-controlled study. Stool samples and vaginal swabs were collected at baseline and after 15 days of consumption of placebo or cranberry beverages, microbiota analyses were performed by Illumina Miseq sequencing following a double index 16S rRNA gene amplicon. All baseline stool samples generally fell in the *Bacteroides* enterotype. Significant increases of *Prevotella* ( $P = 0.04$ ), *Clostridium* XIVa members ( $P = 0.04$ ), *Eggerthella* ( $P = 0.03$ ), and *Bifidobacterium* ( $P =$

0.02) were shown following the cranberry juice intervention; this indicates modulation of the gut microbiota by cranberry components. Baseline vaginal microbiotas fell in three distinct patterns, *Lactobacillus* dominant, diversified microbiome, and *Streptococcus* dysbiosis. Compared with the placebo, the cranberry intervention significantly reduced the abundance of pathogenic *Streptococcus* ( $P = 0.04$ ) in the dysbiosis group and increased commensal bacteria *Anaerococcus*, *Finegoldia*, *Actinomyces*, and *Corynebacterium* in the diversified microbiome and dysbiosis groups. Overall, these data suggest that cranberry consumption may improve vaginal microbiota composition in individuals with dysbiosis. Gut-borne taxa stimulation by the combination of cranberry oligosaccharides and polyphenols present in the cranberry product potentially mediates these beneficial properties.

**Keywords:** Cranberries; Urinary Tract Infection, Gut microbiota, Vaginal microbiota; Dysbiosis

### 3.1 Introduction

Urinary tract infections (UTIs) are one of the most common bacterial infections among women in general, and postmenopausal women are at an increased risk of recurrent infections (Fu, Liska, Talan, & Chung, 2017; Jung & Brubaker, 2019). It is estimated that more than 50% of women will have at least one occurrence of a UTI during their life spans and approximately 20–30% are affected with recurrent infections (Beerepoot & Geerlings, 2016; Fu et al., 2017). While UTIs are generally benign, their prevalence poses medical and economic issues resulting in millions of visits yearly and an estimated cost of \$2.8 billion in medical care expenses in the United States (Simmering, Tang, Cavanaugh, Polgreen, & Polgreen, 2017). In addition, the need for antibiotic treatment is problematic both for the patient's natural microbiota stability and for the community and environment, since antibiotic-resistance can result from such treatments

(Ianiro, Tilg, & Gasbarrini, 2016; Koutzoumis et al., 2020; Morgun et al., 2015; Stapleton, 2016).

While pathogenic bacteria, most often *Escherichia coli* and *Streptococcus*, are the main UTI causative bacteria, it has been proposed that a reduction in the abundance of commensal vaginal microbes may also contribute to an increased UTI risk in postmenopausal women (Reid, 2018; Stapleton, 2016). It is now well understood that the normal vaginal microbiome is dominated by *Lactobacilli*, and the vaginal tract is naturally populated with a relatively dense and diverse collection of bacteria (Mendling, 2016; Ravel et al., 2011; White, Creedon, Nelson, & Wilson, 2011; Witkin & Linhares, 2017). An imbalance of the vaginal microbiome is not only a common cause for bacterial vaginosis (BV) (Stapleton, 2016; White et al., 2011) but may directly affect the potential pathogens to colonize the urethra and/or bladder (Czaja et al., 2009; Stapleton, 2016). Thus, it would appear that asymptomatic changes in the vaginal microbiome may trigger UTI development, either by seeding pathogens or by creating a niche for pathogens to thrive (Czaja et al., 2009; Stapleton, 2016).

The factors shaping the vaginal microbiota composition and stability are still not well known (Braundmeier et al., 2015). In this context, we hypothesize that the vaginal microbiome is strongly influenced by the gut microbiome. There are different lines of evidence to support this hypothesis. First, because of the anatomical proximity, gut microbes in the rectum and distal colon can easily come in contact with the vaginal tract, and indeed several vaginal *Lactobacillus* species are also found in the digestive tract (Marchesi, Silva, Wiese, & Nader-Macías, 2020; Pan, Hidalgo-Cantabrana, Goh, Sanozky-Dawes, & Barrangou, 2020). Second, it has become increasingly evident that metabolites produced by the gut microbiome are highly bioavailable and distributed to any organ/region of the human body and these metabolites may influence the

vaginal ecosystem as well (Possemiers, Bolca, Verstraete, & Heyerick, 2011). Since the gut microbiome is strongly individualized, shaped by external factors, and naturally evolving to some extent, this would in part explain why the underlying causes of UTI and BV are often challenging to identify.

It is plausible that the different responses to cranberry intervention might be the consequence of inter-individual differences in gut microbiome composition. During the last decade, reports of potential positive modulation of the human gut microbiota by berries have emerged (Cai et al., 2019; Lavefve, Howard, & Carbonero, 2020; O'Connor et al., 2019; Rodríguez-Morató, Matthan, Liu, de la Torre, & Chen, 2018). Polymeric cranberry phenolic compounds generally have poor bioavailability. However, when they reach the distal intestine, those molecules are broken down and metabolized by microbiota under anaerobic conditions, a process known as fermentation. Compared to their parent compounds, these bacteria-derived metabolites are highly bioavailable and bioactive. Previous studies showed that  $\gamma$ -valerolactone and its sulphate conjugates (Mena et al., 2017) phenylacetic acid, 3,4-dihydroxyphenylacetic acid (González de Llano et al., 2015), hippuric and  $\alpha$ -hydroxyhippuric acids (González de Llano, Liu, Khoo, Moreno-Arribas, & Bartolomé, 2019) have anti-bacteria effects under experimental conditions. These findings highlight the critical need to understand how cranberries could affect the possible crosstalk between intestinal and other microbiome ecosystems, thereby modulating the impact of dietary cranberries on UTI prevention. This project aimed to assessing the relationships between the gut microbiome and vaginal microbiome, and how cranberry consumption affecting the gut microbiome may also impact the vaginal microbiome. It is expected that in the long term, nutritional studies using other dietary elements (such as prebiotics) may be designed to optimize and maximize the beneficial effects of cranberries.

## 3.2 Materials and Methods

### 3.2.1 Subjects

Twenty-three (23) women volunteered and participated in this study. The eligibility (no menstrual period for at least 12 months) of the participants was determined using a questionnaire. The participants were postmenopausal women between the ages of 50-75 years old who met the inclusion criteria including no recent antibiotics or immunosuppressive therapies; surgery of the stomach, small or large intestines; appendectomy, gastric bypass or gastric banding in the previous 6 months, and diagnosis with any autoimmune diseases.

### 3.2.2 Study Design and sample collection

This study was a randomized, placebo-controlled, crossover, dietary intervention. Ocean Spray Cranberries, Inc. provided the experimental and placebo beverages, which have been validated and used in previous studies (Hsia, Zhang, Beyl, Greenway, & Khoo, 2020; Haiyan Liu, Howell, Zhang, & Khoo, 2019; Maki et al., 2016; Zhao, Liu, Su, Khoo, & Gu, 2020). Both beverages were similar in appearance, taste, and aroma and assigned to volunteers according to computer-generated random orders. Both investigators and participants were blind to the assignment and the products. The products were identified by a random 3-digit code pre-printed on the cap. The participants consumed either cranberry juice or the placebo beverage daily (8 fl oz per day) for 15 days (**Table 3. 1**). Analytical methods for testing the products were previously described (Maki et al., 2016). Biological samples were collected at baseline and after each intervention (**Figure 3. 1**). Subjects were provided with polyester tipped swabs with a neutralizing buffer solution using the Environmental Sampling Kit (Puritan Diagnostics, USA) for collecting vaginal samples and the Precision™ Stool Collector (Cardinal Health, USA) for



collecting stool samples. They were also provided with instructions for self-sampling at each sampling event. Vaginal swabs and stool samples were stored at  $-20^{\circ}\text{C}$  until used for microbiome analysis.

### **3.2.3 DNA extraction**

Microbial DNA was extracted from all samples with the QIAamp DNA stool kit (QIAGEN, Valencia, CA) following the manufacturer's protocol with the addition of cell lysis step to rapidly and completely disrupt cells in the samples to release DNA. The cell lysis was performed by adding 0.1 g of 0.1 mm diameter and 0.1 g of 0.5 mm diameter zirconia-silica beads (BioSpec Products, Bartlesville, OK, United States) to 2 ml tubes. Stool and vaginal samples were transferred to the 2 ml tubes containing zirconia-silica beads. One ml of InhibitEX Buffer (from the QIAamp® Fast DNA Stool Mini Kit) was then added to the tubes. The tubes were subjected to a bead-beater for 60 seconds at maximum speed using a FastPrep®-24 bead-beater (MP Biomedicals, Santa Ana, CA, United States).

The NanoDrop™ 1000 Spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA, United States) was used to determine the DNA concentration. Extracted DNA was then visualized following electrophoresis on a 2% agarose gel in 1X TAE (Tris-acetate ethylenediaminetetraacetic acid) buffer (AMRESCO®, Cleveland, OH, United States).

### **3.2.4 Universal polymerase chain reaction**

16S ribosomal RNA gene was amplified by PCR ( $95^{\circ}\text{C}$  for 3 min, followed by 35 cycles at  $98^{\circ}\text{C}$  for 30 s,  $55^{\circ}\text{C}$  for 30 s, and  $72^{\circ}\text{C}$  for 1 min and a final extension at  $72^{\circ}\text{C}$  for 5 min) using primers (8F 5' – AGAGTTTGATCCTGGCTCAG – 3' and 1541R 5' – AAGGAGGTGATCCAGCCGCA – 3') (Carbonero, Oakley, & Purdy, 2014). PCR reactions

were performed in 25  $\mu$ L mixture containing 3  $\mu$ L of DNA template; 1  $\mu$ L of each universal primer (8F and 1541R); 12.5  $\mu$ L of GoTaq® Green Master Mix (Promega™ Corporation, Wisconsin, USA); and 7.5  $\mu$ L of Nuclease-Free Water. Amplicons were run on agarose gel electrophoresis to confirm the success of the PCR by using 1  $\mu$ L of SYBR safe DNA Gel Stain (Thermo Fisher Scientific, USA) fluorescent dye.

### **3.2.5 Libraries Preparation and Sequencing**

For 16S rRNA gene sequencing analyses, a dual-indexed amplicon strategy was performed using universal primers tagged with Illumina adapters and barcodes as previously described (Aljahdali et al., 2017; Kozich, Westcott, Baxter, Highlander, & Schloss, 2013; Mayta-Apaza et al., 2018). Briefly, PCR was set up in 25  $\mu$ L reaction components with: 3  $\mu$ L of the DNA template; 2.5  $\mu$ L of the AccuPrime™ PCR buffer II; 0.5  $\mu$ L of each index primers; 0.1  $\mu$ L of AccuPrime™ Taq DNA Polymerase (Invitrogen, USA); and 18.4  $\mu$ L of Nuclease-Free Water following the recommended protocol. The PCR consisted of initial denaturation at 95 °C for 3 min, followed by 30 cycles of denaturation at 95 °C for 30 sec, primer annealing at 55 °C for 30 sec, and an extension at 72 °C for 1 min, with a final extension step at 72 °C for 5 min. Correct size amplicons were subjected to purification and normalization with the Invitrogen SequalPrep kits following the manufacturer's protocol. Amplicons were pooled into two different libraries and concentrations were measured by Agilent bioanalyzer and qPCR using the same primers and PerfeCta NGS library quantification kits (Quanta Biosciences, USA). Libraries were then denatured and diluted to appropriate concentrations and sequenced on an Illumina MiSeq.

Resulting sequencing files were analyzed using the Mothur software. Operational taxonomic units (OUT) were determined and classified against reference databases (Ribosomal

Database Project (RDP), SILVA, GenBank). Basic and multivariate statistical analyses were performed in PAST.

### **3.2.6 Statistical analyses**

Statistical analyses including Wilcoxon signed-rank test and Mann-Whitney U test (depending on the availability of fully paired sample sets) were performed to detect significant differences between groups (by convention, differences were considered significant when  $P < 0.05$ ). Non-metric multidimensional scaling (NMDS) and analysis of similarities (ANOSIM) based on count distance metrics (Bray-Curtis similarity index; ANOSIM:  $P < 0.05$ ) were obtained with Past4 software (Al Othaim, Marasini, & Carbonero, 2020; Hammer, Harper, & Ryan, 2001).

## **3.3 Results and Discussion**

The human study was completed with 23 volunteers, and 21 of which provided the full set of samples. The volunteers' demographics are described in **Table 3. 2**.

For the baseline samples, 23 stool and 23 vaginal swabs were collected, 22 of each for the first period samples, and 21 of each for the second period. Therefore, a total of 66 stool samples and 66 vaginal swabs have been collected. However, for several subjects, self-sampling of stool resulted in problematic samples for at least one of the three samples. As these samples did not yield satisfactory levels of microbial DNA, only eight subjects with the full set of samples were included in the analyses.

### 3.3.1 Impact of the cranberry juice on the gut microbiota

Cranberry juice significantly increased the numbers of *Bifidobacterium* ( $P = 0.02$ ) and *Eggerthella* ( $P = 0.03$ ), two Actinobacteria members that have been shown previously to be able to grow and metabolize dietary polyphenols (Fogliano et al., 2011; Koutsos & Lovegrove, 2015; Morais, de Rosso, Estadella, & Pisani, 2016; Queipo-Ortuño et al., 2012). *Bifidobacterium* are also known as probiotics and the increased abundance is considered a sign of improved intestinal health (Cuervo et al., 2014; Rajkumar et al., 2014; Singh et al., 2017). Cranberry juice also resulted in a significant increase of *Prevotella* ( $P = 0.04$ ), which is considered a beneficial genus associated with polysaccharides and fiber consumption (David et al., 2014; De Filippis et al., 2016; Wu et al., 2011) (**Figure 3. 2**).

The most remarkable trend among the Firmicutes was a significant increase of *Clostridium* XIVa ( $P = 0.04$ ) and other known butyrate producers (*Eubacterium*, *Flavonifractor*, and *Subdoligranulum*) after the consumption of cranberry juice (El Aidy, Van den Abbeele, Van de Wiele, Louis, & Kleerebezem, 2013; Rivière, Selak, Lantin, Leroy, & De Vuyst, 2016; Walker, Duncan, Louis, & Flint, 2014) (**Figure 3. 2**). Butyrate is a short-chain fatty acid shown to offer benefits to not only the local intestinal environment, but also contributes to systemic metabolism (Hu Liu et al., 2018; O’Keefe, 2016). *Akkermansia*, a genus that has been associated with a number of health properties, was increased with cranberry consumption in animal models (Anhê et al., 2017; Rodríguez-Daza et al., 2020) and to some extent in humans (Bekiaries, Krueger, Meudt, Shanmuganayagam, & Reed, 2018), it showed a slight but non-significant increase between cranberry and placebo beverages (Dao et al., 2016; Png et al., 2010) (**Figure 3. 2**). It should be noted that all those trends were strongly individualized, and further validation is

warranted in more clinical studies with a larger sample size. Overall, cranberry consumption appears to stimulate beneficial genera.

### **3.3.2 Impact of the cranberry juice on the vaginal microbiota**

The vaginal microbiotas were found to be extremely heterogeneous at baseline, and have therefore been stratified into three groups based on the most abundant genera and their known relation to vaginal health (**Figure 3. 3**):

Group 1 comprises individuals whose baseline vaginal microbiotas were strongly dominated by *Lactobacillus*, which is considered the ideal healthy state (Ravel et al., 2011). Group 2 comprises individuals whose baseline vaginal microbiotas were characterized by diverse and relatively well distributed high number of different taxa, without obvious dominant detrimental genus. Group 3 comprises individuals whose baseline vaginal microbiotas were characterized by high abundance of *Streptococcus*, which is strongly indicative of microbial dysbiosis that could potentially lead to vaginosis or UTI (Leclercq et al., 2016; Tan, Ulett, Steele, Benjamin, & Ulett, 2012; Ulett et al., 2009). Individuals were roughly equally distributed among the 3 groups which allowed statistical analyses for each group.

#### **3.3.2.1 *Lactobacillus* dominant vaginal microbiota**

Consumption of both products appeared to have little impact on the vaginal microbiota of volunteers that have *Lactobacillus* as the major bacterial taxa (**Figure 3. 4**).

#### **3.3.2.2 *Diverse baseline vaginal microbiota***

As expected with the criterion used to segregate the groups, the vaginal microbiota of these subjects was significantly more diverse (**Figure 3. 5**). The number of taxa was significantly

decreased after the consumption of cranberry juice, and even more with the placebo. However, the decrease of the Shannon index was only significant in the placebo group. It can be hypothesized that loss of diversity not associated with *Lactobacillus* dominance may be detrimental to some extent; but the loss of the diversity does not appear to be different between the interventions and thus, it is likely not associated with cranberry components.

Several significant changes were observed in the bacteria abundance at genus level (**Figure 3. 6**). Compared with baseline, consumption of cranberry juice allowed for the maintenance of several moderately abundant Firmicutes (*Lactobacillus*, *Anaerococcus*, *Finegoldia*, and *Peptoniphilus*) and Actinobacteria (*Actinomyces*, *Varibaculum*, *Actinomycetales*, and *Corynebacterium*). In contrast, aforementioned bacteria were significantly reduced following placebo beverage consumption. Different responses between the cranberry and placebo beverage from baseline were statistically significant. The status of these genera is somewhat controversial, but they would be considered as mostly harmless to vaginal health (Boskey, 2001; Ma, Forney, & Ravel, 2012). The placebo beverage led to an unexplainable increase of *Prevotella*, which was different from the response following cranberry beverage intake. *Prevotella* in the gut microbiota are associated with plant-based diets with a possible role in digestion of complex carbohydrates and production of short-chain fatty acids (Geva-Zatorsky et al., 2017; Precup & Vodnar, 2019; Zhu, Sunagawa, Mende, & Bork, 2015). However, the function of *Prevotella* in the vaginal tract is less clear. Some lines of evidence suggested that high levels of SCFAs might promote dysbiosis and inflammation in the vaginal tract (Aldunate et al., 2015; Amabebe & Anumba, 2020). To conclude, cranberry juice appears to maintain the diversity of vaginal microbiota which is a presumably healthier profile for these volunteers.

### 3.3.2.3 *Streptococcus dysbiosis*

Several women ( $n = 7$ ) were found to have a strong dysbiosis of their vaginal microbiota, characterized by a high initial abundance 22% to 81% of *Streptococcus*, a well-known causative agent of vaginosis and UTI (Jass & Reid, 2009; Leclercq et al., 2016; Tan et al., 2012; Ulett et al., 2009).

As suggested by the diversity trends, cranberry juice had a very strong and specific impact on the dysbiosed vaginal microbiota (**Figure 3. 7**). Most notably, although both beverages reduced the abundance of *Streptococcus* compared to baseline, the consumption of cranberry juice significantly reduced the relative abundance of *Streptococcus* to a greater extent, compared to the placebo beverage. Intriguingly, it appeared that *Streptococcus* has been replaced with a range of different Firmicutes (*Anaerococcus* and *Finegoldia*) and Actinobacteria (*Actinobaculum*, *Actinomyces*, *Varibaculum*, and *Corynebacterium*; **Figure 3. 8**). *Streptococcus*, both group A and B, colonization has been associated with an increased risk of vaginitis, bacterial vaginosis, through the female life span (Clark & Atendido, 2005; Verstraelen, Verhelst, Vaneechoutte, & Temmerman, 2011). The presence of this genus also significantly increased chances of vertical transmission to neonates during pregnancy (Puopolo, Lynfield, Cummings, & Committee On Infectious, 2019). The exact mechanism of the reduction of the abundance of *Streptococcus* by cranberries is unknown. However, Feliciano and colleagues showed that cranberry proanthocyanidins (PAC) reduced extra-intestinal pathogenic *E. coli* agglutination and inhibited the invasion of enterocytes (Feliciano, Meudt, Shanmuganayagam, Krueger, & Reed, 2014). Cranberry-derived PACs might impair virulence and quorum sensing of *Pseudomonas aeruginosa*, according to another study (Maisuria, Lopez-de Los Santos, Tufenkji, & Déziel, 2016). In the research performed by O'May and Tufenkji, cranberry PAC and other tannin-

containing materials blocked swarming motility of *P. aeruginosa*, possibly via inhibition of the flagellin gene expression (O'May & Tufenkji, 2011). It can be hypothesized that cranberries and metabolites generated and absorbed in the gut affected *Streptococcus* through multiple mechanisms, inhibition of bacteria growth or viability, disruption of cytoskeleton structure, and etc. These hypotheses will warrant future experiments to explore.

### 3.4 Conclusion

These findings suggest a beneficial impact of cranberry juice on vaginal microbiota with dysbiosis, and possibly its contribution to vaginal health. It can be speculated that *Streptococcus* dysbiosis could be improved directly within the local niche environment by increasing competition of other commensal bacteria or indirectly by decreasing the bacteria migration from the gut reservoir, and thereby restoring a healthier vaginal microbiota. Due to that possible relationship, the beneficial impact of cranberry juice on the gut microbiota is also encouraging for vaginal health. As reported previously, the ageing gut microbiota tends to be less diverse and responsive to dietary changes. Herein, a relatively short-term consumption and daily consumption of cranberry juice shows promising prebiotic effect and may help maintain levels of a wide range of beneficial genera (*Prevotella*, *Bifidobacterium*, and butyrate-producing Firmicutes).

**Acknowledgement:** Funding and cranberry products were provided by Ocean Spray Cranberries, Inc.

**Author Contributions:** Conceptualization, Franck Carbonero; Data curation, Ayoub Al Othaim, Daya Marasini and Franck Carbonero; Formal analysis, Ayoub Al Othaim, Daya Marasini and Franck Carbonero; Funding acquisition, Franck Carbonero; Methodology, Ayoub Al Othaim,




Daya Marasini and Franck Carbonero; Project administration, Daya Marasini and Franck Carbonero; Supervision, Daya Marasini and Franck Carbonero; Visualization, Franck Carbonero; Writing – original draft, Ayoub Al Othaim and Franck Carbonero; Writing – review & editing, Ayoub Al Othaim, Daya Marasini and Franck Carbonero.

**Table 3. 1:** The analytical data for the cranberry and placebo beverages. ND indicates non-detectable


Analytical	Cranberry juice	Placebo	Unit
Anthocyanins	3.09	ND	mg
Flavonols	5.68	ND	mg
Phenolic acids	11.60	ND	mg
Total Phenolics	161	ND	mg
Proanthocyanidins	141	ND	mg
Organic acid	1.74	1.67	g
Sugar	5.94	6.3	g

**Table 3. 2:** Volunteers demographics.


<b>Age (years)</b>	61±7 (min=51; max=74)
<b>BMI</b>	28.5±7.8 (min=18; max=46)
<b>Height (inches)</b>	64.2±2.3 (min=60.2; max=72.2)
<b>Weight (lbs)</b>	170.5±49.7 (min=112; max=278)




**Figure 3. 1:** Timeline of intervention.




**Figure 3. 2:** Impact of cranberry juice and placebo on the relative abundance of genera from Actinobacteria, Bacteroidetes, Verrucomicrobia, and Firmicutes. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 8$ ). Different letters indicate significant difference ( $P < 0.05$ ), whereas N.S. indicates no significant difference.




**Figure 3. 3:** Non-metric Multidimensional Scaling (Bray-Curtis index) showing the three different clusters of baseline vaginal microbiota.




**Figure 3. 4:** Diversity indices and changes in relative abundances in volunteers with *Lactobacillus*-dominated vaginal microbiota. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 8$ ). Different letters indicate significant difference ( $P < 0.05$ ), whereas N.S. indicates no significant difference.




**Figure 3. 5:** Diversity indices changes in subjects with diverse vaginal microbiota at baseline. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 6$ ). Different letters indicate significant difference ( $P < 0.05$ ).



**Figure 3. 6:** Impact of cranberry juice and the placebo on the relative abundance of the major genera in subjects with diverse vaginal microbiota at baseline. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 6$ ). Different letters indicate significant difference ( $P < 0.05$ ), whereas N.S. indicates no significant difference.



**Figure 3. 7:** Diversity indices changes in subjects with *Streptococcus* dysbiosis. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 7$ ). Different letters indicate significant difference ( $P < 0.05$ ), whereas N.S. indicates no significant difference.



**Figure 3. 8:** Impact of cranberry juice and the placebo on the relative abundance of the major genera in subjects with *Streptococcus* dysbiosis at baseline. Values are expressed as mean  $\pm$  standard error (SE) ( $n = 7$ ). Different letters indicate significant difference ( $P < 0.05$ ), whereas N.S. indicates no significant difference.

## References

- Al Othaim, A., Marasini, D., & Carbonero, F. (2020). Impact of increasing concentration of tart and sweet cherries juices concentrates on healthy mice gut microbiota. *Food Frontiers*, 1(3), 224-233. doi:10.1002/fft2.46
- Aldunate, M., Srbinovski, D., Hearps, A. C., Latham, C. F., Ramsland, P. A., Gugasyan, R., . . . Tachedjian, G. (2015). Antimicrobial and immune modulatory effects of lactic acid and short chain fatty acids produced by vaginal microbiota associated with eubiosis and bacterial vaginosis. *Frontiers in physiology*, 6, 164.
- Aljahdali, N., Gadonna-Widehem, P., Delayre-Orthez, C., Marier, D., Garnier, B., Carbonero, F., & Anton, P. M. (2017). Repeated oral exposure to N  $\epsilon$ -carboxymethyllysine, a Maillard reaction product, alleviates gut microbiota dysbiosis in colitic mice. *Digestive Diseases and Sciences*, 62(12), 3370-3384.
- Amabebe, E., & Anumba, D. O. C. (2020). Female gut and genital tract microbiota-induced crosstalk and differential effects of short-chain fatty acids on immune sequelae. *Frontiers in immunology*, 11.
- Anhê, F. F., Nachbar, R. T., Varin, T. V., Vilela, V., Dudonné, S., Pilon, G., . . . Roy, D. (2017). A polyphenol-rich cranberry extract reverses insulin resistance and hepatic steatosis independently of body weight loss. *Molecular metabolism*, 6(12), 1563-1573.
- Beerepoot, M., & Geerlings, S. (2016). Non-antibiotic prophylaxis for urinary tract infections. *Pathogens*, 5(2), 36.
- Bekiares, N., Krueger, C. G., Meudt, J. J., Shanmuganayagam, D., & Reed, J. D. (2018). Effect of sweetened dried cranberry consumption on urinary proteome and fecal microbiome in healthy human subjects. *Omics: a journal of integrative biology*, 22(2), 145-153.
- Boskey, E. R. (2001). Vaginal acidity is produced by vaginal bacteria.
- Braundmeier, A. G., Lenz, K. M., Inman, K. S., Chia, N., Jeraldo, P., Walther-Antônio, M. R. S., . . . White, B. A. (2015). Individualized medicine and the microbiome in reproductive tract. *Frontiers in physiology*, 6, 97-97. doi:10.3389/fphys.2015.00097
- Cai, X., Han, Y., Gu, M., Song, M., Wu, X., Li, Z., . . . Xiao, H. (2019). Dietary cranberry suppressed colonic inflammation and alleviated gut microbiota dysbiosis in dextran sodium sulfate-treated mice. *Food & Function*, 10(10), 6331-6341.
- Carbonero, F., Oakley, B. B., & Purdy, K. J. (2014). Metabolic flexibility as a major predictor of spatial distribution in microbial communities. *PloS one*, 9(1), e85105.
- Clark, L. R., & Atendido, M. (2005). Group B streptococcal vaginitis in postpubertal adolescent girls. *Journal of adolescent health*, 36(5), 437-440.

- Cuervo, A., Valdés, L., Salazar, N., de los Reyes-Gavilán, C. G., Ruas-Madiedo, P., Gueimonde, M., & González, S. (2014). Pilot Study of Diet and Microbiota: Interactive Associations of Fibers and Polyphenols with Human Intestinal Bacteria. *Journal of agricultural and food chemistry*, 62(23), 5330-5336. doi:10.1021/jf501546a
- Czaja, C. A., Stamm, W. E., Stapleton, A. E., Roberts, P. L., Hawn, T. R., Scholes, D., . . . Hooton, T. M. (2009). Prospective Cohort Study of Microbial and Inflammatory Events Immediately Preceding Escherichia coli Recurrent Urinary Tract Infection in Women. *The Journal of Infectious Diseases*, 200(4), 528-536. doi:10.1086/600385
- Dao, M. C., Everard, A., Aron-Wisnewsky, J., Sokolovska, N., Prifti, E., Verger, E. O., . . . Consortium, M. I.-O. (2016). Akkermansia muciniphila and improved metabolic health during a dietary intervention in obesity: relationship with gut microbiome richness and ecology. *Gut*, 65(3), 426. doi:10.1136/gutjnl-2014-308778
- David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., . . . Turnbaugh, P. J. (2014). Diet rapidly and reproducibly alters the human gut microbiome. *Nature*, 505(7484), 559-563. doi:10.1038/nature12820
- De Filippis, F., Pellegrini, N., Vannini, L., Jeffery, I. B., La Stora, A., Laghi, L., . . . Ercolini, D. (2016). High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut*, 65(11), 1812-1821. doi:10.1136/gutjnl-2015-309957
- El Aidy, S., Van den Abbeele, P., Van de Wiele, T., Louis, P., & Kleerebezem, M. (2013). Intestinal colonization: how key microbial players become established in this dynamic process: microbial metabolic activities and the interplay between the host and microbes. *BioEssays*, 35(10), 913-923. doi:10.1002/bies.201300073
- Feliciano, R. P., Meudt, J. J., Shanmuganayagam, D., Krueger, C. G., & Reed, J. D. (2014). Ratio of “A-type” to “B-type” proanthocyanidin interflavan bonds affects extra-intestinal pathogenic Escherichia coli invasion of gut epithelial cells. *Journal of agricultural and food chemistry*, 62(18), 3919-3925.
- Fogliano, V., Corollaro, M. L., Vitaglione, P., Napolitano, A., Ferracane, R., Travaglia, F., . . . Gibson, G. (2011). In vitro bioaccessibility and gut biotransformation of polyphenols present in the water-insoluble cocoa fraction. *Mol Nutr Food Res*, 55 Suppl 1, S44-55. doi:10.1002/mnfr.201000360
- Fu, Z., Liska, D., Talan, D., & Chung, M. (2017). Cranberry reduces the risk of urinary tract infection recurrence in otherwise healthy women: a systematic review and meta-analysis. *The Journal of nutrition*, 147(12), 2282-2288.
- Geva-Zatorsky, N., Sefik, E., Kua, L., Pasman, L., Tan, T. G., Ortiz-Lopez, A., . . . Mathis, D. (2017). Mining the human gut microbiota for immunomodulatory organisms. *Cell*, 168(5), 928-943.



- González de Llano, D., Esteban-Fernández, A., Sánchez-Patán, F., Martínlvarez, P. J., Moreno-Arribas, M., & Bartolomé, B. (2015). Anti-adhesive activity of cranberry phenolic compounds and their microbial-derived metabolites against uropathogenic *Escherichia coli* in bladder epithelial cell cultures. *International Journal of Molecular Sciences*, 16(6), 12119-12130.
- González de Llano, D., Liu, H., Khoo, C., Moreno-Arribas, M. V., & Bartolomé, B. a. (2019). Some new findings regarding the antiadhesive activity of cranberry phenolic compounds and their microbial-derived metabolites against uropathogenic bacteria. *Journal of agricultural and food chemistry*, 67(8), 2166-2174.
- Hammer, Ø., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia electronica*, 4(1), 9.
- Hsia, D. S., Zhang, D. J., Beyl, R. S., Greenway, F. L., & Khoo, C. (2020). Effect of Daily Consumption of Cranberry Beverage on Insulin Sensitivity and Modification of Cardiovascular Risk Factors in Obese Adults: A Pilot Randomized Placebo-Controlled Study. *British Journal of Nutrition*, 1-27.
- Ianiro, G., Tilg, H., & Gasbarrini, A. (2016). Antibiotics as deep modulators of gut microbiota: between good and evil. *Gut*, 65(11), 1906-1915.
- Jass, J., & Reid, G. (2009). Effect of cranberry drink on bacterial adhesion in vitro and vaginal microbiota in healthy females. *The Canadian journal of urology*, 16(6), 4901-4907.
- Jung, C., & Brubaker, L. (2019). The etiology and management of recurrent urinary tract infections in postmenopausal women. *Climacteric*, 22(3), 242-249.
- Koutsos, A., & Lovegrove, J. A. (2015). Chapter 12 - An Apple a Day Keeps the Doctor Away – Inter-Relationship Between Apple Consumption, the Gut Microbiota and Cardiometabolic Disease Risk Reduction. In (pp. 173-194): Elsevier Inc.
- Koutzoumis, D. N., Vergara, M., Pino, J., Buddendorff, J., Khoshbouei, H., Mandel, R. J., & Torres, G. E. (2020). Alterations of the gut microbiota with antibiotics protects dopamine neuron loss and improve motor deficits in a pharmacological rodent model of Parkinson's disease. *Experimental Neurology*, 325, 113159.
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Applied and Environmental Microbiology*, 79(17), 5112-5120. doi:10.1128/AEM.01043-13
- Lavefve, L., Howard, L. R., & Carbonero, F. (2020). Berry polyphenols metabolism and impact on human gut microbiota and health. *Food & Function*, 11(1), 45-65.
- Leclercq, S. Y., Sullivan, M. J., Ipe, D. S., Smith, J. P., Cripps, A. W., & Ulett, G. C. (2016). Pathogenesis of *Streptococcus* urinary tract infection depends on bacterial strain and  $\beta$ -

- hemolysin/cytolysin that mediates cytotoxicity, cytokine synthesis, inflammation and virulence. *Scientific reports*, 6(1), 29000. doi:10.1038/srep29000
- Liu, H., Howell, A. B., Zhang, D. J., & Khoo, C. (2019). A randomized, double-blind, placebo-controlled pilot study to assess bacterial anti-adhesive activity in human urine following consumption of a cranberry supplement. *Food & Function*, 10(12), 7645-7652.
- Liu, H., Wang, J., He, T., Becker, S., Zhang, G., Li, D., & Ma, X. (2018). Butyrate: a double-edged sword for health? *Advances in nutrition*, 9(1), 21-29.
- Ma, B., Forney, L. J., & Ravel, J. (2012). Vaginal microbiome: rethinking health and disease. *Annual review of microbiology*, 66, 371-389.
- Maisuria, V. B., Lopez-de Los Santos, Y., Tufenkji, N., & Déziel, E. (2016). Cranberry-derived proanthocyanidins impair virulence and inhibit quorum sensing of *Pseudomonas aeruginosa*. *Scientific reports*, 6, 30169.
- Maki, K. C., Kaspar, K. L., Khoo, C., Derrig, L. H., Schild, A. L., & Gupta, K. (2016). Consumption of a cranberry juice beverage lowered the number of clinical urinary tract infection episodes in women with a recent history of urinary tract infection. *The American journal of clinical nutrition*, 103(6), 1434-1442.
- Marchesi, A., Silva, J., Wiese, B., & Nader-Macías, M. E. F. (2020). Survival of Beneficial Vaginal Lactobacilli (BVL) to Different Gastrointestinal Tract Conditions. *Current Pharmaceutical Design*.
- Mayta-Apaza, A. C., Pottgen, E., De Bodt, J., Papp, N., Marasini, D., Howard, L., . . . Carbonero, F. (2018). Impact of tart cherries polyphenols on the human gut microbiota and phenolic metabolites in vitro and in vivo. *The Journal of nutritional biochemistry*, 59, 160-172.
- Mena, P., de Llano, D. G., Brindani, N., Esteban-Fernández, A., Curti, C., Moreno-Arribas, M. V., . . . Bartolomé, B. (2017). 5-(3', 4'-Dihydroxyphenyl)- $\gamma$ -valerolactone and its sulphate conjugates, representative circulating metabolites of flavan-3-ols, exhibit anti-adhesive activity against uropathogenic *Escherichia coli* in bladder epithelial cells. *Journal of Functional Foods*, 29, 275-280.
- Mendling, W. (2016). Vaginal microbiota. In *Microbiota of the human body* (pp. 83-93): Springer.
- Morais, C. A., de Rosso, V. V., Estadella, D., & Pisani, L. P. (2016). Anthocyanins as inflammatory modulators and the role of the gut microbiota. *J Nutr Biochem*, 33, 1-7. doi:10.1016/j.jnutbio.2015.11.008
- Morgun, A., Dzutsev, A., Dong, X., Greer, R. L., Sexton, D. J., Ravel, J., . . . Shulzhenko, N. (2015). Uncovering effects of antibiotics on the host and microbiota using transkingdom gene networks. *Gut*, 64(11), 1732-1743.

- O'Keefe, S. J. D. (2016). Diet, microorganisms and their metabolites, and colon cancer. *Nature reviews Gastroenterology & hepatology*, 13(12), 691.
- O'May, C., & Tufenkji, N. (2011). The swarming motility of *Pseudomonas aeruginosa* is blocked by cranberry proanthocyanidins and other tannin-containing materials. *Applied and Environmental Microbiology*, 77(9), 3061-3067.
- O'Connor, K., Morrisette, M., Strandwitz, P., Ghiglieri, M., Caboni, M., Liu, H., . . . Lewis, K. (2019). Cranberry extracts promote growth of Bacteroidaceae and decrease abundance of Enterobacteriaceae in a human gut simulator model. *PloS one*, 14(11), e0224836.
- Pan, M., Hidalgo-Cantabrana, C., Goh, Y. J., Sanozky-Dawes, R., & Barrangou, R. (2020). Comparative analysis of *Lactobacillus gasseri* and *Lactobacillus crispatus* isolated from human urogenital and gastrointestinal tracts. *Frontiers in microbiology*, 10, 3146.
- Png, C. W., Linden, S. K., Gilshenan, K. S., Zoetendal, E. G., McSweeney, C. S., Sly, L. I., . . . Sahlgrenska, A. (2010). Mucolytic bacteria with increased prevalence in IBD mucosa augment in vitro utilization of mucin by other bacteria. *American Journal of Gastroenterology*, 105(11), 2420-2428. doi:10.1038/ajg.2010.281
- Possemiers, S., Bolca, S., Verstraete, W., & Heyerick, A. (2011). The intestinal microbiome: a separate organ inside the body with the metabolic potential to influence the bioactivity of botanicals. *Fitoterapia*, 82(1), 53-66. doi:10.1016/j.fitote.2010.07.012
- Precup, G., & Vodnar, D.-C. (2019). Gut Prevotella as a possible biomarker of diet and its eubiotic versus dysbiotic roles: a comprehensive literature review. *British Journal of Nutrition*, 122(2), 131-140. doi:10.1017/S0007114519000680
- Puopolo, K. M., Lynfield, R., Cummings, J. J., & Committee On Infectious, D. (2019). Management of infants at risk for group B streptococcal disease. *Pediatrics*, 144(2), e20191881.
- Queipo-Ortuño, M. I., Boto-Ordóñez, M., Murri, M., Gomez-Zumaquero, J. M., Clemente-Postigo, M., Estruch, R., . . . Tinahones, F. J. (2012). Influence of red wine polyphenols and ethanol on the gut microbiota ecology and biochemical biomarkers. *The American journal of clinical nutrition*, 95(6), 1323-1334. doi:10.3945/ajcn.111.027847
- Rajkumar, H., Mahmood, N., Kumar, M., Varikuti, S. R., Challa, H. R., & Myakala, S. P. (2014). Effect of probiotic (VSL#3) and omega-3 on lipid profile, insulin sensitivity, inflammatory markers, and gut colonization in overweight adults: a randomized, controlled trial. *Mediators Inflamm*, 2014, 348959. doi:10.1155/2014/348959
- Ravel, J., Gajer, P., Abdo, Z., Schneider, G. M., Sara, S. K. K., McCulle, S. L., . . . Gordon, J. I. (2011). Vaginal microbiome of reproductive-age women. *Proceedings of the National Academy of Sciences of the United States of America*, 108(Supplement 1), 4680-4687. doi:10.1073/pnas.1002611107

- Reid, G. (2018). Has knowledge of the vaginal microbiome altered approaches to health and disease? *F1000Research*, 7.
- Rivière, A., Selak, M., Lantin, D., Leroy, F., & De Vuyst, L. (2016). Bifidobacteria and Butyrate-Producing Colon Bacteria: Importance and Strategies for Their Stimulation in the Human Gut. *Frontiers in microbiology*, 7, 979. doi:10.3389/fmicb.2016.00979
- Rodríguez-Daza, M.-C., Roquim, M., Dudonné, S., Pilon, G., Levy, E., Marette, A., . . . Desjardins, Y. (2020). Berry polyphenols and fibers modulate distinct microbial metabolic functions and gut microbiota enterotype-like clustering in obese mice. *Frontiers in microbiology*, 11, 2032.
- Rodríguez-Morató, J., Matthan, N. R., Liu, J., de la Torre, R., & Chen, C. Y. O. (2018). Cranberries attenuate animal-based diet-induced changes in microbiota composition and functionality: a randomized crossover controlled feeding trial. *The Journal of nutritional biochemistry*, 62, 76-86.
- Simmering, J. E., Tang, F., Cavanaugh, J. E., Polgreen, L. A., & Polgreen, P. M. (2017). The Increase in Hospitalizations for Urinary Tract Infections and the Associated Costs in the United States, 1998-2011. *Open forum infectious diseases*, 4(1), ofw281-ofw281. doi:10.1093/ofid/ofw281
- Singh, R. K., Chang, H. W., Yan, D., Lee, K. M., Ucmak, D., Wong, K., . . . Liao, W. (2017). Influence of diet on the gut microbiome and implications for human health. *J Transl Med*, 15(1), 73. doi:10.1186/s12967-017-1175-y
- Stapleton, A. E. (2016). The Vaginal Microbiota and Urinary Tract Infection. *Microbiology spectrum*, 4(6). doi:10.1128/microbiolspec.UTI-0025-2016
- Tan, C. K., Ulett, K. B., Steele, M., Benjamin, J. W. H., & Ulett, G. C. (2012). Prognostic value of semi-quantitative bacteruria counts in the diagnosis of group B streptococcus urinary tract infection: a 4-year retrospective study in adult patients. *BMC infectious diseases*, 12(1), 273-273. doi:10.1186/1471-2334-12-273
- Ulett, K. B., William, H. B., Jr., Zhuo, F., Xiao, M., Kong, F., Gilbert, G. L., . . . Ulett, G. C. (2009). Diversity of Group B Streptococcus Serotypes Causing Urinary Tract Infection in Adults. *Journal of Clinical Microbiology*, 47(7), 2055-2060. doi:10.1128/JCM.00154-09
- Verstraelen, H., Verhelst, R., Vaneechoutte, M., & Temmerman, M. (2011). Group A streptococcal vaginitis: an unrecognized cause of vaginal symptoms in adult women. *Archives of gynecology and obstetrics*, 284(1), 95-98.
- Walker, A. W., Duncan, S. H., Louis, P., & Flint, H. J. (2014). Phylogeny, culturing, and metagenomics of the human gut microbiota. *Trends in Microbiology*, 22(5), 267-274. doi:10.1016/j.tim.2014.03.001

- White, B. A., Creedon, D. J., Nelson, K. E., & Wilson, B. A. (2011). The vaginal microbiome in health and disease. *Trends in Endocrinology & Metabolism*, 22(10), 389-393. doi:10.1016/j.tem.2011.06.001
- Witkin, S. S., & Linhares, I. M. (2017). Why do lactobacilli dominate the human vaginal microbiota? *BJOG: An International Journal of Obstetrics & Gynaecology*, 124(4), 606-611.
- Wu, G. D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y.-Y., Keilbaugh, S. A., . . . Lewis, J. D. (2011). Linking Long-Term Dietary Patterns with Gut Microbial Enterotypes. *Science*, 334(6052), 105-108. doi:10.1126/science.1208344
- Zhao, S., Liu, H., Su, Z., Khoo, C., & Gu, L. (2020). Identifying Cranberry Juice Consumers with Predictive OPLS-DA Models of Plasma Metabolome and Validation of Cranberry Juice Intake Biomarkers in a Double-Blinded, Randomized, Placebo-Controlled, Cross-Over Study. *Molecular nutrition & food research*, 64(11), 1901242.
- Zhu, A., Sunagawa, S., Mende, D. R., & Bork, P. (2015). Inter-individual differences in the gene content of human gut bacterial species. *Genome biology*, 16(1), 82.

## **CHAPTER 4:**

### **Botanical Diversity and Amount of Dietary Fruits and Vegetables Distinctly Affect the Human Gut Microbiome**

#### **Abstract**

While increasing fruit and vegetable (FV) intake is a near-universal recommendation for improved health outcomes, information on dietary FV the impact amount and diversity on health biomarkers is scarce. FV are a major dietary source of accessible gut microbiota carbohydrates and phytochemicals, however, most studies have focused on single food items or their extracted components, with few holistic studies available. Here, two separate randomized dietary interventions were used to assess the impact of low vs. adequate FV intake and low vs. high botanical diversity on gut microbiota profiles in healthy adults. We hypothesized that increasing FV would result in beneficial modulations to gut microbiota with further increases to benefit those consuming FV from diverse botanical families. The SMART-Diet study was a pilot, randomized, cross-over dietary intervention where male volunteers were randomized to start diets of low FV or adequate FV over 9 days. Stool samples were obtained at day 0, 3, 6 and 9 for each treatment period. In the SMART-Valid study, 21 individuals were provided a low FV lead-in diet for 4 days and then randomly assigned to a high FV diet with either low (11 families) or high botanical diversity (24 families) for an additional 4 days. Stool was collected at baseline and after each diet intervention. Gut microbiota was analyzed using 16S rRNA sequencing performed on an Illumina MiSeq. The Mothur pipeline was used for preliminary data analysis, followed by statistical analyses in PAST. In the SMART-Diet study, the low FV treatment resulted in minimal microbiota alterations, while a significant increase in Bacteroidetes and

decrease in Firmicutes (which peaked at day 6) was observed in the adequate FV treatment group. Intriguingly, the low group experienced a short-term increase in *Bifidobacterium* and *Lactobacillus*, and both treatments incurred significant increases in *Bacteroides* and *Akkermansia* and a decline of *Faecalibacterium*. In the SMART-Valid study, the transition from low to high FV resulted in similar trends as the SMART-Diet study. However, the high botanical treatment resulted in a more diverse gut microbiota, characterized by increased relative abundances of beneficial Firmicutes (Lachnospiraceae, *Faecalibacterium*, and *Clostridium* XIVa). Our results suggest that both the amounts of FV consumed and botanical diversity modulate the gut microbiota. Determining better FV combinations from a gut microbiota perspective, thus, appears as a possible task for future research.

**Keywords:** Fruits, Vegetables, Gut microbiota, Fibers, Polyphenols.

#### 4.1 Introduction

A balanced diet is a substantial factor in promoting human health and limiting chronic disease risks (Cena & Calder, 2020; Fischer, Pallazola, Xun, Cainzos-Achirica, & Michos, 2020; Slavin & Lloyd, 2012; World Health, 2019). Fruits and vegetables (FV) are fundamental components of a healthy diet and have been the keystone foods of dietary recommendations for decades (Dietary Guidelines Advisory, 2015; Lampe, 1999; Phe, 2016; Trichopoulou et al., 2003; Vanderlee, McCrory, & Hammond, 2015; Wallace et al., 2019; World Health, 2019). The U.S. Departments of Health and Human Services (DHHS) and Agriculture (USDA) recommended that half of a meal consists of FV (Dietary Guidelines Advisory, 2015). Generally, FV contain a diverse collection of nutrients and biologically active components that provide potential health benefits, including dietary fibers, phytochemicals, and vitamins (Sharoba,

Farrag, & Abd El-Salam, 2013; Slavin & Lloyd, 2012; Yahia, García-Solís, & Celis, 2019).

Epidemiologic studies have overwhelmingly confirmed the role of FV in the reduction of systemic inflammation and the prevention of various chronic diseases and metabolic disorders such as obesity, diabetes, high blood pressure, cardiovascular diseases, and cancers (Bertoia et al., 2015; Farvid et al., 2016; Farvid et al., 2019; Gehlich et al., 2020; Gluba-Brzózka, Franczyk, & Rysz, 2017; Jurikova et al., 2017; Lee & Park, 2017; Muraki et al., 2013; Mursu, Virtanen, Tuomainen, Nurmi, & Voutilainen, 2014; Tome-Carneiro & Visioli, 2016; Wang et al., 2014; Yokoyama et al., 2014).

FV are an excellent source of dietary fibers, which are primarily credited with imparting the health benefits associated with adequate consumption (Korcz, Kerényi, & Varga, 2018). Many types of dietary fibers support gastrointestinal and digestive health by modulation of the gut microbiota. The gut microbiota has recently emerged as an important regulator of human health, and fermentable fibers serve as a critical energy source for colonic microorganisms, which subsequently result in the production of bioactive metabolites, such as short chain fatty acids (SCFAs) (Flint, Duncan, & Louis, 2017; Holscher, 2017; Makki, Deehan, Walter, & Bäckhed, 2018). A series of meta-analyses and systematic reviews recently summarized in the Lancet concluded that dietary fiber intake is inversely associated with the risk of developing non-communicable diseases (Reynolds et al., 2019); and the strong dose-responsiveness of risk reduction suggests a causal link between fiber and disease protection. Furthermore, fiber rich FV are also high in other bioactive compounds, such as phytochemicals, which may be partially responsible for the widely acknowledged reduction in disease risks and other benefits to human health.



Phytochemicals are a chemically diverse array of secondary metabolites produced by plants. In nature, they play a role in the plant's growth and defense against competitors, pathogens, predators, or environmental factors (Bartwal, Mall, Lohani, Guru, & Arora, 2013; Rasmann & Agrawal, 2009). Phytochemicals are organized according to their chemical structure; therefore, phytochemicals in the same group typically have similar mechanisms of action in the human body (Produce for Better Health, 2015). They exert various physiologic effects in humans, including acting as anti-inflammatory and anti-oxidant agents (Hussain et al., 2016; Li et al., 2014; Williamson, 2017; H. Zhang & Tsao, 2016). Major categories of health-promoting phytochemicals include phenolic compounds, terpenoids, alkaloids, and other nitrogen-containing compounds and sulphur-containing compounds. While some of these compounds, such as polyphenols, are widely distributed among FV (Abbas et al., 2017; Abuajah, Ogbonna, & Osuji, 2015; Edwards et al., 2017; Williamson, 2017), others classes, such as alkaloids and glucosinolates, are much more restricted in occurrence. Like dietary fiber, phytochemicals can positively alter the gut microbiota composition both through provision of conjugated sugars that can be used as an energy source as well as through anti-microbial activity that can selectively inhibit some microorganisms and reduce competition within mixed microbial populations. These effects on the microbiota, along with direct anti-inflammatory and anti-oxidant activity within the intestines, can promote homeostasis within the gastrointestinal tract (Cardona, Andrés-Lacueva, Tulipani, Tinahones, & Queipo-Ortuño, 2013; Kumar Singh et al., 2019; Lavefve, Howard, & Carbonero, 2020; Ozdal et al., 2016). Furthermore, the metabolism of these compounds within the gut may also increase their bioavailability and bioactivity, suggesting that inter-individual variability in microbial composition and capacity to metabolize phytochemicals can influence their response to exposure to these compounds.

Despite near-universal recommendations regarding the intake of FV to improve health outcomes, the average vegetable intake was found to be below the recommendations of 240 g/day in approximately 88% of the countries (Kalmpourtzidou, Eilander, & Talsma, 2020). In the U.S., only one in 10 adults report adequate intake of FV (Lee-Kwan, Moore, Blanck, Harris, & Galuska, 2017). Not only are Americans falling short in their FV consumption, but the variety and botanical diversity of commonly consumed FV is also low. The most consumed whole fruits by Americans include bananas, apples, and berries, while the most consumed vegetables include white potatoes, lettuce, onions, and tomatoes (Produce for Better Health, 2015). Among, these most popular vegetables are provided similar phytochemicals. Therefore, not only is FV intake generally inadequate, but the lack of phytochemical diversity in consumed foods may be limiting potential health benefits, both in gut health and in overall disease prevention. Here, we aim to identify the impact of adequate versus inadequate FV intake on the gut microbiota as well as identify whether botanical diversity can promote microbial diversity in the gut. In this study, two separate randomized short-term dietary interventions were used to assess the impact of low vs. adequate FV intake, according to the USDA recommendations (Dietary Guidelines Advisory, 2015), and low vs. high botanical diversity on gut microbiome profiles in healthy adults. We hypothesized that increasing FV in meals would result in beneficial modulations to the gut microbiota such as increased diversity and increased relative abundance of beneficial genera, with even more marked benefits in those consuming FV from diverse botanical families.

## 4.2 Materials and Methods

### 4.2.1 Subjects

#### *4.2.1.1 SMART-Diet study*

This study was a pilot, randomized, cross-over dietary intervention and was conducted to investigate the impact of low vs. adequate FV intake on the gut microbiota ecosystem. Eleven (11) males between 18-35 years of age at Colorado State University participated in this study; however, only a full set of samples from six (6) participants were used for gut microbiota analysis, and that is what is used for the sequence data. Recruitment was conducted by word of mouth, email, and flyers. Inclusion and exclusion criteria are detailed in **Table 4. 1**. Study protocol was approved by the Colorado State University Institutional Review Board (CSU IRB), Protocol #13-4278H and all participants consented in writing before participating in the study.

#### *4.2.1.2 SMART-Valid study*

This study was a pilot, parallel-arm dietary intervention which was conducted to assess the impact of low vs. high botanical diversity on gut microbiome profiles. This study was completed by twenty-one (21) healthy volunteers, nine (9) males and twelve (12) females between 18-65 years old with BMI's between 20-29.9. Eligibility was determined using a questionnaire. Inclusion and exclusion criteria are outlined in **Table 4. 1**. All participants provided written consent, and the study protocol was approved by the CSU IRB, Protocol #17-7214H.

## 4.2.2 Study Design and sample collection

### 4.2.2.1 *SMART-Diet study*

Participants were randomized into adequate (five cups of FV daily) and low (two cups of FV daily) starting treatments at the beginning of the study. Starting diet treatments spanned nine days followed by a twelve-day wash-out period and a second nine-day treatment intervention (**Figure 4. 1**). Biological samples (blood and stool) and anthropometric measures were recorded during clinical visits. All samples were stored at -80°C prior to processing. Dietary intake information was collected using three-day food logs, which included two weekdays and one weekend day. Diet data was collected prior to the start of both diet treatments (baseline) and during each treatment period (compliance). FV were provided, but remaining foods were self-selected by participants. Participants were given menu suggestions, developed by registered dietitians at Colorado State University's Kendall Reagan Nutrition Center (KRNC), for the remainder of their intake. A sample menu for each treatment is provided in **Appendix 1**. Participants were asked to consume all of the provided foods and the remainder of their diet was *ad libitum*. All provided FV were prepared in kitchen facilities located in the KRNC. Participants were provided with new portions of FV every three days during their clinical visits. To measure compliance, diet interviews were conducted at each clinical visit and food containers and any uneaten food portions (including peels, pits, and skins) were returned.

### 4.2.2.2 *SMART-Valid study*

Participants were assigned to the two intervention groups based upon schedule availability. Those who were available during the earlier intervention schedule followed the low botanical diversity protocol and those who requested the later intervention schedule followed the

high botanical diversity protocol. Baseline diet was assessed using two 24-hour recalls using the ASA24 tool (<https://epi.grants.cancer.gov/asa24/>) the week before the start of the intervention period. Prior to starting the diet intervention, participants collected baseline stool samples, and returned them to the clinic the following day. Participants were instructed to store all stool samples in the refrigerator until they were delivered to the lab. Baseline anthropometric measures were collected as well as a fasting blood sample.

After completing the clinic visit, all participants consumed a pre-specified lead-in low FV diet for a period of four days. Following this, participants provided a second stool sample on the morning of day five. They then began one of two treatment groups: high FV intake with low biodiversity, or high FV intake with high biodiversity. They followed their intervention diets for four days. On day nine of the intervention, participants provided a third stool sample (**Figure 4. 2**). All samples were stored at -80°C prior to processing.

The treatment meals were specified by a pre-designed menu and provided varying amounts of FV depending on the respective treatment groups (**Appendix 2, 3**). The low FV diets were comprised of two to three cups of FV per day coming from six different botanical families. Both of the high FV diets contained five to nine cups of FV daily, with the high biodiversity diet containing foods from 24 different botanical families and the low diversity diet containing foods from 11 different botanical families. All foods were provided through CSU's Housing and Dining Services. Compliance with prescribed menus was determined through pre- and post-meal photos of all meals and snacks.

### **4.2.3 DNA extraction**

Bacterial DNA extraction from stool samples was conducted using commercial QIAamp DNA stool mini kits (QIAGEN, Valencia, CA) following the manufacturer's protocol with the addition of a cell lysis step to rapidly and completely disrupt cells in the samples to release DNA. The cell lysis was performed by adding 0.1 g of 0.1 mm diameter and 0.1 g of 0.5 mm diameter zirconia-silica beads (BioSpec Products, Bartlesville, OK, United States) to 2 ml tubes. Subsequently, Stool content was transferred to the 2 ml tubes containing zirconia-silica beads, and 1 ml of InhibitEX Buffer (from the QIAamp® Fast DNA Stool Mini Kit) was added to the tubes. The tubes were then subjected to a bead-beater for 60 seconds at maximum speed using a FastPrep®-24 bead-beater (MP Biomedicals, Santa Ana, CA, United States).

The NanoDrop™ 1000 Spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA, United States) was used to determine the DNA concentration. Extracted DNA was then visualized following electrophoresis in a 2% agarose gel and 1X TAE (Tris-acetate ethylenediaminetetraacetic acid) buffer (AMRESCO®, Cleveland, OH, United States).

### **4.2.4 Universal polymerase chain reaction**

16S ribosomal RNA gene was amplified by the polymerase chain reaction (PCR) (95 °C for 3 min, followed by 35 cycles at 98 °C for 30 s, 55 °C for 30 s, and 72 °C for 1 min and a final extension at 72 °C for 5 min) using primers (8F 5' - AGAGTTTGATCCTGGCTCAG - 3' and 1541R 5' - AAGGAGGTGATCCAGCCGCA - 3'). PCR reactions were performed in 25 µL mixture containing 3 µL of DNA template; 1 µL of each universal primer (8F and 1541R); 12.5 µL of GoTaq® Green Master Mix (Promega™ Corporation, Wisconsin, USA); and 7.5 µL of Nuclease-Free Water. Agarose gel electrophoresis was then accomplished to confirm the success

of the PCR by using 1  $\mu$ L of SYBR Safe DNA Gel Stain (Thermo Fisher Scientific, USA) fluorescent dye.

#### **4.2.5 Amplicon Library Preparation and Sequencing**

For 16S rRNA gene sequencing (ITag) analyses, universal primers tagged with Illumina adapters and barcodes were used for PCR from the DNA extracts (Kozich, Westcott, Baxter, Highlander, & Schloss, 2013). Briefly, the PCR dual-indexed strategy was performed in 25  $\mu$ L reaction components with 3  $\mu$ L of the DNA template; 2.5  $\mu$ L of the AccuPrime™ PCR buffer II; 0.5  $\mu$ L of each index primers; 0.1  $\mu$ L of AccuPrime™ Taq DNA Polymerase (Invitrogen, USA); and 18.4  $\mu$ L of Nuclease-Free Water following the manufacture's protocol. The PCR consisted of initial denaturation at 95 °C for 3 min, followed by 30 cycles of denaturation at 95 °C for 30 sec, primer annealing at 55 °C for 30 sec, and an extension at 72 °C for 1 min, with a final extension step at 72 °C for 5 min. Correct size amplicons were subjected to purification and normalization using Invitrogen SequalPrep kits following the manufacturer's protocol. Amplicons concentrations were measured by Agilent bioanalyzer after pooling as two different libraries.

For accurate concentration, amplicon libraries were quantified by qPCR using the same primers and PerfeCta NGS library quantification kits (Quanta Biosciences, USA) following manufacturer instructions. Libraries were then denatured and diluted to appropriate concentrations and sequenced on an Illumina MiSeq.

Resulting sequencing files were analyzed using the Mothur v.1.41.1 pipeline (Schloss et al., 2009). Operational taxonomic units (OTU) were determined and classified against reference databases (Ribosomal Database Project (RDP), SILVA,)). Basic and multivariate statistical

analyses were performed in PAST (Al Othaim, Marasini, & Carbonero, 2020; Hammer, Harper, & Ryan, 2001).

#### **4.2.6 Short-chain fatty acids analyses**

Frozen stool samples from the SMART-Valid study were extracted for SCFAs by mixing 0.75 g of frozen feces with acidified water (pH 2.5) and sonicating for 10 min. as described previously (Sheflin et al., 2015; Weir et al., 2013). Briefly, samples were analyzed on a 6890 series Gas Chromatograph with a flame ionization detector (GC\_FID; Agilent). Samples were injected at a 10:1 split ratio; the inlet was held at 22°C and the transfer line was held at 230°C. Separation of SCFAs was achieved on a 30m TG-WAX-A column (0.25-mm ID, 0.25-μm film thickness; Thermo Scientific) by using a temperature program of 100°C for 1 min, increased by 8°C/min to 180°C, held at 180°C for 1 min, increased by 200°C at 20°C/min, and held at 200°C for 5 min. Helium carrier flow was held at 1.2 mL/min. Commercial standards of acetic acid, propionic acid and butyric acid, (Sigma, St. Louis, MO, USA) were used to confirm compound identities.

#### **4.2.7 Statistical analysis**

Statistical analyses including Wilcoxon signed-rank test and Mann-Whitney U test (depending on the availability of fully paired sample sets) were performed to detect significant differences between groups (by convention, differences were considered significant when  $P < 0.05$ ). Non-Metric Multidimensional scaling (NMDS) based on count distance metrics (Bray-Curtis similarity index; ANOSIM:  $P < 0.05$ ) were considered to be significant similarities between groups using Past4 software.



## 4.3 Results and Discussion

### 4.3.1 SMART-Diet study: Longitudinal assessment of the impact of low and adequate FV intake

Using NMDS visualization of Bray-Curtis distances, short-term changes to FV intake did not significantly impact the general gut microbiota profiles (**Figure 4. 3**). These results are in line with other studies which revealed no significant impact from a vegetarian diet on the microbial community composition in healthy and obese individuals (Kim, Hwang, Park, & Bae, 2013; Losasso et al., 2018). Likewise, a study showed that short-term dietary changes could alter the gut in as little as 24 hours, but that the diet high in animal products had a much greater immediate impact on the microbiota than the plant-based diet (David et al., 2014). It is clear from the NMDS that the individuality of each subject had a stronger impact than the diet assignment; specifically, half of the subjects were responders to the diet change, while the other half were not responders (**Figure 4. 4**). This observation is also logical considering the numerous reports of metatypes with differing response to dietary phytochemicals (Cortés-Martín et al., 2020; Mayta-Apaza et al., 2018) as well as differential responses to fiber and other plant-derived oligosaccharides (Davis, Martínez, Walter, Goin, & Hutkins, 2011; G. D. Wu et al., 2011).

Both diet interventions significantly increased the gut microbiota diversity (Shannon index), but the increase was gradual with the low FV diet (**Figure 4. 5**). Improving the gut microbiota diversity is generally considered a healthy outcome, whereas dysbiosis, loss of gut microbiota, was associated with many diseases and metabolic disorders (Kriss, Hazleton, Nusbacher, Martin, & Lozupone, 2018). A study reported that the fiber intake increased the gut

microbiota diversity and richness in overweight pregnant women (Röytiö, Mokkala, Vahlberg, & Laitinen, 2017).

When comparing phyla-level changes across the different FV intake periods, both treatments showed similar trends. The low FV diet had more heterogeneity, probably because of responder/non-responder effects. Generally, both diets showed reductions in the Firmicutes/Bacteroidetes ratio, although the maximum change from baseline occurred on different days (peak at day 6 for adequate; peaks at day 3 and 9 for low). During the low FV diet, the Firmicutes relative abundance was significantly decreased from baseline to day 9, although the Bacteroidetes were not significantly changed. In comparison, from baseline to day 9 on the adequate FV diet, there was a decrease in Firmicutes which became apparent by day 6, and an increase in Bacteroidetes, which appeared by day 3 of the diet intervention (**Figure 4. 6**).

Alteration of the Firmicutes/Bacteroidetes ratio, associated with adequate FV intake, may indicate a beneficial change to the gut microbiota. Many animal and human studies have demonstrated that the Firmicutes/Bacteroidetes ratio is increased in the microbiota of obese individuals compared to lean individuals, making it a potential indicator/biomarker for obesity (Barlow, Yu, & Mathur, 2015; Kim et al., 2013; Koliada et al., 2017; Xu, Li, Zhang, & Zhang, 2012; Zou et al., 2020). However, other studies have not observed this correlation with obesity (Hu et al., 2015; Magne et al., 2020; Tims et al., 2013; Walters, Xu, & Knight, 2014). A recent review suggests that differences in sample processing, participant demographics, and lack of control in other variables that influence the microbiota may account for some of these discrepancies in the literature, but that it is currently difficult to interpret the Firmicutes/Bacteroidetes ratio in light of human health and obesity status (Magne et al., 2020).

Genus-level microbial dynamics were mostly mirrored in both diet interventions. Among Bacteroidetes, the *Bacteroides*, *Prevotella*, *Alistipes*, and *Parabacteroides* genera were increased by both diet interventions compared to baseline, whereas *Faecalibacterium*, *Blautia*, *Dorea*, and Lachnospiraceae\_unclarified, which belong to Firmicutes phyla, were decreased (**Figure 4. 7, 4. 8**). Both adequate and low FV diets induced the relative abundance of butyrate producing bacteria *Oscillibacter*, *Clostridium\_IV*, and *Clostridium\_XIVa* (Gophna, Konikoff, & Nielsen, 2017; Moens & De Vuyst, 2017). A significant increase in *Akkermansia* relative abundance with a slight decrease in *Bifidobacterium* relative abundance was observed after the adequate FV diet (**Figure 4. 9, 4. 10**). The gut commensal *Akkermansia muciniphila* is a potential probiotic bacterium that is associated with systemic effects on host metabolism and immune response (Ansaldi et al., 2019; T. Zhang, Li, Cheng, Buch, & Zhang, 2019). It promises to be a therapeutic target in microbiota-related diseases, such as colitis, metabolic syndrome, immune diseases and cancer (T. Zhang et al., 2019).

#### **4.3.2 SMART-Valid: Impact of botanical diversity of a high FV diet**

In line with SMART-Diet study, we observed that the strong gut microbiota individuality prevented the detection of significantly different clusters by NMDS and ANOSIM (**Figure 4. 11**).

Both diets increased taxonomic diversity, but only significantly in the case of the high botanical diversity (**Figure 4. 12**). It can be hypothesized that this is a reflection of the diversity of glycans and phytochemicals leading to the stimulation of rare taxa with very specific metabolic abilities. The gut microbiota diversity (Shannon index) increased slightly after both

botanical diversity diets but was more marked with high botanical diversity (**Figure 4. 12**). This observation was not significant possibly because of the short dietary intervention period.

Regardless of botanical diversity, similar trends were observed at the phylum level. The Firmicutes/Bacteroidetes ratio was induced by increasing the relative abundance of Firmicutes and decreasing the relative abundance of Bacteroidetes. The Proteobacteria level was reduced in both intervention but was only significant in the low botanical diversity diet. It is commonly considered that higher abundance of Proteobacteria is a microbial signature of disease (Rizzatti, Lopetuso, Gibiino, Binda, & Gasbarrini, 2017). Similar response of Actinobacteria was observed in both groups. The low FV diet increased the abundance of Actinobacteria; however, the trends dropped with the high FV diet (**Figure 4. 13**).

Surprisingly, the low botanical diversity diet led to a decrease of the relative abundance of *Prevotella* and *Butyricimonas*, whereas the relative abundance of *Alistipes* was increased (**Figure 4. 14**). It has been reported that *Butyricimonas*, a butyrate producing bacteria, plays a potential role in maintaining the gut function and enhancing the intestinal barrier integrity (Tomsett et al., 2020; E. Wu, Jingzhu, Lingpeng, & Yaqin, 2020). The reduction of the abundance of *Butyricimonas* was found to be linked with the incidence of irritable bowel syndrome (IBS) (E. Wu et al., 2020; Yang et al., 2017). It has been found that the higher-fiber diet increased the relative abundance of *Butyricimonas* and *Prevotella* (de Moraes et al., 2017; Tomsett et al., 2020).

The low botanical diversity diet, however, led to a significant increase in the relative abundance of the butyrate producers *Faecalibacterium* and Lachnospiraceae\_unclarified as well as a suggestive increase in the *Clostridium\_XIVa* level. The results also showed a significant

decrease in the relative abundance of some other genera belonging to Firmicutes phyla including *Blautia*, *Ruminococcus*, *Dorea*, and *Clostridium\_XI* with a suggestive decrease in the *Clostridium\_IV* abundance. Similar trends were observed for the high botanical diversity diet but they were not significant (**Figure 4. 15**). Regardless of the botanical diversity, high amounts of FV interestingly limited the increase of commensal bacteria *Bifidobacterium* and *Akkermansia* (**Figure 4. 16**). The possible explanation is that these genera were outcompeted with higher abundance of other bacteria in the presence of higher amounts of FV in the diets. The low botanical diversity diet significantly decreased the relative abundance of the family Enterobacteriaceae (**Figure 4. 16**). The decrease of Enterobacteriaceae abundance was not significant with the high botanical diversity diet, but the abundance was very low in the baseline. Enterobacteriaceae family includes opportunistic pathogens that are known to cause morbidity and mortality among humans (Sassone-Corsi et al., 2016). It was reported that the relative abundance of Enterobacteriaceae was significantly low in vegan individuals (Zimmer et al., 2012).

A general limitation of DNA-based microbiota analysis is that any trends observed do not necessarily reflect the fermentation metabolic rates (Fricker, Podlesny, & Fricke, 2019; Wurm et al., 2018). To better assess them, we measured the SCFAs in stool samples from subjects of this study (**Figure 4. 17**). Remarkably, while only slight microbiota changes were observed, SCFAs levels were strikingly different between the low and high botanical diversity group. SCFAs intriguingly trended slightly down in response to low botanical diversity. The high botanical diversity resulted in significantly higher SCFA levels. It can be hypothesized that effective fermentative metabolism improvements require a higher botanical diversity in concert with a

higher amount of dietary FV. The composition of diet has rarely been the focus in such studies, and our data demonstrate it should be.


On the other hand, acetate, propionate and particularly butyrate were each significantly increased by the high botanical diversity diet. This outcome clearly indicates that botanical diversity is likely to be beneficial in stimulating the gut microbiota and the fermentative metabolism. All SCFAs have been associated with beneficial health properties (Chambers, Preston, Frost, & Morrison, 2018; De Filippis et al., 2016; Gill, Van Zelm, Muir, & Gibson, 2018). Models derived from different animal models can explain this observation, with cross-feeding and intra-genus diversity potentially at play (Ríos-Covián et al., 2016; Solden et al., 2018).

#### **4.4 Conclusion**


General trends were similar in terms of diversity indices which indicate the potential impact of FV consumption in enhancing the gut microbiota ecosystem for the participants regardless of the amount. Intriguingly, at phylum level, changes were almost opposite between the two studies as the SMART-Diet study showed a reduction of the Firmicutes/Bacteroidetes ratio whereas the SMART-Valid study resulted in an increase in the Firmicutes/Bacteroidetes ratio. This might be due to differences in the type and the amount of meals provided to the participants in each study.

**Table 4. 1:** Inclusion/Exclusion criteria for the SMART-Diet and SMART-Valid Studies.

	SMART-Diet	SMART-Valid
<b>Inclusion Criteria</b>	<ul style="list-style-type: none"> <li>• Male</li> <li>• 18-40 years old</li> <li>• BMI between 20-29.9</li> <li>• Non-smoker</li> </ul>	<ul style="list-style-type: none"> <li>• 18-65 years old</li> <li>• BMI between 20-29.9</li> <li>• Non-smoker</li> </ul>
<b>Exclusion Criteria</b>	<ul style="list-style-type: none"> <li>• Female</li> <li>• Outside of the age and BMI requirements</li> <li>• Smokers</li> <li>• Dietary restrictions and/or food allergies</li> <li>• Metabolic or intestinal disease diagnosis</li> <li>• Use of antibiotics, probiotics, dietary supplements and/or prescription drugs in previous 2 months</li> <li>• Inability to abstain from recreational drugs and limit alcohol to 1 beverage/day.</li> </ul>	<ul style="list-style-type: none"> <li>• Outside of the age and BMI requirements</li> <li>• Smokers</li> <li>• Strong food aversions</li> <li>• Dietary restrictions and/or food allergies</li> <li>• Metabolic or intestinal disease diagnosis</li> <li>• Use of antibiotics, probiotics, dietary supplements and/or prescription drugs in previous 2 months</li> <li>• Inability to abstain from recreational drugs and limit alcohol, coffee, and tea to 1 beverage total/day.</li> </ul>





**Figure 4. 1:** SMART-Diet study design and intervention.




**Figure 4. 2:** SMART-Valid study design and intervention.






**Figure 4. 3:** Impact on general microbiota profiles of the low and adequate FV treatment interventions at day 3, 6, and 9 compared to subjects' baseline. Gut microbiota data are shown on non-metric multidirectional scaling (NMDS) (Bray-Curtis similarity index; ANOSIM: ( $P < 0.05$ )).




**Figure 4. 4:** Impact on individuals' gut microbiota profiles (NMDS) of the low and high FV botanical biodiversity diet interventions. Gut microbiota data are shown on non-metric multidirectional scaling (NMDS) (Bray-Curtis similarity index; ANOSIM: ( $P < 0.05$ )).

### Low FV diet




### Adequate FV diet




**Figure 4. 5:** Impact on diversity indices (Number of taxa, Shannon index) of the low and adequate FV treatment interventions at day 3, 6, and 9 compared to subjects' baseline. Values are expressed as mean  $\pm$  standard error (*SE*). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.

### Low FV diet




### Adequate FV diet




**Figure 4. 6:** Impact of low FV and adequate FV diets at the phylum level for each time point. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.


### Low FV diet




### Adequate FV diet



**Figure 4. 7:** Impact of low FV and adequate FV diets on the composition of genera that belong to the Bacteroidetes phyla. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.




**Figure 4. 8:** Impact of low FV and adequate FV diets on the composition of genera that belong to the Firmicutes phyla. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.





**Figure 4. 9:** Impact of low FV and adequate FV diets on the composition of genera that belong to the Firmicutes phyla. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.






**Figure 4. 10:** Impact of low FV and adequate FV diets on the composition of *Bifidobacterium*, *Akkermansia*, and Enterobacteriaceae. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.






**Figure 4. 11:** Impact on general microbiota profiles (NMS) of the low and high FV botanical biodiversity diet interventions. Gut microbiota data are shown on non-metric multidirectional scaling (NMS) (Bray-Curtis similarity index; ANOSIM: ( $P < 0.05$ )).


### Low botanical diversity




### High botanical diversity



**Figure 4. 12:** Impact on diversity indices (Number of taxa, Shannon index) of the low and high FV botanical diversity diet interventions. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.




**Figure 4. 13:** Impact of the low and high FV botanical diversity consumption at the phylum level. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.




**Figure 4. 14:** Impact of the low and high FV botanical diversity consumption of the genera belong to the Bacteroidetes phyla. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.


### Low botanical diversity



### High botanical diversity



**Figure 4. 15:** Impact of the low and high FV botanical diversity consumption of the genera belong to the Firmicutes phyla. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.



### Low botanical diversity




Figure 4. 15 (Cont.)

### High botanical diversity






Figure 4. 15 (Cont.)




**Figure 4. 16:** Impact of the low and high FV botanical diversity consumption of *Bifidobacterium*, *Akkermansia*, and *Enterobacteriaceae* Unclassified. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.



### Low botanical diversity



### High botanical diversity



**Figure 4. 17:** Impact of low and high FV botanical diversity consumption on SCFAs levels. Values are expressed as mean  $\pm$  standard error (SE). Different letters indicate significant difference ( $p < 0.05$ ), whereas N.S. indicates no significant difference.

## References

- Abbas, M., Saeed, F., Anjum, F. M., Afzaal, M., Tufail, T., Bashir, M. S., . . . Suleria, H. A. R. (2017). Natural polyphenols: An overview. *International Journal of Food Properties*, 20(8), 1689-1699.
- Abuajah, C. I., Ogbonna, A. C., & Osuji, C. M. (2015). Functional components and medicinal properties of food: a review. *Journal of food science and technology*, 52(5), 2522-2529.
- Al Othaim, A., Marasini, D., & Carbonero, F. (2020). Impact of increasing concentration of tart and sweet cherries juices concentrates on healthy mice gut microbiota. *Food Frontiers*, 1(3), 224-233. doi:10.1002/fft2.46
- Ansaldo, E., Slayden, L. C., Ching, K. L., Koch, M. A., Wolf, N. K., Plichta, D. R., . . . Moon, J. J. (2019). Akkermansia muciniphila induces intestinal adaptive immune responses during homeostasis. *Science*, 364(6446), 1179-1184.
- Barlow, G. M., Yu, A., & Mathur, R. (2015). Role of the gut microbiome in obesity and diabetes mellitus. *Nutrition in clinical practice*, 30(6), 787-797.
- Bartwal, A., Mall, R., Lohani, P., Guru, S. K., & Arora, S. (2013). Role of secondary metabolites and brassinosteroids in plant defense against environmental stresses. *Journal of plant growth regulation*, 32(1), 216-232.
- Bertoia, M. L., Mukamal, K. J., Cahill, L. E., Hou, T., Ludwig, D. S., Mozaffarian, D., . . . Rimm, E. B. (2015). Changes in intake of fruits and vegetables and weight change in United States men and women followed for up to 24 years: analysis from three prospective cohort studies. *PLoS Med*, 12(9), e1001878.
- Cardona, F., Andrés-Lacueva, C., Tulipani, S., Tinahones, F. J., & Queipo-Ortuño, M. I. (2013). Benefits of polyphenols on gut microbiota and implications in human health. *The Journal of nutritional biochemistry*, 24(8), 1415-1422. doi:10.1016/j.jnutbio.2013.05.001
- Cena, H., & Calder, P. C. (2020). Defining a healthy diet: evidence for the role of contemporary dietary patterns in health and disease. *Nutrients*, 12(2), 334.
- Chambers, E. S., Preston, T., Frost, G., & Morrison, D. J. (2018). Role of gut microbiota-generated short-chain fatty acids in metabolic and cardiovascular health. *Current nutrition reports*, 7(4), 198-206.
- Cortés-Martín, A., García-Villalba, R., García-Mantrana, I., Rodríguez-Varela, A., Romo-Vaquero, M., Collado, M. C., . . . Selma, M. V. (2020). Urolithins in Human Breast Milk after Walnut Intake and Kinetics of Gordonibacter Colonization in Newly Born: The Role of Mothers' Urolithin Metabotypes. *Journal of agricultural and food chemistry*, 68(45), 12606-12616.

- David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., . . . Turnbaugh, P. J. (2014). Diet rapidly and reproducibly alters the human gut microbiome. *Nature*, 505(7484), 559-563. doi:10.1038/nature12820
- Davis, L. M. G., Martínez, I., Walter, J., Goin, C., & Hutkins, R. W. (2011). Barcoded pyrosequencing reveals that consumption of galactooligosaccharides results in a highly specific bifidogenic response in humans. *PloS one*, 6(9), e25200.
- De Filippis, F., Pellegrini, N., Vannini, L., Jeffery, I. B., La Storia, A., Laghi, L., . . . Ercolini, D. (2016). High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut*, 65(11), 1812-1821. doi:10.1136/gutjnl-2015-309957
- de Moraes, A. C. F., Fernandes, G. R., da Silva, I. T., Almeida-Pititto, B., Gomes, E. P., Pereira, A. d. C., & Ferreira, S. R. G. (2017). Enterotype may drive the dietary-associated cardiometabolic risk factors. *Frontiers in cellular and infection microbiology*, 7, 47.
- Dietary Guidelines Advisory, C. (2015). *Dietary guidelines for Americans 2015-2020*: Government Printing Office.
- Edwards, C. A., Havlik, J., Cong, W., Mullen, W., Preston, T., Morrison, D. J., & Combet, E. (2017). Polyphenols and health: Interactions between fibre, plant polyphenols and the gut microbiota. *Nutrition bulletin*, 42(4), 356-360.
- Farvid, M. S., Chen, W. Y., Michels, K. B., Cho, E., Willett, W. C., & Eliassen, A. H. (2016). Fruit and vegetable consumption in adolescence and early adulthood and risk of breast cancer: population based cohort study. *BMJ*, 353.
- Farvid, M. S., Chen, W. Y., Rosner, B. A., Tamimi, R. M., Willett, W. C., & Eliassen, A. H. (2019). Fruit and vegetable consumption and breast cancer incidence: Repeated measures over 30 years of follow-up. *International journal of cancer*, 144(7), 1496-1510.
- Fischer, N. M., Pallazola, V. A., Xun, H., Cainzos-Achirica, M., & Michos, E. D. (2020). The evolution of the heart-healthy diet for vascular health: A walk through time. *Vascular Medicine*, 25(2), 184-193.
- Flint, H. J., Duncan, S. H., & Louis, P. (2017). The impact of nutrition on intestinal bacterial communities. *Current opinion in microbiology*, 38, 59-65.
- Fricke, A. M., Podlesny, D., & Fricke, W. F. (2019). What is new and relevant for sequencing-based microbiome research? A mini-review. *Journal of advanced research*, 19, 105-112.
- Gehlich, K. H., Beller, J., Lange-Asschenfeldt, B., Köcher, W., Meinke, M. C., & Lademann, J. (2020). Consumption of fruits and vegetables: Improved physical health, mental health, physical functioning and cognitive health in older adults from 11 European countries. *Aging & Mental Health*, 24(4), 634-641.

- Gill, P. A., Van Zelm, M. C., Muir, J. G., & Gibson, P. R. (2018). Short chain fatty acids as potential therapeutic agents in human gastrointestinal and inflammatory disorders. *Alimentary pharmacology & therapeutics*, 48(1), 15-34.
- Gluba-Brzózka, A., Franczyk, B., & Rysz, J. (2017). Vegetarian diet in chronic kidney disease—A friend or foe. *Nutrients*, 9(4), 374.
- Gophna, U., Konikoff, T., & Nielsen, H. B. (2017). Oscillospira and related bacteria—From metagenomic species to metabolic features. *Environmental microbiology*, 19(3), 835-841.
- Hammer, Ø., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia electronica*, 4(1), 9.
- Holscher, H. D. (2017). Dietary fiber and prebiotics and the gastrointestinal microbiota. *Gut microbes*, 8(2), 172-184.
- Hu, H.-J., Park, S.-G., Jang, H. B., Choi, M.-G., Park, K.-H., Kang, J. H., . . . Cho, S.-H. (2015). Obesity alters the microbial community profile in Korean adolescents. *PloS one*, 10(7).
- Hussain, T., Tan, B., Yin, Y., Blachier, F., Myrlene, C. B. T., & Rahu, N. (2016). Oxidative Stress and Inflammation: What Polyphenols Can Do for Us? *Oxidative medicine and cellular longevity*, 2016. doi:10.1155/2016/7432797
- Jurikova, T., Mlcek, J., Skrovankova, S., Sumczynski, D., Sochor, J., Hlavacova, I., . . . Orsavova, J. (2017). Fruits of black chokeberry *Aronia melanocarpa* in the prevention of chronic diseases. *Molecules*, 22(6), 944.
- Kalmpourtzidou, A., Eilander, A., & Talsma, E. F. (2020). Global vegetable intake and supply compared to recommendations: a systematic review. *Nutrients*, 12(6), 1558.
- Kim, M. S., Hwang, S. S., Park, E. J., & Bae, J. W. (2013). Strict vegetarian diet improves the risk factors associated with metabolic diseases by modulating gut microbiota and reducing intestinal inflammation. *Environmental microbiology reports*, 5(5), 765-775.
- Koliada, A., Syzenko, G., Moseiko, V., Budovska, L., Puchkov, K., Perederiy, V., . . . Tkach, S. (2017). Association between body mass index and Firmicutes/Bacteroidetes ratio in an adult Ukrainian population. *BMC microbiology*, 17(1), 120.
- Korcz, E., Kerényi, Z., & Varga, L. (2018). Dietary fibers, prebiotics, and exopolysaccharides produced by lactic acid bacteria: potential health benefits with special regard to cholesterol-lowering effects. *Food & Function*, 9(6), 3057-3068.
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Applied and Environmental Microbiology*, 79(17), 5112-5120. doi:10.1128/AEM.01043-13

- Kriss, M., Hazleton, K. Z., Nusbacher, N. M., Martin, C. G., & Lozupone, C. A. (2018). Low diversity gut microbiota dysbiosis: drivers, functional implications and recovery. *Current opinion in microbiology*, 44, 34-40.
- Kumar Singh, A., Cabral, C., Kumar, R., Ganguly, R., Kumar Rana, H., Gupta, A., . . . Pandey, A. K. (2019). Beneficial effects of dietary polyphenols on gut microbiota and strategies to improve delivery efficiency. *Nutrients*, 11(9), 2216.
- Lampe, J. W. (1999). Health effects of vegetables and fruit: assessing mechanisms of action in human experimental studies. *Am J Clin Nutr*, 70(3 Suppl), 475s-490s.  
doi:10.1093/ajcn/70.3.475s
- Lavefve, L., Howard, L. R., & Carbonero, F. (2020). Berry polyphenols metabolism and impact on human gut microbiota and health. *Food & Function*, 11(1), 45-65.
- Lee-Kwan, S. H., Moore, L. V., Blanck, H. M., Harris, D. M., & Galuska, D. (2017). Disparities in state-specific adult fruit and vegetable consumption—United States, 2015. *MMWR. Morbidity and mortality weekly report*, 66(45), 1241.
- Lee, Y., & Park, K. (2017). Adherence to a vegetarian diet and diabetes risk: a systematic review and meta-analysis of observational studies. *Nutrients*, 9(6), 603.
- Li, A.-N., Li, S., Zhang, Y.-J., Xu, X.-R., Chen, Y.-M., & Li, H.-B. (2014). Resources and biological activities of natural polyphenols. *Nutrients*, 6(12), 6020-6047.
- Losasso, C., Eckert, E. M., Mastroilli, E., Villiger, J., Mancin, M., Patuzzi, I., . . . Pernthaler, J. (2018). Assessing the influence of vegan, vegetarian and omnivore oriented Westernized dietary styles on human gut microbiota: a cross sectional study. *Frontiers in microbiology*, 9, 317.
- Magne, F., Gotteland, M., Gauthier, L., Zazueta, A., Pessoa, S., Navarrete, P., & Balamurugan, R. (2020). The Firmicutes/Bacteroidetes Ratio: A Relevant Marker of Gut Dysbiosis in Obese Patients? *Nutrients*, 12(5), 1474.
- Makki, K., Deehan, E. C., Walter, J., & Bäckhed, F. (2018). The impact of dietary fiber on gut microbiota in host health and disease. *Cell host & microbe*, 23(6), 705-715.
- Mayta-Apaza, A. C., Pottgen, E., De Bodt, J., Papp, N., Marasini, D., Howard, L., . . . Carbonero, F. (2018). Impact of tart cherries polyphenols on the human gut microbiota and phenolic metabolites in vitro and in vivo. *The Journal of nutritional biochemistry*, 59, 160-172.
- Moens, F., & De Vuyst, L. (2017). Inulin-type fructan degradation capacity of Clostridium cluster IV and XIVa butyrate-producing colon bacteria and their associated metabolic outcomes. *Beneficial microbes*, 8(3), 473-490.

- Muraki, I., Imamura, F., Manson, J. E., Hu, F. B., Willett, W. C., van Dam, R. M., & Sun, Q. (2013). Fruit consumption and risk of type 2 diabetes: results from three prospective longitudinal cohort studies. *BMJ*, 347, f5001.
- Mursu, J., Virtanen, J. K., Tuomainen, T.-P., Nurmi, T., & Voutilainen, S. (2014). Intake of fruit, berries, and vegetables and risk of type 2 diabetes in Finnish men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *The American journal of clinical nutrition*, 99(2), 328-333.
- Ozdam, T., Sela, D. A., Xiao, J., Boyacioglu, D., Chen, F., & Capanoglu, E. (2016). The reciprocal interactions between polyphenols and gut microbiota and effects on bioaccessibility. *Nutrients*, 8(2), 78.
- Phe. (2016). Government Dietary Recommendations: Government Recommendations for Energy and Nutrients for Males and Females Aged 1-18 Years and 19+ Years. In: PHE London.
- Produce for Better Health, F. (2015). State of the Plate, 2015 Study on America's Consumption of Fruit and Vegetables.
- Rasmann, S., & Agrawal, A. A. (2009). Plant defense against herbivory: progress in identifying synergism, redundancy, and antagonism between resistance traits. *Current opinion in plant biology*, 12(4), 473-478.
- Reynolds, A., Mann, J., Cummings, J., Winter, N., Mete, E., & Te Morenga, L. (2019). Carbohydrate quality and human health: a series of systematic reviews and meta-analyses. *The Lancet*, 393(10170), 434-445.
- Ríos-Covián, D., Ruas-Madiedo, P., Margolles, A., Gueimonde, M., De Los Reyes-gavilán, C. G., & Salazar, N. (2016). Intestinal short chain fatty acids and their link with diet and human health. *Frontiers in microbiology*, 7, 185.
- Rizzatti, G., Lopetuso, L. R., Gibiino, G., Binda, C., & Gasbarrini, A. (2017). Proteobacteria: a common factor in human diseases. *BioMed research international*, 2017.
- Röytiö, H., Morkkala, K., Vahlberg, T., & Laitinen, K. (2017). Dietary intake of fat and fibre according to reference values relates to higher gut microbiota richness in overweight pregnant women. *British Journal of Nutrition*, 118(5), 343-352.
- Sassone-Corsi, M., Nuccio, S.-P., Liu, H., Hernandez, D., Vu, C. T., Takahashi, A. A., . . . Raffatellu, M. (2016). Microcins mediate competition among Enterobacteriaceae in the inflamed gut. *Nature (London)*, 540(7632), 280-283. doi:10.1038/nature20557
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., . . . Robinson, C. J. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology*, 75(23), 7537-7541.

- Sharoba, A. M., Farrag, M. A., & Abd El-Salam, A. M. (2013). Utilization of some fruits and vegetables waste as a source of dietary fiber and its effect on the cake making and its quality attributes. *Journal of Agroalimentary Processes and Technologies*, 19(4), 429-444.
- Sheflin, A. M., Borresen, E. C., Wdowik, M. J., Rao, S., Brown, R. J., Heuberger, A. L., . . . Ryan, E. P. (2015). Pilot dietary intervention with heat-stabilized rice bran modulates stool microbiota and metabolites in healthy adults. *Nutrients*, 7(2), 1282-1300.
- Slavin, J. L., & Lloyd, B. (2012). Health benefits of fruits and vegetables. *Advances in nutrition*, 3(4), 506-516.
- Solden, L. M., Naas, A. E., Roux, S., Daly, R. A., Collins, W. B., Nicora, C. D., . . . Jørgensen, B. (2018). Interspecies cross-feeding orchestrates carbon degradation in the rumen ecosystem. *Nature microbiology*, 3(11), 1274-1284.
- Tims, S., Derom, C., Jonkers, D. M., Vlietinck, R., Saris, W. H., Kleerebezem, M., . . . Zoetendal, E. G. (2013). Microbiota conservation and BMI signatures in adult monozygotic twins. *The ISME journal*, 7(4), 707-717.
- Tome-Carneiro, J., & Visioli, F. (2016). Polyphenol-based nutraceuticals for the prevention and treatment of cardiovascular disease: Review of human evidence. *Phytomedicine*, 23(11), 1145-1174.
- Tomsett, K. I., Barrett, H. L., Dekker, E. E., Callaway, L. K., McIntyre, D. H., & Dekker Nitert, M. (2020). Dietary Fiber Intake Alters Gut Microbiota Composition but Does Not Improve Gut Wall Barrier Function in Women with Future Hypertensive Disorders of Pregnancy. *Nutrients*, 12(12), 3862.
- Trichopoulou, A., Naska, A., Antoniou, A., Friel, S., Trygg, K., & Turrini, A. (2003). Vegetable and fruit: the evidence in their favour and the public health perspective. *Int J Vitam Nutr Res*, 73(2), 63-69. doi:10.1024/0300-9831.73.2.63
- Vanderlee, L., McCrory, C., & Hammond, D. (2015). Awareness and knowledge of recommendations from Canada's Food Guide. *Canadian Journal of Dietetic Practice and Research*, 76(3), 146-149.
- Wallace, T. C., Bailey, R. L., Blumberg, J. B., Burton-Freeman, B., Chen, C. O., Crowe-White, K. M., . . . Lewis, R. (2019). Fruits, vegetables, and health: A comprehensive narrative, umbrella review of the science and recommendations for enhanced public policy to improve intake. *Critical Reviews in Food Science and Nutrition*, 1-38.
- Walters, W. A., Xu, Z., & Knight, R. (2014). Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS letters*, 588(22), 4223-4233.
- Wang, X., Ouyang, Y., Liu, J., Zhu, M., Zhao, G., Bao, W., & Hu, F. B. (2014). Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer:

- systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ*, 349, g4490.
- Weir, T. L., Manter, D. K., Sheflin, A. M., Barnett, B. A., Heuberger, A. L., & Ryan, E. P. (2013). Stool microbiome and metabolome differences between colorectal cancer patients and healthy adults. *PloS one*, 8(8), e70803.
- Williamson, G. (2017). The role of polyphenols in modern nutrition. *Nutrition bulletin*, 42(3), 226-235.
- World Health, O. (2019). *Healthy diet*. Retrieved from [https://apps.who.int/iris/bitstream/handle/10665/325828/EMROPUB\\_2019\\_en\\_23536.pdf](https://apps.who.int/iris/bitstream/handle/10665/325828/EMROPUB_2019_en_23536.pdf)
- Wu, E., Jingzhu, S., Lingpeng, P., & Yaqin, L. (2020). Comparison of the Gut Microbiota Disturbance in Rat Models of Irritable Bowel Syndrome Induced by Maternal Separation and Multiple Early-life Adversity.
- Wu, G. D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y.-Y., Keilbaugh, S. A., . . . Lewis, J. D. (2011). Linking Long-Term Dietary Patterns with Gut Microbial Enterotypes. *Science*, 334(6052), 105-108. doi:10.1126/science.1208344
- Wurm, P., Dörner, E., Kremer, C., Spranger, J., Maddox, C., Halwachs, B., . . . Högenauer, C. (2018). Qualitative and quantitative DNA-and RNA-based analysis of the bacterial stomach microbiota in humans, mice, and gerbils. *Msystems*, 3(6).
- Xu, P., Li, M., Zhang, J., & Zhang, T. (2012). Correlation of intestinal microbiota with overweight and obesity in Kazakh school children. *BMC microbiology*, 12(1), 283.
- Yahia, E. M., García-Solís, P., & Celis, M. E. M. (2019). Contribution of fruits and vegetables to human nutrition and health. In *Postharvest physiology and biochemistry of fruits and vegetables* (pp. 19-45): Elsevier.
- Yang, C., Qu, Y., Fujita, Y., Ren, Q., Ma, M., Dong, C., & Hashimoto, K. (2017). Possible role of the gut microbiota–brain axis in the antidepressant effects of (R)-ketamine in a social defeat stress model. *Translational psychiatry*, 7(12), 1-11.
- Yokoyama, Y., Nishimura, K., Barnard, N. D., Takegami, M., Watanabe, M., Sekikawa, A., . . . Miyamoto, Y. (2014). Vegetarian diets and blood pressure: a meta-analysis. *JAMA internal medicine*, 174(4), 577-587.
- Zhang, H., & Tsao, R. (2016). Dietary polyphenols, oxidative stress and antioxidant and anti-inflammatory effects. *Current Opinion in Food Science*, 8, 33-42.
- Zhang, T., Li, Q., Cheng, L., Buch, H., & Zhang, F. (2019). Akkermansia muciniphila is a promising probiotic. *Microbial biotechnology*, 12(6), 1109-1125.



- Zimmer, J., Lange, B., Frick, J.-S., Sauer, H., Zimmermann, K., Schwartz, A., . . . Enck, P. (2012). A vegan or vegetarian diet substantially alters the human colonic faecal microbiota. *European Journal of Clinical Nutrition*, 66(1), 53-60.
- Zou, Y., Ju, X., Chen, W., Yuan, J., Wang, Z., Aluko, R. E., & He, R. (2020). Rice bran attenuated obesity via alleviating dyslipidemia, browning of white adipocytes and modulating gut microbiota in high-fat diet-induced obese mice. *Food & Function*, 11(3), 2406-2417.

## GENERAL CONCLUSION

The relation between gut microbiota and human health has been increasingly recognized in the last few decades. Gut microbiota plays critical roles in the overall health of the host by, for instance, contributing to metabolism, maintaining homeostasis, providing energy, and educating the immune system. Fruits and vegetables are rich in dietary fibers and polyphenols which aid in the modifying and reshaping of the gut microbiota ecosystem. Most of the microbiological and nutritional studies have traditionally focused on investigating the impact of extracted compounds on gut microbiota profiles instead of whole foods. Therefore, this work brings attention to the possible impact of whole foods — including cherry juices, cranberries juice, and the amounts and botanical diversity of fruit and vegetable diets — on the gut microbiota composition.

The impact of cherry juices in gut microbiota in mice was notably influenced by the variation in the juices concentrations. Increasing cherry juices concentration was positively correlated with some considerably beneficial bacteria, including *Akkermansia* and *Barnesiella*, whereas the opposite response was observed for the *Bacteroides* population.

Cranberry juice consumption impacted the gut and vaginal microbiota in postmenopausal women. Cranberry juice intake resulted in an increase of the relative abundance of *Bifidobacterium*, a well-known probiotic bacterium. The clear beneficial impact of cranberry juice was the improvement of dysbiosed vaginal microbiota by reducing the relative abundance of *Streptococcus*.

Fruit and vegetable consumption enhanced the gut microbiota ecosystem for the participants regardless of the amount. Intriguingly, the effects of fruit and vegetable consumption at the phylum level were almost opposite between the two studies, the SMART-Diet and SMART-Valid. The SMART-Diet study showed a reduction of the Firmicutes/Bacteroidetes

ratio whereas the SMART-Valid study resulted in an increase in the Firmicutes/Bacteroidetes ratio. Remarkably, a higher botanical diversity of fruits and vegetables was clearly associated with higher short-chain fatty acids levels, suggesting that this botanical diversity may be needed to fully realize health benefits mediated by the microbiome fermentative activities.

## APPENDIXES

### Appendix 1: SMART-Diet Sample Menu.

HIGH					
Day	Food Item	Serving	Day	Food Item	Serving
1	Pre-Washed Salad Mix	1 Cup	1	Grapes	1/2 Cup
	Frozen Corn-Off Cob	1 Cup		Banana	1 Cup
	Baby Carrots	1 Cup		Blueberries	1/2 Cup
2	Chopped Celery Stalks	1 Cup	2	Peach or Green Apple (Group 2)	1 Cup
	Tomato Cucumber Salad	1/2 Cup		Strawberries	1/2 Cup
	Chicken Vegetable Stir Fry	1 1/2 Cups		Pineapple Chunks	1/2 Cup
3	Broccoli Floret & Cucumber Mix	1 Cup	3	Orange	1 Cup
	Asparagus Spears	1 Cup		Red Apple	1 Cup
	Marinara Sauce (No Meat)	1 Cup			
LOW					
Day	Food Item	Serving	Day	Food Item	Serving
1	Pre-Washed Salad Mix	1 Cup	1	Green Apple	1 Cup
2	Pre-Washed Salad Mix	1 Cup	2	Banana	1 Cup
3	Pre-Washed Salad Mix	1 Cup	3	Orange	1 Cup

### Appendix 2: Low Diversity Cohort Menus.

#### Low Diversity Cohort Menu:

**Lead-in: 6 Total Botanical families:** *Rutaceae* (orange), *Roasaceae* (apple), *Solanaceae* (peppers, potato, tomato), *Asteraceae* (lettuce), *Poaceae* (corn), *Alliaceae* (onion).

**High FV: 11 Total Botanical families:** *Rutaceae* (orange), *Roasaceae* (apple, peaches, pears, berries), *Solanaceae* (peppers, potato, tomato), *Asteraceae* (lettuce), *Poaceae* (corn), *Alliaceae* (onion), *Muscaceae* (banana), *Brassicaceae* (broccoli, cauliflower), *Fabaceae/Leguminosae* (beans, soybeans, chickpeas), *Cucurbitaceae* (cucumber, watermelon, squash), *Apiaceae/Umbelliferae* (carrots, celery).

Condition	Day	Meal	Menu
Lead in	1	Breakfast	1/4 plate (1/2 cup) scrambled eggs, wheat bagel with tablespoon plain cream cheese or butter, 1/4 plate (1/2 cup) breakfast potatoes, 8 oz glass of skim milk
		Lunch	Deli turkey/ham sandwich (3oz meat) on wheat bread (1 slice cheese, mustard or light mayo, lettuce, tomato), apple with 2 tablespoons peanut butter
		Dinner	1/2 plate (2 cups) penne pasta with 1/2 cup red meat sauce with 2 tablespoons of parmesan cheese, 1 breadstick
		Snack/Dessert	Cup of ice cream

**Appendix 2 (Cont.)**

Condition	Day	Meal	Menu
	2	Breakfast	1/2 plate Scrambled eggs with cheese, 2 slices of wheat toast lightly buttered, 8oz glass of orange juice
		Lunch	2 slices of pepperoni pizza, small side salad (romaine lettuce or iceberg lettuce) with tablespoon of dressing, cup of corn chowder soup (1 ½ ladle fulls)
		Dinner	1 piece garlic grilled salmon, ¼ plate (1/2 cup) potato au gratin dinner roll
		Snack/Dessert	Cookie (any kind but oatmeal raisin) with 8oz glass of skim milk
	3	Breakfast	cup of oatmeal (approximately 1 ½ ladle fulls) with ~1 oz milk, 2 tablespoons brown sugar or honey, 2 turkey sausage
		Lunch	3 beef tacos w/ salsa, ½ cup tortilla chips, 2 tablespoons nacho cheese
		Dinner	2 plum glazed chicken thighs, baked potato w/ tablespoon chz and tablespoon sour cream, side salad (romaine or iceberg lettuce) with dressing, 8oz glass of skim milk
		Snack/Dessert	apple with 2 tablespoons peanut butter
	4	Breakfast	2 pancakes/french toast tablespoon butter and syrup, 8oz glass of milk, 2 turkey sausage
		Lunch	Hamburger (no cheese), ¼ (1/2 cup) plate fries with 2 tablespoons of ketchup, orange, ½ cup flavored yogurt
		Dinner	3 slices of cheese pizza, 1/2C bell peppers or side salad (romaine or iceberg lettuce) with tablespoon ranch, apple
		Snack/Dessert	None
High FV	1	Breakfast	¼ plate (1/2 cup) scrambled eggs with cheese, 2 pieces of wheat toast lightly butter, banana
		Lunch	Chicken burrito- (1 whole wheat tortilla, ask for extra grilled veggies, shredded cheese, lettuce, salsa), 1/4 plate (1/2cup) of beans, ¼ plate (1/2 cup) rice
		Dinner	(2 pieces or “boats”) Zucchini stuffed with tomato, and faro, dinner roll, 1 cup of berries (no blueberries) or melon, 8oz glass of skim milk
		Snack/Dessert	2 cookies (any kind but oatmeal raisin)

## Appendix 2 (Cont.)

Condition	Day	Meal	Menu
	2	Breakfast	¼ plate scrambled eggs with cheese, ¼ plate breakfast potatoes, 2 veggie sausage, 2 slices of toast lightly buttered or with peanut butter, 1 cup of melon,
		Lunch	2 black bean tamales with 2 tablespoons salsa, 1 cup (1 ½ ladle fulls) butternut squash bisque, side salad, (no spinach) with dressing, cookie (any kind but oatmeal raisin)
		Dinner	carved roast sirloin (3oz) with tablespoon creamy horseradish, ½ plate (1 cup) steamed fresh vegetables, dinner roll, apple, 8oz glass skim milk
		Snack/Dessert	½ cup or 1 Cup n celery/ carrots/ broccoli/ bell peppers
	3	Breakfast	¼ plate scrambled eggs with salsa, 2 pieces of toast lightly buttered, orange
		Lunch	~ ¼ plate (1 cup) whole wheat pasta with (¼ cup) red meat sauce, and tablespoon of parmesan cheese, ½ plate (1 cup) steamed broccoli, breadstick, ½ cup yogurt with ¼ cup berries, (no blueberries)
		Dinner	tandoori chicken, ¼ plate (½ cup) basmati rice, ¼ plate (½ cup) roasted cauliflower, 8oz glass of milk, side salad, (no spinach) with dressing
		Snack/Dessert	banana, with tablespoon peanut butter
	4	Breakfast	1 cup plain Greek yogurt, ½ cup peaches or berries (no blueberries), 1/2 cup granola, 2 veggie sausages
		Lunch	Falafel, gyro sandwich with tzatiki, ¼ plate (½ cup) quinoa tabbouleh, 1 cup cucumbers, with tablespoon hummus
		Dinner	Cobb Salad, 1 piece of cornbread with tablespoon butter, 8oz glass of chocolate milk, 1 cup watermelon
		Snack/Dessert	½ cup carrots /bell peppers /broccoli /celery

## Appendix 3: High Diversity Cohort Menus.

### High Diversity Cohort Menu:

**Lead-in: 6 Total Botanical families:** *Rutaceae* (orange), *Roasaceae* (apple), *Solanaceae* (peppers, potato, tomato), *Asteraceae* (lettuce), *Poaceae* (corn), *Alliaceae* (onion).

**High FV: 24 Total Botanical Families:** *Rutaceae* (orange), *Roasaceae* (apple/berries/peach/pear), *Solanaceae* (peppers, potato, tomato), *Asteraceae* (lettuce), *Poaceae* (corn), *Alliaceae* (onion), *Muscaceae* (banana), *Brassicaceae* (broccoli, cauliflower), *Fabaceae/Leguminosae* (beans, peas, soybeans), *Cucurbitaceae* (cucumber, melon, squash), *Apiaceae/Umbelliferae* (carrots, celery), *Chenopodiaceae* (spinach), *Actinidaceae* (kiwi), *Ericaceae* (blueberry, cranberry), *Convolvulaceae* (sweet potatoes), *Vitaceae* (grape), *Anacardiaceae* (mango), *Arecaceae* (date, coconut), *Bromeliaceae* (pineapple), *Caricaceae* (papaya), *Asparaaaceae* (asparagus), *Dioscoreaceae* (yam), *Agaricaceae* (mushrooms), *Grossulariaceae* (black currant).

Condition	Day	Meal	Menu
Lead in	1	Breakfast	1/4 plate (1/2 cup) scrambled eggs, wheat bagel with tablespoon low fat plain cream cheese or butter, 1/4 plate (1/2 cup) jalapeno cheddar potato rounds, 8 oz glass of skim milk
		Lunch	Deli turkey/ham sandwich (3oz meat) on wheat bread (1 slice cheese, mustard or light mayo, lettuce, tomato), apple, with 2 tablespoons peanut butter, 1/4 plate (1/2 cup) steamed corn
		Dinner	1/2 plate (2 cups) penne pasta with 1/2 cup meat sauce, and 2 tablespoons of parmesan cheese, 1 breadstick
		Snack/Dessert	Cup of ice cream
	2	Breakfast	1/2 plate (1 cup) Scrambled eggs with cheese, 2 slices of wheat toast lightly buttered, 8oz glass of orange juice
		Lunch	2 slices of pepperoni pizza, small side salad, (romaine or iceberg lettuce) with tablespoon of dressing (no raspberry vinaigrette), 2 focaccia sticks, 1/2 cup (roughly size of tennis ball) greek yogurt drizzled with honey
		Dinner	Tempura chicken with 2 tablespoons honey mustard, 1/4 plate (1/2 cup) mashed potatoes , dinner roll
		Snack/Dessert	Cookie (any kind but oatmeal raisin) with 8oz glass of skim milk
	3	Breakfast	cup of oatmeal (approximately 1.5 ladle fulls) with ~1 oz milk, 2 tablespoons brown sugar, 1 turkey sausage patty
		Lunch	3 tacos w/ red salsa or pico de galo, 1/2 cup (roughly size of tennis ball) tortilla chips, 2 tablespoons nacho cheese
		Dinner	Spicy marinated cod, baked potato, with tablespoon cheese and tablespoon sour cream, side salad, (romaine or iceberg lettuce) with dressing (not raspberry vinaigrette), 8oz glass of skim milk
		Snack/Dessert	Apple with 2tbsp peanut butter
	4	Breakfast	2 cinnamon vanilla pancakes with butter and syrup, 8oz glass of skim milk, 1/4 cup scrambled eggs or hard boiled egg
		Lunch	Hamburger (no cheese), 1/4 plate fries (1/2 cup) with 2 tablespoons of ketchup, orange
		Dinner	2 slices of cheese pizza, 1/2 cup bell peppers or side salad (romaine or iceberg lettuce), with tablespoon ranch, apple
		Snack/Dessert	lowfat vanilla yogurt

**Appendix 3 (Cont.)**

Condition	Day	Meal	Menu
High FV	1	Breakfast	¼ plate (1/2 cup) plate scrambled eggs with cheese, ~15 grapes, 2 veggie sausage, oatmeal yam muffin
		Lunch	Vegetarian burrito bowl: cilantro lime rice, black beans, 2 tablespoons shredded cheese, extra peppers and onions, lettuce, salsa, w/ tropical smoothie (made with 1C frozen papaya & mango, 4oz OJ, 1/4C plain greek yogurt)
		Dinner	Chicken Florentine, ½ plate (1 cup) roasted potatoes and vegetables (sweet potatoes), spinach side salad (include sliced mushrooms), with dressing
		Snack/Dessert	1/2C carrots/celery and ½ cup dried figs, 8oz glass of skim milk
	2	Breakfast	1 cup (1 ½ ladle full) oatmeal with 1oz milk, ¼ cup dried black currants, and 2 tablespoons brown sugar, ½ cup pineapple
		Lunch	Chipotle lime tilapia with ¼ cup chipotle pineapple salsa, ¼ plate (½ cup) Jamaican peas and rice, ¼ plate (½ cup) asparagus
		Dinner	Meatloaf, ¼ plate (1/2 cup) mashed potatoes, ½ plate, (1 cup) sautéed spinach w/ garbanzo beans, 8oz lowfat chocolate milk
		Snack/Dessert	½ cup of mushrooms and banana with 1- 2 tablespoons peanut butter
	3	Breakfast	¼ plate scrambled eggs, bagel with tablespoon cream cheese, 1 cup melon
		Lunch	Salmon with ¼ cup tropical fruit salsa, ¼ plate (½ cup) aztec blend or rice pilaf, ½ plate (1 cup) steamed fresh vegetables
		Dinner	Plate of salad, (include raisins, cheese, and nuts of some sort) cup of vegetable soup of the day, (1 ½ ladles full), 1 piece of cornbread, 8 oz glass of skim milk
		Snack/Dessert	2 kiwi, yam muffin, 2 date coconut bites



**Appendix 3 (Cont.)**

<b>Condition</b>	<b>Day</b>	<b>Meal</b>	<b>Menu</b>
	4	Breakfast	Banana walnut pancakes topped w/ whipped cream, ½ cup peaches/pears/apple/strawberries, 8oz glass of milk
		Lunch	2 fish or chicken tacos with ~ 3 tablespoons mango salsa, ½ cup black beans
		Dinner	¼ plate (1 cup) pasta with marinara sauce, and 2 tablespoons parmesan, ½ plate (1 cup) roasted cauliflower, side salad (romaine lettuce, sliced mushrooms, raisins, any other veg) with tablespoon dressing
		Snack/Dessert	1 cup greek yogurt with 1 cup blueberries and sweet potato chips

#### Appendix 4:



Office of Research Compliance

To: Franck Carbonero  
FR: Craig Coon  
Date: March 10th, 2017  
Subject: IACUC Approval  
Expiration Date: March 9th, 2018

The Institutional Animal Care and Use Committee (IACUC) has APPROVED your protocol # **17057**: *Impact of varying amounts of concentrate tart cherry juice on healthy mice gut microbiota*.

In granting its approval, the IACUC has approved only the information provided. Should there be any further changes to the protocol during the research, please notify the IACUC in writing (via the Modification form) prior to initiating the changes. If the study period is expected to extend beyond March 9th, 2018 you can submit a modification to extend project up to 3 years, or submit a new protocol. By policy the IACUC cannot approve a study for more than 3 years at a time.

The following individuals are approved to work on this study: . Please submit personnel additions to this protocol via the modification form prior to their start of work.

The IACUC appreciates your cooperation in complying with University and Federal guidelines involving animal subjects.

CNC/aem

## Appendix 5:



Office of Research Compliance  
Institutional Review Board

May 9, 2017

### MEMORANDUM

TO: Franck Carbonero  
Ayoub AlOthaim  
Days Marasini

FROM: Ro Windwalker  
IRB Coordinator

RE: New Protocol Approval

IRB Protocol #: 17-04-631

Protocol Title: *Cranberry Juice and its Effects on Vaginal and Gut Microbiota in Postmenopausal Women: A Pilot Cross-Over Trial of a Dietary Preventive for Urinary Tract Infection*

Review Type: ☐ EXEMPT ☒ EXPEDITED ☐ FULL IRB

Approved Project Period: Start Date: 05/08/2017 Expiration Date: 04/27/2018

Your protocol has been approved by the IRB. Protocols are approved for a maximum period of one year. If you wish to continue the project past the approved project period (see above), you must submit a request, using the form *Continuing Review for IRB Approved Projects*, prior to the expiration date. This form is available from the IRB Coordinator or on the Research Compliance website (<https://vpred.uark.edu/units/rscp/index.php>). As a courtesy, you will be sent a reminder two months in advance of that date. However, failure to receive a reminder does not negate your obligation to make the request in sufficient time for review and approval. Federal regulations prohibit retroactive approval of continuation. Failure to receive approval to continue the project prior to the expiration date will result in Termination of the protocol approval. The IRB Coordinator can give you guidance on submission times.

**This protocol has been approved for 40 participants.** If you wish to make any modifications in the approved protocol, including enrolling more than this number, you must seek approval prior to implementing those changes. All modifications should be requested in writing (email is acceptable) and must provide sufficient detail to assess the impact of the change.

If you have questions or need any assistance from the IRB, please contact me at 109 MLKG Building, 5-2208, or [irb@uark.edu](mailto:irb@uark.edu).